

# HOMOTOPY EXPONENTS FOR LARGE $H$ -SPACES

WOJCIECH CHACHÓLSKI, WOLFGANG PITTSCH, JÉRÔME SCHERER,  
AND DON STANLEY

ABSTRACT. We show that  $H$ -spaces with finitely generated cohomology, as an algebra or as an algebra over the Steenrod algebra, have homotopy exponents at all primes. This provides a positive answer to a question of Stanley.

## INTRODUCTION

A simply connected space is elliptic if both its rational homotopy and rational homology are finite. Moore's conjecture, see for example [9], predicts that elliptic complexes have an exponent at any prime  $p$ , meaning that there is a bound on the  $p$ -torsion in the graded group of all homotopy groups. Any finite  $H$ -space is known to be elliptic as it is rationally equivalent to a finite product of (odd dimensional) spheres. Relying on results by James [6] and Toda [11] about the homotopy groups of spheres, the fourth author (re)proved in [10] Long's result that finite  $H$ -spaces have an exponent at any prime [7]. He proved in fact a stronger result which holds for example for  $H$ -spaces for which the mod  $p$  cohomology is finite. He also asked whether this would hold for finitely generated cohomology rings. The aim of this note is to give a positive answer to this question and provide a way larger class of  $H$ -spaces which have homotopy exponents.

**Theorem 0.1.** *1.2 Let  $X$  be a connected and  $p$ -complete  $H$ -space such that  $H^*(X; \mathbb{F}_p)$  is finitely generated as an algebra over the Steenrod algebra. Then  $X$  has an exponent at  $p$ .*

This class of  $H$ -spaces is optimal in the sense that  $H$ -spaces with a larger mod  $p$  cohomology, such as an infinite product of Eilenberg-Mac Lane spaces

---

2000 *Mathematics Subject Classification.* Primary 55P45; Secondary 55Q99, 55S10, 55U20.

The first author is supported in part by NSF grant DMS-0296117, Vetenskapsrådet grant 2001-4296, and Göran Gustafssons Stiftelse. The second and third authors are partially supported by FEDER/MEC grant MTM2007-61545. The second author is supported by the program Ramón y Cajal, MEC, Spain.

$K(\mathbb{Z}/p^n, n)$ , will not have in general an exponent at  $p$ . As a corollary, we obtain the desired result. In fact we obtain the following global theorem.

**Theorem 0.2.** *1.4 Let  $X$  be a connected  $H$ -space such that  $H^*(X; \mathbb{Z})$  is finitely generated as an algebra. Then  $X$  has an exponent at each prime  $p$ .*

The methods we use are based on the deconstruction techniques of the third author in his joint work with Castellana and Crespo, [3]. Our results on homotopy exponents should also be compared with the computations of homological exponents done with Clément, [4]. Whereas such  $H$ -spaces always have homotopy exponents, they almost never have homological exponents. The only simply connected  $H$ -spaces for which the 2-torsion in  $H_*(X; \mathbb{Z})$  has a bound are products of mod 2 finite  $H$ -spaces with copies of the infinite complex projective space  $\mathbb{C}P^\infty$  and  $K(\mathbb{Z}, 3)$ .

#### ACKNOWLEDGMENTS

This project originated during a workshop at the CRM, Barcelona, held in the emphasis year on algebraic topology (2007-08). We would like to thank the organizers for making it possible to meet in such a pleasant atmosphere.

#### 1. HOMOTOPY EXPONENTS

Our starting point is the fact that mod  $p$  finite  $H$ -spaces have always homotopy exponents. The following is a variant of Stanley's [10, Corollary 2.9]. Whereas he focused on spaces localized at a prime, we will stick to  $p$ -completion in the sense of Bousfield and Kan, [2]. Since the  $p$ -localization map  $X \rightarrow X_{(p)}$  is a mod  $p$  homology equivalence, his result implies the following.

**Proposition 1.1** (Stanley). *Let  $p$  be a prime and  $X$  be a  $p$ -complete and connected  $H$ -space such that  $H^*(X; \mathbb{F}_p)$  is finite. Then  $X$  has an exponent at  $p$ .*

We will not repeat the proof, but let us sketch the main steps. Let us consider a decomposition of  $X$  by  $p$ -complete cells, i.e.  $X$  is obtained by attaching cones along maps from  $(S^n)_p^\wedge$ . The natural map  $X \rightarrow \Omega\Sigma X$  factors then through the loop spaces on a wedge  $W$  of a finite numbers of such  $p$ -completed spheres, up to multiplying by some integer  $N$ : the composite  $X \rightarrow \Omega\Sigma X \xrightarrow{N} \Omega\Sigma X$  is homotopic to  $X \rightarrow \Omega W \rightarrow \Omega\Sigma X$ . The proof goes by induction on the number of  $p$ -complete cells and the key ingredient here is Hilton's description of the loop space on a wedge of spheres, [5]. Note that the suspension of a map between spheres is torsion except for the multiples of the identity. This idea to "split off" all the cells

of  $X$  up to multiplication by some integer is dual to Arlettaz' way to split off Eilenberg-Mac Lane spaces in  $H$ -spaces with finite order  $k$ -invariants, [1, Section 7]. The final step relies on the classical results by James, [6], and Toda, [11], that spheres do have homotopy exponents at all primes.

**Theorem 1.2.** *Let  $X$  be a connected and  $p$ -complete  $H$ -space such that  $H^*(X; \mathbb{F}_p)$  is finitely generated as an algebra over the Steenrod algebra. Then  $X$  has an exponent at  $p$ .*

*Proof.* A connected  $H$ -space such that  $H^*(X; \mathbb{F}_p)$  is finitely generated as an algebra over the Steenrod algebra can always be seen as the total space of an  $H$ -fibration  $F \rightarrow X \rightarrow Y$  where  $Y$  is an  $H$ -space with finite mod  $p$  cohomology and  $F$  is a  $p$ -torsion Postnikov piece whose homotopy groups are finite direct sums of copies of cyclic groups  $\mathbb{Z}/p^r$  and Prüfer groups  $\mathbb{Z}_{p^\infty}$ , [3, Theorem 7.3]. This is a fibration of  $H$ -spaces and  $H$ -maps, so that we obtain another fibration  $F_p^\wedge \rightarrow X \rightarrow Y_p^\wedge$  by  $p$ -completing it. The base space  $Y_p^\wedge$  now satisfies the assumptions of Proposition 1.1. It has therefore an exponent at  $p$ . The homotopy groups of the fiber  $F_p^\wedge$  are finite direct sums of cyclic groups  $\mathbb{Z}/p^n$  and copies of the  $p$ -adic integers  $\mathbb{Z}_p^\wedge$ . Thus  $F_p^\wedge$  has an exponent at  $p$  as well. The homotopy long exact sequence of the fibration allows us to conclude.  $\square$

We see here how the  $p$ -completeness assumption plays an important role. The space  $K(\mathbb{Z}_{p^\infty}, 1)$  for example has obviously no exponent at  $p$ , but its  $p$ -completion is  $K(\mathbb{Z}_p^\wedge, 2) = (\mathbb{C}P^\infty)_p^\wedge$ , which is a torsion free space. The mod  $p$  cohomology of  $K(\mathbb{Z}_{p^\infty}, 1)$  is a polynomial ring on one generator in degree 2, we must thus also work with  $p$ -complete spaces to give an answer to Stanley's question [10, Question 2.10].

**Corollary 1.3.** *Let  $X$  be a connected and  $p$ -complete  $H$ -space such that  $H^*(X; \mathbb{F}_p)$  is finitely generated as an algebra. Then  $X$  has an exponent at  $p$ .*

In fact, when the mod  $p$  cohomology is finitely generated, the fiber  $F$  in the fibration described in the proof of Theorem 1.2 is a single Eilenberg-Mac Lane space  $K(P, 1)$ . Thus the typical example of an  $H$ -space with finitely generated mod  $p$  cohomology is the 3-connected cover of a simply connected finite  $H$ -space ( $P$  is  $\mathbb{Z}_{p^\infty}$  in this case). Likewise, the typical example in Theorem 1.2 are highly connected covers of finite  $H$ -spaces. This explains why such spaces have homotopy exponents!

If one does not wish to work at one prime at a time and prefers to find a global condition which permits to conclude that a certain class of spaces have exponents at all primes, one must replace mod  $p$  cohomology by integral cohomology.

**Theorem 1.4.** *Let  $X$  be a connected  $H$ -space such that  $H^*(X; \mathbb{Z})$  is finitely generated as an algebra. Then  $X$  has an exponent at each prime  $p$ .*

*Proof.* Since the integral cohomology groups are finitely generated it follows from the universal coefficient exact sequence (see [8]) that the integral homology groups are also finitely generated. Since  $X$  is an  $H$ -space we may use a standard Serre class argument to conclude that so are the homotopy groups. Therefore the  $p$ -completion map  $X \rightarrow X_p^\wedge$  induces an isomorphism on the  $p$ -torsion at the level of homotopy groups. The theorem is now a direct consequence of the next lemma.  $\square$

**Lemma 1.5.** *Let  $X$  be a connected space. If  $H^*(X; \mathbb{Z})$  is finitely generated as an algebra, then so is  $H^*(X; \mathbb{F}_p)$ .*

*Proof.* Let  $u_1, \dots, u_r$  generate  $H^*(X; \mathbb{Z})$  as an algebra. Consider the universal coefficients short exact sequences

$$0 \rightarrow H^n(X; \mathbb{Z}) \otimes \mathbb{Z}/p \rightarrow H^n(X; \mathbb{F}_p) \xrightarrow{\partial} \text{Tor}(H^{n+1}(X; \mathbb{Z}); \mathbb{Z}/p) \rightarrow 0.$$

Since  $H^*(X; \mathbb{Z})$  is finitely generated as an algebra it is degree-wise finitely generated as a group and therefore  $\text{Tor}(H^*(X; \mathbb{Z}); \mathbb{Z}/p)$  can be identified with the ideal of elements of order  $p$  in  $H^*(X; \mathbb{Z})$ . This ideal must be finitely generated since  $H^*(X; \mathbb{Z})$  is Noetherian. Choose generators  $a_1, \dots, a_s$ . Each  $a_i$  corresponds to a pair  $\alpha_i, \beta\alpha_i$  in  $H^*(X; \mathbb{F}_p)$ , where  $\beta$  denotes the Bockstein.

We claim that the elements  $\alpha_1, \dots, \alpha_s$  together with the mod  $p$  reduction of the algebra generators, denoted by  $\bar{u}_1, \dots, \bar{u}_r$ , generate  $H^*(X; \mathbb{F}_p)$  as an algebra. Let  $x \in H^*(X; \mathbb{F}_p)$  and write its image  $\partial(x) = \sum \lambda_j a_j$  with  $\lambda_j = \lambda_j(u)$  a polynomial in the  $u_i$ 's. Define now  $\bar{\lambda}_j = \lambda_j(\bar{u}) \in H^*(X; \mathbb{F}_p)$  to be the corresponding polynomial in the  $\bar{u}_i$ 's. As the action of  $H^*(X; \mathbb{Z})$  on the ideal  $\text{Tor}(H^*(X; \mathbb{Z}); \mathbb{Z}/p)$  factors through the mod  $p$  reduction map  $H^*(X; \mathbb{Z}) \rightarrow H^*(X; \mathbb{F}_p)$ , the element  $x - \sum \bar{\lambda}_j \alpha_j$  belongs to the kernel of  $\partial$ , i.e. it lives in the image of the mod  $p$  reduction. It can be written therefore as a polynomial  $\bar{\mu}$  in the  $\bar{u}_i$ 's. Thus  $x = \bar{\mu} + \sum \bar{\lambda}_j \alpha_j$ .  $\square$

#### REFERENCES

- [1] D. Arlettaz, *Algebraic K-theory of rings from a topological viewpoint*, Publ. Mat. **44(1,3)** (2000), 3–84.
- [2] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Springer-Verlag, Berlin, 1972, Lecture Notes in Mathematics, Vol. 304.
- [3] N. Castellana, J. A. Crespo, and J. Scherer, *Deconstructing Hopf spaces*, Invent. Math. **167(1)** (2007), 1–18.
- [4] A. Clément and J. Scherer, *Homology exponents for  $h$ -spaces*, Rev. Iberoam. Mat. (2009).

- [5] P. J. Hilton, *On the homotopy groups of the union of spheres*, J. London Math. Soc. **30** (1955), 154–172.
- [6] I. M. James, *The suspension triad of a sphere*, Ann. of Math. (2) **63** (1956), 407–429.
- [7] J. Long, *Thesis, Princeton university*, Ph.D. thesis, 1978.
- [8] W. J. R. Mitchell, *Dual modules*, Math. Proc. Cambridge Philos. Soc. **84**(1) (1978), 21–24.
- [9] P. Selick, *Moore conjectures*, Algebraic topology—rational homotopy (Louvain-la-Neuve, 1986), Lecture Notes in Math., vol. 1318, Springer, Berlin, 1988, pp. 219–227.
- [10] D. Stanley, *Exponents and suspension*, Math. Proc. Cambridge Philos. Soc. **133**(1) (2002), 109–116.
- [11] H. Toda, *On the double suspension  $E^2$* , J. Inst. Polytech. Osaka City Univ. Ser. A. **7** (1956), 103–145.

WOJCIECH CHACHÓLSKI  
K.T.H. MATEMATIK  
S-10044 STOCKHOLM, SWEDEN  
*E-mail address:* `wojtek@math.kth.se`

WOLFGANG PITSCH  
UNIVERSITAT AUTÒNOMA DE BARCELONA  
DEPARTAMENT DE MATEMÀTIQUES  
E-08193 BELLATERRA, SPAIN  
*E-mail address:* `pitsch@mat.uab.es`

JÉRÔME SCHERER  
UNIVERSITAT AUTÒNOMA DE BARCELONA  
DEPARTAMENT DE MATEMÀTIQUES  
E-08193 BELLATERRA, SPAIN  
*E-mail address:* `jscherer@mat.uab.es`

DON STANLEY  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
UNIVERSITY OF REGINA, COLLEGE WEST 307.14  
REGINA, SASKATCHEWAN, CANADA S4S 0A2  
*E-mail address:* `Donald.Stanley@uregina.ca`