

EMBEDDINGS OF DI_2 IN F_4

CARLES BROTO AND JESPER M. MØLLER

ABSTRACT. We show that there is only one embedding of BDI_2 in BF_4 at the prime $p = 3$, up to self maps of BDI_2 . We also describe the effect of the group of self-equivalences of BF_4 at the prime $p = 3$ on this embedding and then show that the Friedlander's exceptional isogeny composed with a suitable Adams map is an involution of BF_4 whose homotopy fixed point set coincide with BDI_2

1. INTRODUCTION

A p -compact group is a p -complete analogue in homotopy theory of the concept of compact Lie group. The notion was introduced by Dwyer and Wilkerson in [12]. A p -compact group is a p -complete space BX whose loop space is \mathbf{F}_p -finite, i.e. has finite mod p cohomology. The p -completion of the classifying space of a compact connected Lie group is an example of a p -compact group but there are other, exotic, examples. Among other notions that p -compact groups share with compact Lie groups are that of a maximal torus, Weyl group, and maximal torus normalizer (see [12] for the definitions). There is also a well defined concept of homomorphism or monomorphism between p -compact groups. Since the introduction of this concept one main issue has been the description of the exotic examples. These are simply connected and irreducible p -compact groups that do not appear as the p -completion of a compact Lie group. Contrary to what happens with compact Lie groups these exotic examples can only be defined as p -complete spaces at a certain given prime p . We will focus our attention on one of these examples, namely, BDI_2 . This is a 3-compact group with mod 3 cohomology ring

$$(1.1) \quad H^*(BDI_2; \mathbf{F}_3) \cong P[x_{12}, x_{16}], \quad P^1 x_{12} = x_{16},$$

the rank two Dickson algebra at the prime 3.

Date: February 24, 1999.

1991 Mathematics Subject Classification. 55R35, 55P15, 55P10.

Key words and phrases. Homotopy Lie groups, Dickson algebra.

C. Broto is partially supported by DGES grant 97-0203.

Theorem 1.2. *Any non-trivial homomorphism from DI_2 to F_4 is a monomorphism and the group of automorphisms of DI_2 acts simply transitively on the non-empty set of monomorphisms from DI_2 to F_4 .*

There is then a monomorphism $\mathrm{B}\alpha: \mathrm{BDI}_2 \rightarrow \mathrm{BF}_4$ and it is unique up to composition with automorphisms of DI_2 . We can rephrase it by saying that F_4 contains a unique copy of DI_2 .

If we compose the Friedlander's exceptional isogeny of F_4 ,

$$\mathrm{B}\varphi: \mathrm{BF}_4 \rightarrow \mathrm{BF}_4$$

[14], with the Adams map $\psi^{1/\zeta}$, where $\zeta^2 = -2$, then $\mathrm{B}\tau = \mathrm{B}\psi^{1/\zeta} \circ \mathrm{B}\varphi$ is a homotopy involution of BF_4 that fixes $\mathrm{B}\alpha: \mathrm{BDI}_2 \rightarrow \mathrm{BF}_4$ up to homotopy.

Theorem 1.3. *There is an action of $\mathbf{Z}/2$ on BF_4 induced up to homotopy by $\mathrm{B}\tau$, for which $\mathrm{B}\alpha: \mathrm{BDI}_2 \rightarrow \mathrm{BF}_4$ induces a homotopy equivalence*

$$\mathrm{BDI}_2 \simeq (\mathrm{BF}_4)^{h\mathbf{Z}/2}$$

that identifies BDI_2 with a homotopy fix point set of BF_4 .

Notice that in particular the Bousfield-Kan spectral sequence for the homotopy groups of $(\mathrm{BF}_4)^{h\mathbf{Z}/2}$ degenerates to

$$(1.4) \quad \pi_*(\mathrm{BDI}_2) \cong \pi_*(\mathrm{BF}_4)^{\mathbf{Z}/2}.$$

As a further corollary we obtain

Corollary 1.5. *For any 3-complete space X ,*

$$\mathrm{map}(X, \mathrm{BDI}_2) \simeq \mathrm{map}(X, \mathrm{BF}_4)^{h\mathbf{Z}/2}.$$

And applying this result to the homogeneous space $\mathrm{F}_4/\mathrm{DI}_2$,

Corollary 1.6 (Harper [15]). *At the prime 3 there is a splitting of H -spaces*

$$\mathrm{F}_4 \simeq \mathrm{DI}_2 \times \mathrm{F}_4/\mathrm{DI}_2.$$

Some historical considerations about the construction of this space will clarify our motivation. The way exotic examples are obtained is as realizations of graded polynomial algebras over \mathbf{F}_p as mod p cohomology rings of p -completed spaces BX which will turn out to be p -compact groups. Steenrod [31] already pointed out the importance of realizing polynomial algebras as cohomology rings. The idea was reconducted by Clark-Ewing [8] and Wilkerson [34] who noticed the relevance of invariant theory to the realization problem.

Dickson algebras at odd primes are invariant algebras $D(n) \cong \mathbf{F}_p[t_1, \dots, t_n]^{\mathrm{GL}_n(\mathbf{F}_p)}$ by the action of the full general linear group of n by n matrices over \mathbf{F}_p on a polynomial algebra on n independent generators

of degree 2. They are shown to be again polynomial algebras of n independent generators of degrees $2(p^n - p^i)$ [10, 28, 33]. In [28] the authors already proved that Dickson algebras at odd primes can only appear as a cohomology ring if $n = 1$, in which case we obtain the cohomology of a p -local sphere of dimension $2(p - 1)$ or in case $n = 2$ and $p = 3$, when $D(2) = \mathbf{F}_3[t_1, t_2]^{\mathrm{GL}_2(\mathbf{F}_3)} \cong \mathbf{F}_3[x_{12}, x_{16}]$ is the algebra of equation (1.1).

A space BDI_2 realizing $D(2)$ as mod 3 cohomology ring was first obtained by Zabrodsky in [36]. His method was technical and quite involved. It consisted in realizing geometrically the 3-local homotopy classes of BF_4 , the classifying space of the exceptional Lie group F_4 , that remain invariant under the action of the Friedlander's exceptional isogeny. A more intelligible construction was provided by Aguadé [2]. The structure of BDI_2 as a 3-compact group was exposed in [23]. However one important feature of Zabrodsky's construction remained unexplained in these new approaches. Zabrodsky's construction came together with a map $\mathrm{BDI}_2 \rightarrow \mathrm{BF}_4$, after 3-completion, turning $H^*(\mathrm{BDI}_2; \mathbf{F}_3)$ into a finitely generated $H^*(\mathrm{BF}_4; \mathbf{F}_3)$ -module, thus representing a monomorphism of 3-compact groups. The original motivation of this note was an attempt to explain not the methods of Zabrodsky but the fact that BDI_2 was obtained as a sort of invariant gadget in BF_4 . This is achieved in Theorem 1.2 and equation (1.4).

The paper is organized as follows. Sections 2 and 3 are devoted to a careful investigation of p -compact group homomorphisms from DI_2 to F_4 . This leads to the proof of Theorem 1.2. In Section 4 we compute the cohomological structure of the homogeneous space F_4/DI_2 . It turns out to coincide with that of the Harper's molecule at the prime three, and this, in turn implies that both spaces have the same homotopy type. Details are included in an Appendix for convenience of the reader. Sections 5 and 6 are devoted to the Friedlander's exceptional isogeny of $(BF_4)_{1/2}$. It is shown that suitably composed with an Adams map becomes an involution that fixes BDI_2 . The effect of these maps in cohomological generators is determined and this leads to the proof of Theorem 1.3 and corollaries 1.5 and 1.6. Some straightforward calculations, mainly in Section two, were done with the aid of a computer using *magma* system. It should be easy to perform the same calculations with any other suitable program.

The authors are grateful to Bill Dwyer for a helpful discussion that led to the final form of Theorem 1.3, and also to Kasper Andersen for his help with the *magma* system. We kindly acknowledge the support of the Centre de Recerca Matemàtica of the Institut d'Estudis Catalans and the Matematisk Institut of the Københavns Universitet that put the necessary means for the authors to meet together.

2. ADMISSIBLE HOMOMORPHISMS

Let X_1 and X_2 be p -compact groups [12] with maximal tori $i_1: T_1 = T(X_1) \rightarrow X_1$, $i_2: T_2 = T(X_2) \rightarrow X_2$, and corresponding Weyl groups $W_1 = W(X_1)$, $W_2 = W(X_2)$, respectively. Write $\text{Rep}(X_1, X_2) = [\text{BX}_1, \text{BX}_2]$ for the set of conjugacy classes of morphisms $X_1 \rightarrow X_2$. The restriction map

$$\text{Rep}(X_1, X_2) \xrightarrow{\overline{i_1}} \text{Rep}(T_1, X_2) \xleftarrow[\cong]{i_2} W_2 \backslash \text{Rep}(T_1, T_2)$$

where $\overline{i_1}$ and the bijection i_2 [22, 3.4.(1)] are induced by i_1 and i_2 , takes $f \in \text{Rep}(X_1, X_2)$ to $f|_{T_1} = W_2 T(f)$ where $T(f)$ is any lift

$$(2.1) \quad \begin{array}{ccc} T_1 & \xrightarrow{T(f)} & T_2 \\ i_1 \downarrow & & \downarrow i_2 \\ X_1 & \xrightarrow{f} & X_2 \end{array}$$

of f to a morphism between the maximal tori.

It is sometimes useful to know that certain morphisms $T_1 \rightarrow T_2$ are *not* of the form $T(f)$ for any $f: X_1 \rightarrow X_2$.

Lemma 2.2. (Cf. [18, 1.8]) *Let $f: X_1 \rightarrow X_2$ be a p -compact group morphism where p is odd and X_1 is connected. Assume that*

- $\pi_1(T(f))$ is injective, and
- p divides the order of the Weyl group W_1 .

Then p does not divide $\pi_1(T(f))$ in $\text{Hom}(\pi_1(T_1), \pi_1(T_2))$.

Proof. By fixed point theory [13, 2.10, 2.14], f lifts to a morphism $N_p(f): \text{Syl}_p(N_1) \rightarrow \text{Syl}_p(N_2)$ of the p -normalizers. The assumption that $\pi_1(T(f))$ be injective implies, since W_1 is faithfully represented in $\pi_1(T_1)$ [12, 9.7], that $\pi_0(N_p(f))$ embeds the Sylow p -subgroup of W_1 into W_2 .

Choose a monomorphism $\mu: \mathbf{Z}/p \rightarrow \text{Syl}_p(N_1)$ such that also $\pi_0(\mu): \mathbf{Z}/p \rightarrow \text{Syl}_p(W_1)$ is injective. This is possible since the epimorphism $\text{Syl}_p(N_1) \rightarrow \text{Syl}_p(W_1)$ admits a section when p is odd [3]. Note that the composition $N_p(f)\mu$ is a monomorphism since it induces a monomorphism on component groups. Consider now the commutative

diagram

$$\begin{array}{ccccc}
 & & \text{Syl}_p(N_1) & \xrightarrow{N_p(f)} & \text{Syl}_p(N_2) \\
 & \nearrow \mu & \downarrow j_p & & \downarrow \\
 \mathbf{Z}/p & \xrightarrow{j_p \mu} & X_1 & \xrightarrow{f} & X_2 \\
 & \searrow \mu' & \uparrow & & \uparrow i_2 \\
 & & T_1 & \xrightarrow{T(f)} & T_2
 \end{array}$$

where μ' is a lift of $j_p \mu$ [12, 4.7, 5.6]. Since $N_p(f)\mu$ is monomorphic, so is $i_2 T(f)\mu'$ by commutativity of the diagram. However, this map would be trivial were $\pi_1(T(f))$ divisible by p . \square

This lemma is also true for $p = 2$ provided the extra assumption

- there exists a monomorphism $\mathbf{Z}/2 \rightarrow \text{Syl}_2(W_1)$ that factors through $\text{Syl}_2(N_1)$

is added.

We now return to the general situation of (2.1). For any element w_1 of the Weyl group W_1 of X_1 ,

$$i_2(T(f) \circ w_1) = i_2 \circ T(f) \circ w_1 = f \circ i_1 \circ w_1 = f \circ i_1 = i_2 \circ T(f) = i_2(T(f))$$

so there must exist some Weyl group element $w_2 \in W_2$ such that $T(f) \circ w_1 = w_2 \circ T(f)$. Thus the lift $T(f)$ lies in the subset $W_2 \setminus \text{Adm}(T_1, T_2)$ where

Definition 2.3. $\text{Adm}(T_1, T_2) = \{a \in \text{Rep}(T_1, T_2) \mid a \cdot W_1 \subseteq W_2 \cdot a\}$

The fundamental group functor can be used to identify this set $\text{Adm}(T_1, T_2)$ of *admissible representations* with the corresponding set $\text{Adm}(\pi_1(T_1), \pi_1(T_2)) \subseteq \text{Hom}(\pi_1(T_1), \pi_1(T_2))$ of *admissible homomorphisms*, i.e. homomorphisms $a: \pi_1(T_1) \rightarrow \pi_1(T_2)$ such that $a \cdot W_1 \subseteq W_2 \cdot a$.

We shall now determine the admissible homomorphisms in the particular case where $p = 3$, $X_1 = \text{DI}_2$ [23, 2], and $X_2 = \text{F}_4$ viewed as a 3-compact group.

The construction of BDI_2 exhibits monomorphisms [23]

$$(2.4) \quad T_1 \rightarrow \text{SU}(3) \rightarrow \text{DI}_2$$

of maximal rank where the morphism $T_1 = T(\text{SU}(3)) \rightarrow \text{SU}(3)$ is a maximal torus for $\text{SU}(3)$. Take the composite morphism (2.4) as the maximal torus for DI_2 .

Let $i_2: T_2 = T(\text{F}_4) \rightarrow \text{F}_4$ be a maximal torus for the simple compact Lie group F_4 (defining a maximal torus for the corresponding 3-compact group).

There is a standard identification

$$\pi_1(T_1) \xrightarrow{\cong} \Sigma_0(\mathbf{Z}_3^3) = \{(x_1, x_2, x_3) \in \mathbf{Z}_3^3 \mid x_1 + x_2 + x_3 = 0\}$$

and we shall use $\{(1, -1, 0), (0, 1, -1)\}$ as a basis for the free \mathbf{Z}_3 -module $\Sigma_0(\mathbf{Z}_3^3)$ thus identifying $\Sigma_0(\mathbf{Z}_3^3)$ with \mathbf{Z}_3^2 . Under this identification we may assume [30, 7.3, Example 1] [23] that the Weyl group $W_1 = W(\mathrm{DI}_2)$ of DI_2 corresponds to the subgroup

$$W(\mathrm{DI}_2) = \langle \alpha, \sigma, \tau \rangle \subseteq \mathrm{Aut}(\Sigma_0(\mathbf{Z}_3^3)) = \mathrm{GL}(2, \mathbf{Z}_3)$$

with generators

$$\alpha = \frac{1}{1+\zeta} \begin{pmatrix} -1+\zeta & -\zeta \\ \zeta & 1-\zeta \end{pmatrix}, \quad \sigma = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \quad \tau = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

where $\zeta^2 + 2 = 0$ and $\zeta \equiv 1 \pmod{3}$. The subgroup $W(\mathrm{SU}(3)) = \langle \sigma, \tau \rangle \cong \Sigma_3$ is the Weyl group of $\mathrm{SU}(3)$.

Similarly, there is a standard identification [6] between the fundamental group $\pi_1(T_2)$ of the maximal torus for F_4 and the free abelian group

$$\Sigma_2(\mathbf{Z}^4) = \{(x_1, x_2, x_3, x_4) \in \mathbf{Z}^4 \mid x_1 + x_2 + x_3 + x_4 \in 2\mathbf{Z}\}$$

Under this identification, the Weyl group $W_2 = W(F_4)$ of F_4 is carried to the group (of order $1152 = 384 \cdot 3$)

$$(2.5) \quad W(F_4) = W(\mathrm{B}_4)E \cup W(\mathrm{B}_4)H_1 \cup W(\mathrm{B}_4)H_2$$

where $W(\mathrm{B}_4)$ is the reflection group (of order $384 = 2^4 \cdot 4!$) of all signed permutation matrices, and H_1 and H_2 are the matrices

$$H_1 = \frac{1}{2} \begin{pmatrix} -1 & 1 & 1 & 1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \\ 1 & 1 & 1 & -1 \end{pmatrix}, \quad H_2 = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}$$

Note that the group $W(\mathrm{B}_4)$ and the matrices H_1 and H_2 preserve the submodule $\Sigma_2(\mathbf{Z}^4)$ of \mathbf{Z}^4 . The operators H_1 and H_2 satisfy the relations $H_1^2 = E = H_2^2$, $H_2H_1 = -H_2$, $H_1H_2 = \mathrm{diag}(-1, 1, 1, 1)H_1$. For F_4 viewed as a 3-compact group with maximal torus T_2 , extensions of scalars provides an identification

$$\pi_1(T_2) \xrightarrow{\cong} \Sigma_2(\mathbf{Z}^4) \otimes \mathbf{Z}_3 \xrightarrow{\cong} \mathbf{Z}^4 \otimes \mathbf{Z}_3 \cong \mathbf{Z}_3^4$$

taking the Weyl group $W(F_4)$ onto the reflection group $W(F_4) < \mathrm{GL}(4, \mathbf{Z}_3)$ as defined in (2.5).

The linear map $A(v): \Sigma_0(\mathbf{Z}_3^3) \rightarrow \mathbf{Z}_3^4$ with matrix

$$(2.6) \quad A(v) = \begin{pmatrix} -\zeta & 1 \\ \zeta & 1 - \zeta \\ 0 & 1 + \zeta \\ -2 & 1 \end{pmatrix}, \quad v \in \mathbf{Z}_3,$$

is admissible with respect to $W(\text{DI}_2)$ and $W(\text{F}_4)$ since $A(v)w = \chi(w)A(v)$ for all $w \in W(\text{DI}_2)$ where $\chi: W(\text{DI}_2) \rightarrow W(\text{F}_4)$ is the group homomorphism with the values

$$\begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad \frac{1}{2} \begin{pmatrix} -1 & -1 & 1 & -1 \\ 1 & -1 & -1 & -1 \\ -1 & 1 & -1 & -1 \\ 1 & 1 & 1 & -1 \end{pmatrix}, \quad \frac{1}{2} \begin{pmatrix} -1 & -1 & 1 & 1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & -1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

on the generators α, σ , and τ , respectively.

Note that $A(v)$ and $-A(v)$ lie in the same orbit under the action of $W(\text{F}_4)$ as $-E \in W(\text{F}_4)$.

Lemma 2.7. *Let v be a 3-adic integer.*

1. *The linear homomorphism $A: \Sigma_0(\mathbf{Z}_3^3) \rightarrow \mathbf{Z}_3^4$ is admissible with respect to $W(\text{DI}_2)$ and $W(\text{F}_4)$ if and only if $A \in W(\text{F}_4)A(v)$ for some 3-adic integer $v \in \mathbf{Z}_3$.*
2. *The linear map $A(v)$ is split injective if and only if $v \in \mathbf{Z}_3^*$ is a 3-adic unit.*
3. *The map*

$$\begin{aligned} \{\pm 1\} \backslash \mathbf{Z}_3^* &\rightarrow W(\text{F}_4) \backslash \text{Hom}_{\mathbf{Z}_3}(\Sigma_0(\mathbf{Z}_3^3), \mathbf{Z}_3^4) \\ \pm v &\rightarrow W(\text{F}_4)A(v) \end{aligned}$$

is injective.

The proof, which we omit, is by direct (computer assisted) computation.

3. CONSTRUCTING MAPS OUT OF BDI_2

Let $\mathbf{I} = \mathbf{I}(W(\text{DI}_2), W(\text{SU}(3)))$ be the category with two objects, 0 and 1, and morphism sets $\mathbf{I}(0, 0) = N_{W(\text{DI}_2)}(W(\text{SU}(3))/W(\text{SU}(3))) \cong Z(W(\text{DI}_2)) \cong \mathbf{Z}/2$, $\mathbf{I}(1, 1) = W(\text{DI}_2)$, $\mathbf{I}(0, 1) = W(\text{DI}_2)/W(\text{SU}(3))$, and $\mathbf{I}(1, 0) = \emptyset$.

The space BDI_2 can be constructed [23, 3.5] as the homotopy colimit of the \mathbf{I}^{op} -space

$$(3.1) \quad (\mathbf{Z}/2)^{\text{op}} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \text{BSU}(3) \xleftarrow{W(\text{SU}(3))^{\text{op}} \backslash W(\text{DI}_2)^{\text{op}}} \text{BT} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} W(\text{DI}_2)^{\text{op}}$$

where $\mathbf{Z}/2$ acts on $\mathrm{BSU}(3)$ as $\{\psi^{\pm 1}\}$. We shall use diagram (3.1) in connection with Wojtkowiak obstruction theory [35] to prove existence and uniqueness of certain maps out of BDI_2 .

The set of conjugacy classes of monomorphisms, $\mathrm{Mono}(\mathrm{DI}_2, \mathbb{F}_4)$ of monomorphisms $\mathrm{DI}_2 \hookrightarrow \mathbb{F}_4$ will turn out to be faithfully represented in the set $\mathrm{Mono}(\mathrm{SU}(3), \mathbb{F}_4)$ of conjugacy classes of monomorphisms $\mathrm{SU}(3) \hookrightarrow \mathbb{F}_4$ which we now describe [25]:

There exists a bijection

$$\begin{aligned} \{(u, v) \in (\mathbf{Z}_3)^2 \mid u + v \equiv 1 \pmod{3}\} &\rightarrow \mathrm{Mono}(\mathrm{SU}(3), \mathbb{F}_4) \\ (u, v) &\rightarrow e\psi^{(u, v)} \end{aligned}$$

where $\psi^{(u, v)}$ is the composite morphism

$$\mathrm{SU}(3) \xrightarrow{\Delta} \mathrm{SU}(3) \times \mathrm{SU}(3) \xrightarrow{\psi^u \times \psi^v} \mathrm{SU}(3) \times \mathrm{SU}(3) \rightarrow \mathrm{SU}(3, 3)$$

and $e: \mathrm{SU}(3, 3) \rightarrow \mathbb{F}_4$ is the inclusion, described in [18, 3.3], of $\mathrm{SU}(3, 3) = \mathrm{SU}(3) \times_{\mathbf{Z}/3} \mathrm{SU}(3)$ as a maximal rank subgroup of \mathbb{F}_4 . (The central $\mathbf{Z}/3$ in $\mathrm{SU}(3) \times \mathrm{SU}(3)$ is generated by $(zE, z^{-1}E)$ where $z \neq 1$ is a third root of unity.) Furthermore, the injective restriction map

(3.2)

$$\mathrm{Mono}(\mathrm{SU}(3), \mathbb{F}_4) \rightarrow \mathrm{Mono}(T(\mathrm{SU}(3)), \mathbb{F}_4) = W(\mathbb{F}_4) \backslash \mathrm{Hom}_{\mathbf{Z}_3}(\Sigma_0(\mathbf{Z}_3^3), \mathbf{Z}_3^4)$$

takes the monomorphism $e\psi^{(u, v)}$ to

$$(3.3) \quad W(\mathbb{F}_4) \begin{pmatrix} -u & v \\ u & -u + v \\ 0 & u + v \\ -2v & v \end{pmatrix}$$

Lemma 3.4. *Let $f \in \mathrm{Mono}(\mathrm{DI}_2, \mathrm{SU}(3))$ be a monomorphism. Then the restriction to $\mathrm{SU}(3)$,*

$$f|_{\mathrm{SU}(3)} = e\psi^{(\zeta v, v)}$$

for a uniquely determined 3-adic unit $v \in \mathbf{Z}_3^*$, $v \equiv -1 \pmod{3}$.

Proof. According to (2.7), the restriction of f to the maximal torus,

$$f|_{T(\mathrm{DI}_2)} \in \mathrm{Mono}(T(\mathrm{DI}_2), \mathbb{F}_4) = W(\mathbb{F}_4) \backslash \mathrm{Hom}_{\mathbf{Z}_3}(\Sigma_0(\mathbf{Z}_3^3), \mathbf{Z}_3^4)$$

is of the form $f|_{T(\mathrm{DI}_2)} = W(\mathbb{F}_4)A(v)$ for some 3-adic unit $v \in \mathbf{Z}_3^*$, uniquely determined up to sign. (The 3-adic number v must be a unit because $A(v)$ is split injective as $f|_{T(\mathrm{DI}_2)}$ is a monomorphism [26] [24].) But this means that the restriction $f|_{\mathrm{SU}(3)} = e\psi^{(\zeta v, v)}$ for a uniquely determined 3-adic unit $v \in \mathbf{Z}_3^*$, $v \equiv -1 \pmod{3}$. \square

Let now $v \in \mathbf{Z}_3^*$ be any 3-adic unit such that $v \equiv -1 \pmod{3}$. Then the two homotopy classes $e\psi^{(\zeta v, v)} \in [\mathbf{BSU}(3), \mathbf{BF}_4]$ and $W(\mathbf{F}_4)A(v) \in [\mathbf{BT}(\mathbf{SU}(3)), \mathbf{BF}_4]$ form a homotopy coherent pair of maps out of the \mathbf{I}^{op} -space (3.1) in the sense that

- $e\psi^{(\zeta v, v)}|_{T(\mathbf{SU}(3))} = W(\mathbf{F}_4)A(v)$
- $W(\mathbf{F}_4)A(v)$ is $W(\mathbf{DI}_2)$ -invariant
- $e\psi^{(\zeta v, v)}$ is $\langle \psi^{-1} \rangle$ -invariant

where the last property follows from the computation

$$e\psi^{(\zeta v, v)}\psi^{-1} = e(\psi^{-1} \times \psi^{-1})\psi^{(\zeta v, v)} = e\psi^{(\zeta v, v)}$$

which uses the identity $e(\psi^{-1} \times \psi^{-1}) = e$ from [18, 3.3]. Therefore, there is an induced \mathbf{I} -space

(3.5)

$$\mathbf{Z}/2 \circlearrowleft \mathbf{BC}_{\mathbf{F}_4}(e\psi^{(\zeta v, v)}) \xrightarrow{W(\mathbf{DI}_2)/W(\mathbf{SU}(3))} \mathbf{BC}_{\mathbf{F}_4}(W(\mathbf{F}_4)A(v)) \circlearrowright_{W(\mathbf{DI}_2)}$$

of connected mapping spaces where

$$\mathbf{BC}_{\mathbf{F}_4}(e\psi^{(\zeta v, v)}) = \text{map}(\mathbf{BSU}(3), \mathbf{BF}_4)_{\mathbf{B}e\psi^{(\zeta v, v)}} \simeq \mathbf{BZ}(\mathbf{SU}(3))$$

$$\mathbf{BC}_{\mathbf{F}_4}(W(\mathbf{F}_4)A(v)) = \text{map}(\mathbf{BT}(\mathbf{SU}(3)), \mathbf{BF}_4)_{\mathbf{B}(W(\mathbf{F}_4)A(v))} \simeq \mathbf{BT}(\mathbf{F}_4)$$

as $e\psi^{(\zeta v, v)}$ is centric and p -toric [25]. These two spaces are simple so we may apply the homotopy functor π_t to (3.5) to obtain the \mathbf{I} -module $\underline{\pi}_t(v)$.

Lemma 3.6. $\lim_{\mathbf{I}}^{-s} \underline{\pi}_t(v) = 0$ for all $s \leq 0$, $t \geq 0$, $s + t \geq -1$ and for all 3-adic units $v \in \mathbf{Z}_3^*$ with $v \equiv -1 \pmod{3}$.

Proof. For the \mathbf{I} -module $\underline{\pi}_1(v)$,

$$\mathbf{Z}/2 \circlearrowleft \mathbf{Z}/3 \xrightarrow{W(\mathbf{DI}_2)/W(\mathbf{SU}(3))} 0 \circlearrowright_{W(\mathbf{DI}_2)}$$

we have $\lim_{\mathbf{I}}^{-s} \underline{\pi}_1(v) = H^{-s}(\mathbf{Z}/2; \mathbf{Z}/3)$ which is trivial since $\mathbf{Z}/2 = \langle \psi^{-1} \rangle$ acts non-trivially on the center $\mathbf{Z}/3 = Z(\mathbf{SU}(3))$. For the \mathbf{I} -module $\underline{\pi}_2(v)$,

$$\mathbf{Z}/2 \circlearrowleft 0 \xrightarrow{W(\mathbf{DI}_2)/W(\mathbf{SU}(3))} \mathbf{Z}_3^4 \circlearrowright_{W(\mathbf{DI}_2)}$$

we have $\lim_{\mathbf{I}}^0 \underline{\pi}_2(v) = 0$, $\lim_{\mathbf{I}}^1 \underline{\pi}_2(v) = (\mathbf{Z}_3^4)^{Z(W(\mathbf{DI}_2) \times W(\mathbf{SU}(3)))} = 0$ as $-E$ belongs to the center $Z(W(\mathbf{DI}_2))$, and $\lim_{\mathbf{I}}^{-s} \underline{\pi}_2(v) = 0$ for $s \leq -2$ by [24, 3.8]. \square

With this lemma in place we are ready for one of the main results of this paper.

Theorem 3.7. *The following hold for the set $\text{Mono}(\text{DI}_2, \mathbb{F}_4)$ of conjugacy classes of monomorphisms $\text{DI}_2 \rightarrow \mathbb{F}_4$:*

1. *The restriction map*

$$\text{Mono}(\text{DI}_2, \mathbb{F}_4) \rightarrow \text{Mono}(\text{SU}(3), \mathbb{F}_4)$$

is an injection with $\{e\psi^{(\zeta v, v)} \mid v \in \mathbf{Z}_3^, v \equiv -1 \pmod{3}\}$ as its image.*

2. *The restriction map*

$$\text{Mono}(\text{DI}_2, \mathbb{F}_4) \rightarrow \text{Mono}(T(\text{DI}_2), \mathbb{F}_4)$$

is an injection with $\{W(\mathbb{F}_4)A(v) \mid v \in \mathbf{Z}_3^, v \equiv -1 \pmod{3}\}$ as its image.*

Proof. 1. This follows from Wojtkowiak's obstruction theory [35] as $\lim_{\mathbf{I}}^{-s} \underline{\pi}_t(v) = 0$ when $s + t = 0$ and $s + t = -1$.

2. The map is a composition

$$\text{Mono}(\text{DI}_2, \mathbb{F}_4) \rightarrow \text{Mono}(\text{SU}(3), \mathbb{F}_4) \rightarrow \text{Mono}(T(\text{SU}(3)), \mathbb{F}_4)$$

of two injections (3.2). □

Let $f(v) : \text{DI}_2 \rightarrow \mathbb{F}_4$, $v \in \mathbf{Z}_3^*$, $v \equiv -1 \pmod{3}$, denote the unique extension to DI_2 of the monomorphism $e\psi^{(\zeta v, v)} : \text{SU}(3) \rightarrow \mathbb{F}_4$ on $\text{SU}(3)$.

- Corollary 3.8.**
1. *The group $\text{Out}(\text{DI}_2)$ acts simply transitively on the set $\text{Mono}(\text{DI}_2, \mathbb{F}_4)$ of monomorphisms.*
 2. *The centralizer of any monomorphism $\text{DI}_2 \rightarrow \mathbb{F}_4$ is trivial.*
 3. *The Weyl group [23] of any monomorphism $\text{DI}_2 \rightarrow \mathbb{F}_4$ is trivial.*

Proof. 1. This is clear since [23] $\text{Out}(\text{DI}_2) \cong \{v \in \mathbf{Z}_3^* \mid v \equiv 1 \pmod{3}\}$ and

$$f(v)\psi^u = e\psi^{(\zeta v, v)}\psi^u = e\psi^{(\zeta vu, vu)} = f(vu)$$

so that $f(v)\psi^u = f(v) \Leftrightarrow u = 1$.

2. The E_2 -page of the Bousfield-Kan spectral sequence, $E_s^{st} = \lim_{\mathbf{I}}^{-s} \underline{\pi}_t(v)$, converging to $\pi_{s+t}(\text{BC}_{\mathbb{F}_4}(f(v)\text{DI}_2))$ vanishes completely (3.6).

3. The Weyl group $W_{\mathbb{F}_4}(f(v)\text{DI}_2)$ is [23] the stabilizer subgroup for $\text{Out}(\text{DI}_2)$ acting on $f(v) \in \text{Mono}(\text{DI}_2, \mathbb{F}_4)$. We have just seen that this stabilizer is trivial. □

In a way, (3.8.1) says that \mathbb{F}_4 contains a unique copy of DI_2 .

Finally, we determine the set $\text{Rep}(\text{DI}_2, \mathbb{F}_4) = [\text{BDI}_2, \text{BF}_4]$ of all maps up to homotopy.

Proposition 3.9. *Any non-trivial morphism of DI_2 to \mathbb{F}_4 is a monomorphism.*

Proof. Let $f: \text{DI}_2 \rightarrow \text{F}_4$ be any non-trivial morphism. Then $f|T(\text{DI}_2) = W(\text{F}_4)A(v) \in [\text{BT}(\text{DI}_2), \text{BF}_4]$ for some 3-adic integer $v \in \mathbf{Z}_3$. This 3-adic integer, v , must be non-zero as f is non-trivial [22, 6.7] and even a 3-adic unit by (2.2). This shows that $A(v)$ is split injective and hence [26, 3.6] [24, 5.2] that f is a monomorphism. \square

Proof of Theorem 1.2. We conclude from (3.7, 3.8, 3.9) that

$$\text{Rep}(\text{DI}_2, \text{F}_4) = \{0\} \cup \text{Mono}(\text{DI}_2, \text{F}_4)$$

is in bijection with the set $\{0\} \cup \mathbf{Z}_3^*/\{\pm 1\}$. \square

In the following, we let $\text{B}\alpha: \text{BDI}_2 \rightarrow \text{BF}_4$ denote the monomorphism $\text{B}f(-1)$ corresponding (3.7) to the admissible homomorphism with matrix $A(-1)$ (2.6).

4. THE HOMOGENEOUS SPACE F_4/DI_2

In this section we compute the mod 3 cohomology of the exotic homogeneous space F_4/DI_2 .

The unstable mod 3 cohomology of F_4 is known since Borel [5]:

$$H^*(\text{F}_4; \mathbf{F}_3) \cong E[z_3, z_7, z_{11}, z_{15}] \otimes P[z_8]/(z_8^3)$$

with $P^1 z_3 = z_7$, $\beta z_7 = z_8$ and $P^1 z_{11} = z_{15}$.

The 3-complete space BDI_2 realizes the rank 2 mod 3 Dickson algebra meaning that

$$H^*(\text{BDI}_2; \mathbf{F}_3) \cong P[x_{12}, x_{16}]$$

with $P^1 x_{12} = x_{16}$.

The homogeneous space F_4/DI_2 is defined as the homotopy fibre of the map $\text{BDI}_2 \xrightarrow{\text{B}\alpha} \text{BF}_4$, and we have a sequence of fibrations

$$(4.1) \quad \text{DI}_2 \xrightarrow{\Omega \text{B}\alpha} \text{F}_4 \xrightarrow{\pi} \text{F}_4/\text{DI}_2 \xrightarrow{j} \text{BDI}_2 \xrightarrow{\text{B}\alpha} \text{BF}_4$$

We now investigate the value of the functor $H^*(-; \mathbf{F}_3)$ on this sequence.

Proposition 4.2. *The effects on mod 3 cohomology of the maps π and j of (4.1) are as follows:*

1. *The unstable mod 3 cohomology algebra of F_4/DI_2 is*

$$H^*(\text{F}_4/\text{DI}_2; \mathbf{F}_3) \cong E[z_3, z_7] \otimes P[z_8]/(z_8^3)$$

with $P^1 z_3 = z_7$ and $\beta z_7 = z_8$.

2. $H^*(\pi; \mathbf{F}_3)$ is the obvious inclusion

$$\begin{aligned} H^*(F_4/DI_2; \mathbf{F}_3) &\cong E[z_3, z_7] \otimes P[z_8]/(z_8^3) \\ &\subseteq E[z_3, z_7, z_{11}, z_{15}] \otimes P[z_8]/(z_8^3) \cong H^*(F_4; \mathbf{F}_3) \end{aligned}$$

of algebras.

3. $H^{>0}(j; \mathbf{F}_3)$ is the trivial homomorphism $H^{>0}(\text{BDI}_2; \mathbf{F}_3) \rightarrow H^{>0}(F_4/DI_2; \mathbf{F}_3)$.

Proof. It is a calculation with the Serre spectral sequence for the fibration $F_4 \xrightarrow{\pi} F_4/DI_2 \xrightarrow{j} \text{BDI}_2$. This starts at $E_2^{p,q} \cong H^p(\text{BDI}_2; \mathbf{F}_3) \otimes H^q(F_4; \mathbf{F}_3)$.

By degree reasons the classes z_3, z_7 and z_8 in the vertical edge are permanent cycles. Assume that $z_{11} \in E_2^{0,*}$ is a permanent cycle. Then $z_{15} = P^1 z_{11}$ is also a permanent cycle and therefore the spectral sequence collapses at E_2 . But this is impossible because the fibre F_4/DI_2 of the monomorphism $\text{B}\alpha$ is \mathbf{F}_3 -finite [12].

Hence, the class z_{11} transgresses to $\pm x_{12} \in H^*(\text{BDI}_2; \mathbf{F}_3) \cong E_2^{*,0}$ and then $z_{15} = P^1 z_{11}$ transgresses to $\pm x_{16} = P^1(\pm x_{12})$. After this last, the spectral sequence collapses to $E_\infty^{*,*} \cong E_\infty^{0,*} \cong E[z_3, z_7] \otimes P[z_8]/(z_8^3)$ and the edge homomorphisms are an injection

$$H^*(F_4/DI_2; \mathbf{F}_3) \cong E_\infty^{0,*} \cong E[z_3, z_7] \otimes P[z_8]/(z_8^3) \hookrightarrow E_2^{0,*} \cong H^*(F_4; \mathbf{F}_3)$$

and the trivial homomorphism

$$H^*(\text{BDI}_2; \mathbf{F}_3) \cong E_2^{*,0} \rightarrow E_\infty^{0,*} \cong \mathbf{F}_3 \subset H^*(F_4/DI_2; \mathbf{F}_3)$$

which proves the proposition. \square

Remark 4.3. The mod 3 cohomology of F_4/DI_2 coincides with that of Harper's molecule [16] at the prime 3 and this implies that they are actually homotopy equivalent up to 3-completion. Details will be worked out in Section 7.

5. FRIEDLANDER'S EXCEPTIONAL ISOGENY

In [14], E.M. Friedlander showed the existence of a self-homotopy equivalence $\text{B}\varphi$ of $(\text{BF}_4)_{1/2}$ that restricts to the maximal torus to the isogeny $\text{BT}(\varphi): \text{BT}(\text{F}_4) \rightarrow \text{BT}(\text{F}_4)$ determined by the matrix

$$\pi_2(\text{BT}(\varphi)) = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

acting on $\pi_2(\text{BT}(\text{F}_4)) \cong \mathbf{Z}_3^4$. The automorphism $\text{B}\varphi$ of the 3-compact group BF_4 satisfies the relation $\text{B}\varphi \circ \text{B}\varphi \simeq \text{B}\psi^2$. To see this, note that

$W(F_4)T(\varphi)^2 = W(F_4)T(\psi^2)$ and recall [21] that BF_4 has N -determined automorphisms.

Proposition 5.1. *There is a homotopy commutative diagram*

$$(5.2) \quad \begin{array}{ccc} \text{BDI}_2 & \xrightarrow{\text{B}\alpha} & \text{BF}_4 \\ \text{B}\psi^\zeta \downarrow & & \downarrow \text{B}\varphi \\ \text{BDI}_2 & \xrightarrow{\text{B}\alpha} & \text{BF}_4 \end{array}$$

where $\zeta^2 = -2$ and $\text{B}\psi^\zeta \in [\text{BDI}_2, \text{BDI}_2]$ is the corresponding Adams map.

Proof. According to Theorem 3.7 it is enough to check the restriction to the maximal torus of BDI_2 , and in fact,

$$\begin{aligned} \pi_2(\text{BT}(\varphi)) \cdot \pi_2(\text{BT}(\alpha)) &= \begin{pmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{pmatrix} \cdot \begin{pmatrix} \zeta & -1 \\ -\zeta & -1 + \zeta \\ 0 & -1 - \zeta \\ 2 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 2\zeta & -\zeta \\ 0 & -2 + \zeta \\ -2 & -\zeta \\ 2 & -2 - \zeta \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ 0 & -\zeta - 1 \\ -\zeta & 1 \\ -\zeta & -\zeta + 1 \end{pmatrix} \cdot \zeta \in W(F_4)(\pi_2(\text{BT}(\alpha)) \cdot \zeta) \end{aligned}$$

hence $\text{B}\varphi \circ \text{B}\alpha = \text{B}\alpha \circ \text{B}\psi^\zeta$. \square

Proposition 5.3. *The isomorphism $H^*(\text{B}\varphi; \mathbf{Q}_3)$ of*

$$H^*(\text{BF}_4; \mathbf{Q}_3) \cong P[x_4, x_{12}, x_{16}, x_{24}]$$

is, up to decomposables, determined by

$$\begin{aligned} \text{B}\varphi^*(x_4) &= 2x_4, & \text{B}\varphi^*(x_{12}) &= -8x_{12}, \\ \text{B}\varphi^*(x_{16}) &= 16x_{16}, & \text{B}\varphi^*(x_{24}) &= -64x_{24}. \end{aligned}$$

Proof. This is a long but straightforward calculation, easier done with the aid of a computer. This result coincides also with calculation done by Adams and Mahmud [1, Table 2.14]. \square

Define $\text{B}\tau: \text{BF}_4 \rightarrow \text{BF}_4$ to be the automorphism $\text{B}\tau = \text{B}\psi^{1/\zeta} \circ \text{B}\varphi$. Then $(\text{B}\tau)^2$ is homotopic to the identity and $\text{B}\tau \circ \text{B}\alpha = \text{B}\alpha$ by (5.1).

Corollary 5.4. *$H^4(\text{B}\tau; \mathbf{Z}_3)$ is multiplication by -1 on $H^4(\text{BF}_4; \mathbf{Z}_3)$.*

Proof. The effect of $H^4(\text{B}\tau; \mathbf{Q}_3)$ on $H^4(\text{BF}_4; \mathbf{Q}_3)$ is given by

$$H^4(\text{B}\tau; \mathbf{Q}_3) = H^4(\text{B}\varphi; \mathbf{Q}_3) \circ H^4(\text{B}\psi^{1/\zeta}; \mathbf{Q}_3) = 2 \cdot \zeta^{-2} = -1$$

by (5.3) and because the unstable Adams operation ψ^λ induces multiplication by λ^i in degree $2i$. \square

Let $\tau: F_4/DI_2 \rightarrow F_4/DI_2$ denote the self-homotopy equivalence of the exotic homogeneous space F_4/DI_2 induced by $B\tau$ on BF_4 and the identity on BDI_2 . This map makes the diagram

$$\begin{array}{ccccc} F_4/DI_2 & \xrightarrow{j} & BDI_2 & \xrightarrow{B\alpha} & BF_4 \\ \downarrow \tau & & \parallel & & \downarrow B\tau \\ F_4/DI_2 & \xrightarrow{j} & BDI_2 & \xrightarrow{B\alpha} & BF_4 \end{array}$$

commute up to homotopy.

Corollary 5.5. *The involution $H^*(\tau; \mathbf{F}_3)$ of $H^*(F_4/DI_2; \mathbf{F}_3)$ sends the generators z_3, z_7 , and z_8 to $-z_3, -z_7$, and $-z_8$, respectively.*

Proof. That $H^*(\tau; \mathbf{F}_3)(z_3) = -z_3$ follows from (5.4) since (4.2.2) the mod 3 cohomology of F_4/DI_2 embeds in the mod 3 cohomology of F_4 . As the two other generators are linked to z_3 by Steenrod operations (4.2.1), $H^*(\tau; \mathbf{F}_3)(z_3)$ must also act as multiplication by -1 on them. \square

We shall return to the homotopy involution $B\tau$ in Section 6.

6. BDI_2 AS A HOMOTOPY FIXED POINT SPACE

The automorphism $B\tau = B\psi^{1/\zeta} \circ B\varphi$ of BF_4 is a homotopy involution, $(B\tau)^2 \simeq 1$, that homotopy fixes BDI_2 in the sense that the diagram

$$(6.1) \quad \begin{array}{ccc} BDI_2 & \xrightarrow{B\alpha} & BF_4 \\ \parallel & & \downarrow B\tau \\ BDI_2 & \xrightarrow{B\alpha} & BF_4 \end{array}$$

commutes up to homotopy (5.1). This suggests that BDI_2 should be fixed under $B\tau$ in some sense. To make this precise, we need to rigidify the above two properties.

Proposition 6.2. *We may assume that the maps $B\alpha$ and $B\tau$ satisfy the identities $B\tau \circ B\alpha = B\alpha$ and $B\tau \circ B\tau = 1$.*

Proof. Let \mathbf{D} be the category

$$\cdot \xrightarrow{a} \cdot \begin{array}{c} \circlearrowright \\ b \end{array}$$

with two objects and two non-identity morphisms, a and b , subject to the relations $ba = a$ and $bb = 1$. The \mathbf{D} -diagram

$$(6.3) \quad BDI_2 \xrightarrow{B\alpha} BF_4 \begin{array}{c} \circlearrowright \\ B\tau \end{array}$$

in the homotopy category of spaces is centric; in fact, all the relevant mapping spaces are weakly contractible (3.8.2) [23, 13]. Hence (6.3) has an essentially unique realization in the category of spaces [11, 1.1]. \square

With such specific realizations of $B\alpha$ and $B\tau$, let F_4/DI_2 denote the homotopy fibre of $B\alpha$ over any point fixed by $B\tau$, e.g. any point in the image of $B\alpha$. Then F_4/DI_2 is $\mathbf{Z}/2$ -space and there is a homotopy fibre sequence

$$F_4/DI_2 \xrightarrow{j} BDI_2 \xrightarrow{B\alpha} BF_4$$

of $\mathbf{Z}/2$ -spaces and $\mathbf{Z}/2$ -maps which induces [7, XI.7.1] a homotopy fibre sequence

$$(6.4) \quad (F_4/DI_2)^{h\mathbf{Z}/2} \xrightarrow{j^{h\mathbf{Z}/2}} BDI_2^{h\mathbf{Z}/2} \xrightarrow{(B\alpha)^{h\mathbf{Z}/2}} BF_4^{h\mathbf{Z}/2}$$

of homotopy fixed point spaces. Here,

$$BDI_2^{h\mathbf{Z}/2} \simeq \text{map}(B\mathbf{Z}/2, BDI_2) \simeq BDI_2$$

since $\mathbf{Z}/2$ acts trivially on the 3-complete space BDI_2 . As to the base space, the Bousfield-Kan spectral sequence for the homotopy fixed point space $BF_4^{h\mathbf{Z}/2}$ degenerates to the formula $\pi_*(BF_4^{h\mathbf{Z}/2}) = \pi_*(BF_4)^{\mathbf{Z}/2}$. In particular, this space is non-empty and connected and its homotopy consists of the invariant part of the homotopy for BF_4 , cf. [36]. The aim is to show that the fibre of (6.4) is contractible in order to obtain a homotopy equivalence between BDI_2 and $BF_4^{h\mathbf{Z}/2}$.

Proposition 6.5. *The fibre, $(F_4/DI_2)^{h\mathbf{Z}/2}$, of fibration sequence (6.4) is contractible.*

Proof. By the general result at the end of this section (6.8), it suffices to show that the space E sitting in the homotopy pull back diagram

$$(6.6) \quad \begin{array}{ccc} E & \longrightarrow & F_4/DI_2 \\ \downarrow & & \downarrow (1, \tau) \\ F_4/DI_2 & \xrightarrow{\Delta} & (F_4/DI_2)^2 \end{array}$$

is contractible. The Eilenberg-Moore spectral sequence $E_r^{**} \Rightarrow H^*(E; \mathbf{F}_3)$ associated to (6.6) is a second quadrant cohomological spectral sequence with

$$E_2^{-pq} = \text{Tor}_p^{R \otimes R}(R, R)^q$$

where $R = H^*(F_4/DI_2; \mathbf{F}_3)$ and where R is a right $R \otimes R$ -module through the cup product $\mu: R \otimes R \rightarrow R$ and a left $R \otimes R$ -module through $\mu \circ (1 \otimes \tau^*)$. We shall show that $E_2^{**} = \mathbf{F}_3$.

Let κ be the algebra isomorphism of $R \otimes R$ given by

$$\kappa(z_i \otimes 1) = z_i \otimes 1 - 1 \otimes z_i, \quad \kappa(1 \otimes z_i) = 1 \otimes z_i$$

Then κ satisfies the identity $\varepsilon \otimes 1 = \mu\kappa$, where $\varepsilon: R \rightarrow \mathbf{F}_3$ is the augmentation homomorphism, and hence (6.10)

$$E_2^{-p*} = \mathrm{Tor}_p^R(\mathbf{F}_3, R)$$

where R is a left R -module via the algebra morphism

$$R \xrightarrow{1 \otimes \eta} R \otimes R \xrightarrow{\kappa} R \otimes R \xrightarrow{1 \otimes \tau^*} R \otimes R \xrightarrow{\mu} R$$

which takes z_i to $-z_i$, $i = 1, 2, 3$. In particular, R is a free R -module and the vanishing of the E_2 -page for the Eilenberg-Moore spectral sequence follows. \square

Proof of Theorem 1.3. By (6.4, 6.5), the monomorphism $B\alpha: \mathrm{BDI}_2 \rightarrow \mathrm{BF}_4$ induces a homotopy equivalence $\mathrm{BDI}_2 \simeq (\mathrm{BF}_4)^{h\mathbf{Z}/2}$. \square

Proof of (1.4) and Corollary 1.5. It is a general rule that $\mathrm{map}(X, Y^{hG}) = \mathrm{map}(X, Y)^{hG}$ and also that $\pi_i(M^{hG}) = \pi_i(M)^G$ when M is p -complete and $p \nmid |G|$. \square

Proof of Corollary 1.6. As a consequence of (1.5) and exactness of the Barratt-Puppe sequence, the map $j: \mathrm{F}_4/\mathrm{DI}_2 \rightarrow \mathrm{BDI}_2$ is null-homotopic. The splitting is now constructed using a section of $\pi: \mathrm{F}_4 \rightarrow \mathrm{F}_4/\mathrm{DI}_2$ and the H -space structure of F_4 . \square

In the homotopy splitting from Corollary 1.6 of F_4 , $\mathrm{F}_4/\mathrm{DI}_2 \simeq K(3)$ is one of Harper's H -spaces (7.5) and $\mathrm{DI}_2 \simeq B_3(3)$ is one of the Mimura-Toda bundles of completed spheres over spheres [20].

We finish this section with the general results that we used above to compute the homotopy fixed point space $(\mathrm{F}_4/\mathrm{DI}_2)^{h\mathbf{Z}/2}$.

First, let G be a finite group and K a G -space. Define E to be the homotopy pullback

$$(6.7) \quad \begin{array}{ccc} E & \longrightarrow & K \\ \downarrow & & \downarrow \mu \\ K & \xrightarrow{\Delta} & \mathrm{map}(G, K) \end{array}$$

where Δ takes $x \in K$ to the constant map with value x and the action map μ takes x to the map $\mu(k)(h) = h^{-1}x$, $h \in G$. These two maps are equivariant if we equip $\mathrm{map}(G, K)$ with the action $(gu)(h) = u(g^{-1}h)$, $g, h \in G$, $u \in \mathrm{map}(G, K)$. Thus (6.7) is a diagram of G -spaces and G -maps.

Theorem 6.8. *If K is BG-null, the G -map $E \rightarrow K$ induces a weak homotopy equivalence $E^{hG} \rightarrow K^{hG}$ of homotopy fixed point spaces.*

Proof. Since the homotopy fixed point functor commutes with homotopy pull-backs [7, XI.4.3], there is an induced homotopy pull-back diagram

$$\begin{array}{ccc} E^{hG} & \longrightarrow & K^{hG} \\ \downarrow & & \downarrow \mu^{hG} \\ K^{hG} & \xrightarrow{\Delta^{hG}} & \text{map}(G, K)^{hG} \end{array}$$

of homotopy fixed point spaces. The map $\Delta^{hG}: K^{hG} \rightarrow \text{map}(G, K)^{hG}$ can be identified to the evaluation map $\text{map}(BG, K) \rightarrow K$. Indeed, $K^{hG} = \text{map}(BG, K)$, since the action is trivial in the lower left corner, and $\text{map}(G, K)^{hG} = \text{map}(G_{hG}, K) = \text{map}(*, K) = K$. If K is BG-null, Δ^{hG} is a homotopy equivalence and so is then the top horizontal map $E^{hG} \rightarrow K^{hG}$. \square

By an argument dual to that of [19, 5.1] [27, p. 282], we may identify E and the homotopy limit of the diagram $K \xrightarrow{G} K$ consisting of the $|G|$ maps $K \rightarrow K$ given by $x \rightarrow gx$ for $g \in G$.

The final result that we used above was an algebraic computation ensuring the vanishing of an Eilenberg-Moore spectral sequence.

Let R be a unital and augmented (graded) algebra over a field k . The structure maps

$$\mu: R \otimes R \rightarrow R, \quad \eta: k \rightarrow R, \quad \varepsilon: R \rightarrow k$$

make R a right $R \otimes R$ -module and k a right R -module.

For any right R -module A , the tensor product $A \otimes R$ is a right $R \otimes R$ -module with right $R \otimes R$ -multiplication given by

$$(a \otimes r) \cdot (s \otimes t) = as \otimes rt$$

for all elements $a \in A$, $r, s, t \in R$ (modified by the usual sign in the graded case). When $A = k$, in particular, right multiplication in $k \otimes R = R$ by $s \otimes t \in R \otimes R$ is right multiplication in R by $(\varepsilon \otimes 1)(s \otimes t) = \varepsilon(s)t$.

Lemma 6.9. *Let κ be an algebra isomorphism of $R \otimes R$ and $\otimes_R(R \otimes R)$ the left adjoint functor to the forgetful functor induced by the algebra homomorphism $\kappa \circ (1 \otimes \eta): R \rightarrow R \otimes R$. Then*

1. $A \otimes_R(R \otimes R) = A \otimes R$ with right $R \otimes R$ -multiplication $(a \otimes r)(s \otimes t) = (a \otimes r) \cdot \kappa^{-1}(s \otimes t)$, and
2. $\text{Tor}_p^R(A, B) = \text{Tor}_p^{R \otimes R}(A \otimes_R(R \otimes R), B)$ and $\text{Ext}_R^p(A, B) = \text{Ext}_{R \otimes R}^p(A \otimes_R(R \otimes R), B)$,

for all right R -modules A , all left $R \otimes R$ -modules B , and all $p \geq 0$.

Proof. We have

$$U(\kappa(1 \otimes \eta)) = U(1 \otimes \eta)U(\kappa) \quad \text{and} \quad U(\kappa)L(\kappa(1 \otimes \eta)) = L(1 \otimes \eta)$$

where $L(f)$ denotes the left adjoint to the forgetful functor $U(f)$ induced by the algebra homomorphism f . Note that $L(1 \otimes \eta)(A) = A \otimes R$. It follows that $L(1 \otimes \eta)$ is exact; it also takes R -projectives to $R \otimes R$ -projectives because it is left adjoint to a (right) exact functor [32, 2.3.10]. The same is true of the functor $L(\kappa(1 \otimes \eta))$ which differs from $L(1 \otimes \eta)$ by an isomorphism. Similarly, $U(\kappa(1 \otimes \eta))$ is an exact functor that takes injectives to injectives. Therefore, the identities

$$A \otimes_R UB = LA \otimes_{R \otimes R} B, \quad \text{Hom}_R(A, UB) = \text{Hom}_{R \otimes R}(LA, B)$$

where $L = L(\kappa(1 \otimes \eta))$ and $U = U(\kappa(1 \otimes \eta))$, prolong to the identities of (6.9.2). \square

Corollary 6.10. *Suppose in addition to the assumptions of (6.9) that $\varepsilon \otimes 1 = \mu\kappa$. Then $\kappa \circ (1 \otimes \eta)$ induces an isomorphism*

$$\text{Tor}_p^R(k, B) = \text{Tor}_p^{R \otimes R}(R, B)$$

for all left $R \otimes R$ -modules B and all $p \geq 0$.

Proof. Note that $L(1 \otimes \eta)(k) = U(\varepsilon \otimes 1)(R)$ where R is considered a module over itself. Hence $L(\kappa(1 \otimes \eta))(k) = U(\kappa^{-1})L(1 \otimes \eta)(k) = U(\kappa^{-1})U(\varepsilon \otimes 1)(R) = U((\varepsilon \otimes 1)\kappa^{-1})(R) = U(\mu)(R)$ which is R as an $R \otimes R$ -module. \square

7. APPENDIX: HARPER'S MOLECULE

This section contains some comments on John Harper's work on finite H -spaces. Our starting point will be the theorem below and our main result (7.3) says that the space $K(p)$ is cohomologically unique in that it is determined up to homotopy equivalence by cohomological information. This will enable us (7.5) to identify $K(3)$ and the fibre F_4/DI_2 of the embedding $B\alpha: BDI_2 \rightarrow BF_4$.

Theorem 7.1. ([16, Theorem B]) *For each odd prime p there exists a simply connected finite complex $K(p)$ whose p -localization is an H -space and with*

$$\begin{aligned} H^*(K(p); \mathbf{F}_p) &\cong E[x_3, x_{2p+1}] \otimes P[x_{2p+2}]/(x_{2p+2})^p, \\ P^1 x_3 &= x_{2p+1}, \quad \beta x_{2p+1} = x_{2p+2}. \end{aligned}$$

It turns out that the p -completed homotopy type of the space $K(p)$ is determined by its mod p cohomology as an algebra over the Steenrod algebra. This is proved by means of classical homotopy theory methods and we are sure that it is known to J. Harper and perhaps other people. We will provide a prove for the sake of completeness.

For this aim we sketch the construction of this p -completed homotopy type by G.E. Cooke and L. Smith [9]. They first introduce the stable two stage Postnikov system

$$\begin{array}{ccc} A(p) & \xrightarrow{\quad\quad\quad} & * \\ \downarrow & & \downarrow \\ K(\mathbf{Z}/p, 2p^2 + 1) & \xrightarrow{\Psi} & K(\mathbf{Z}/p, 2p^2 + 2p - 1) \times K(\mathbf{Z}, 2p^2 + 2p + 1) \end{array}$$

where $\Psi = (P^1\iota, \beta P^1\beta\iota)$ (Notice that $\beta P^1\beta\iota \in H^{2p^2+2p+1}(\mathbf{Z}/p, 2p^2+1; \mathbf{Z}/p)$ is a mod p reduction of an integral class.) This is constructed in such a way that the composition

$$K(\mathbf{Z}, 3) \xrightarrow{P^p P^1\iota} K(\mathbf{Z}/p, 2p^2 + 1) \xrightarrow{\Psi} K(\mathbf{Z}/p, 2p^2 + 2p - 1) \times K(\mathbf{Z}, 2p^2 + 2p + 1)$$

is null-homotopic and therefore the first map lifts (non-uniquely) to φ_p :

$$\begin{array}{ccc} & & A(p) \\ & \nearrow \varphi_p & \downarrow \\ K(\mathbf{Z}, 3) & \xrightarrow{P^p P^1\iota} & K(\mathbf{Z}/p, 2p^2 + 1) \end{array}$$

Now define $Y(p)$ as the homotopy fibre

$$\begin{array}{ccc} Y(p) & \xrightarrow{\quad\quad\quad} & * \\ \downarrow q_p & & \downarrow \\ K(\mathbf{Z}, 3) & \xrightarrow{\varphi_p} & A(p) \end{array}$$

of the map φ_p .

The cohomology of the two stage Postnikov system $A(p)$ is computed from results of [29]:

$$H^*(A(p); \mathbf{F}_p) \cong E[\iota] \otimes P[\beta\iota, P^1\beta\iota] \otimes H$$

where ι is restricted from the fundamental class of $K(\mathbf{Z}/p, 2p^2 + 1)$ and, in particular, $\deg \iota = 2p^2 + 1$ and H is a Hopf algebra that is $(2p^2 + 4p - 2)$ -connected. Now, $\varphi_p^*(\iota) = P^p P^1\iota_3$, $\varphi_p^*(\beta\iota) = \beta P^p P^1\iota_3$ and $\varphi_p^*(P^1\beta\iota) =$

$P^1\beta P^p P^1\iota_3 = (\beta P^1\iota_3)^p$, from which it is computed the cohomology of $Y(p)$ in low dimensions:

Proposition 7.2. ([9, 1.2]) *In dimensions less than or equal to $2p^2 + 2p + 2$ the mod p cohomology of $Y(p)$ coincides with $E[u, P^1u] \otimes P[\beta P^1u]/(\beta P^1u)^p$ where $u = q_p^*\iota_3$.*

It then follows that stage $2p^2 + 2p + 2$ in the homology decomposition tower [17, Chp. 8] [4] for $Y(p)$ is a space $K(p) = Y(p)^{2p^2+2p+2}$ with

$$H^*(K(p); \mathbf{F}_p) \cong E[u, P^1u] \otimes P[\beta P^1u]/(\beta P^1u)^p.$$

Dually, the Postnikov tower for $K(p)$ has the form

$$\begin{array}{ccccc} & & \vdots & & \\ & & \downarrow & & \\ & & K(p)_{2p^2+2p+3} & & \\ & \nearrow & \downarrow & & \\ K(p) & \longrightarrow & K(p)_{2p^2+2p+2} & \longrightarrow & K(\pi_{2p^2+2p+3}(K(p)), 2p^2 + 2p + 4) \\ & \searrow & \downarrow & & \\ & & Y(p) & \longrightarrow & K(\pi_{2p^2+2p+2}(K(p)), 2p^2 + 2p + 3) \end{array}$$

because $K(p)_{2p^2+2p+1} = Y(p)^{2p^2+2p+2}_{2p^2+2p+1} = Y(p)_{2p^2+2p+1} = Y(p)$ appears at stage $2p^2 + 2p + 1$.

We now come to our main result of this section which is homotopy uniqueness at the prime p for $K(p)_p^\wedge$.

Theorem 7.3. *Let X be a p -complete space with cohomology $H^*(X; \mathbf{F}_p) \cong H^*(K(p); \mathbf{F}_p)$ as algebras over the Steenrod algebra. Then $X \simeq K(p)_p^\wedge$.*

Proof. Assume that X is a p -complete space with

$$H^*(X; \mathbf{F}_p) \cong E[x_3, x_{2p+1}] \otimes P[x_{2p+2}]/(x_{2p+2})^p$$

satisfying $P^1x_3 = x_{2p+1}$ and $\beta x_{2p+1} = x_{2p+2}$. Notice that X is 2-connected and since the mod p cohomology is finite dimensional, each $\pi_i(X)$ and $H^i(X; \mathbf{Z}_p)$ is a finitely generated \mathbf{Z}_p -module.

The Bockstein spectral sequence with p -adic coefficients applies. It collapses at

$$B_\infty \cong B_2 \cong E[x_3, x_{2p+1}(x_{2p+2})^{p-1}]$$

and shows that the top integral class lies in the group $H_{2p^2+2p+2}(X; \mathbf{Z}_p) \cong H^{2p^2+2p+2}(X; \mathbf{Z}_p) \cong \mathbf{Z}_p$. (The cohomological dimension $\text{cd}_{\mathbf{Z}_p}(X) = 2p^2 +$

$2p + 2$.) In particular, $H^{>2p^2+2p+2}(X; M) = 0$ for any \mathbf{Z}_p -module M and thus $[X, K(p)_p^\wedge] = [X, Y(p)_p^\wedge]$ by obstruction theory.

We now use the p -completed version of the Cooke-Smith construction of $K(p)$. Let

$$X \xrightarrow{x_3} K(\mathbf{Z}_p, 3)$$

represent a generator of $H^3(X; \mathbf{Z}_p) \cong \mathbf{Z}_p$.

Claim 7.4. $[X, A(p)_p^\wedge] = \{*\}$.

Proof. This follows from the exact sequence of sets

$$\begin{aligned} 0 &= H^{2p^2+2p-2}(X; \mathbf{F}_p) \times H^{2p^2+2p}(X; \mathbf{F}_p) \\ &\rightarrow [X, A(p)_p^\wedge] \rightarrow H^{2p^2+1}(X; \mathbf{F}_p) = 0 \end{aligned}$$

obtained by mapping X into the fibration $K(\mathbf{Z}/p, 2p^2+2p-2) \times K(\mathbf{Z}_p, 2p^2+2p) \rightarrow A(p)_p^\wedge \rightarrow K(\mathbf{Z}/p, 2p^2+1)$. \square

Since $\varphi_p x_3$ is nullhomotopic by Claim 7.4, $x_3: X \rightarrow K(\mathbf{Z}_p, 3)$ lifts to a map

$$f: X \rightarrow Y(p)_p^\wedge$$

that satisfies $f^*(u) = x_3$. Hence f^* is an isomorphism in dimensions $\leq 2p^2 + 2p + 2$ and the corresponding map $X \rightarrow K(p)_p^\wedge$ an isomorphism on mod p cohomology. \square

In the special case where $p = 3$, we note that Harper's molecule $K(3)_3^\wedge$ and the exotic homogeneous space F_4/DI_2 have isomorphic mod 3 cohomology algebras over the Steenrod algebra. Thus we may conclude that these two spaces are homotopy equivalent.

Proposition 7.5. F_4/DI_2 and $K(3)_3^\wedge$ are homotopy equivalent 3-complete spaces.

We do not know if $K(p)$ for $p > 3$ is an exotic homogeneous space with respect to some monomorphism of p -compact groups.

REFERENCES

- [1] J.F. Adams and Z. Mahmud, *Maps between classifying spaces*, Invent. Math. **35** (1976), 1–41.
- [2] J. Aguadé, *Constructing modular classifying spaces*, Israel J. Math. **66** (1989), 23–40.
- [3] K.K.S. Andersen, *Cohomology of Weyl groups with applications to topology*, Master's thesis, Københavns Universitet, August 1997.
- [4] Marc Aubry, *Homotopy theory and models*, Birkhäuser Verlag, Basel, 1995, Based on lectures held at a DMV seminar in Blaubeuren by H. J. Baues, S. Halperin and J.-M. Lemaire.

- [5] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math. (2) **57** (1953), 115–207.
- [6] N. Bourbaki, *Groupes et algèbres de Lie, Chp. 9*, Masson, Paris, 1982.
- [7] A.K. Bousfield and D.M. Kan, *Homotopy limits, completions and localizations*, 2nd ed., Lecture Notes in Mathematics, vol. 304, Springer-Verlag, Berlin–Heidelberg–New York–London–Paris–Tokyo, 1987.
- [8] A. Clark and J.R. Ewing, *The realization of polynomial algebras as cohomology rings*, Pacific J. Math **50** (1974), 425–434.
- [9] George E. Cooke and Larry Smith, *On realizing modules over the Steenrod algebra*, J. Pure Appl. Algebra **13** (1978), no. 1, 71–100.
- [10] L.E. Dickson, *A fundamental system of invariants of the general modular linear group with a solution of the form problem*, Trans. Amer. Math. Soc. **12** (1911), 75–98.
- [11] W.G. Dwyer and D.M. Kan, *Centric maps and realizations of diagrams in the homotopy category*, Proc. Amer. Math. Soc. **114** (1992), 575–584.
- [12] W.G. Dwyer and C.W. Wilkerson, *Homotopy fixed point methods for Lie groups and finite loop spaces*, Ann. of Math. (2) **139** (1994), 395–442.
- [13] ———, *The center of a p -compact group*, The Čech Centennial. Contemporary Mathematics, vol. 181 (Providence, Rhode Island) (M. Cenk and H. Miller, eds.), American Mathematical Society, 1995, pp. 119–157.
- [14] Eric M. Friedlander, *Exceptional isogenies and the classifying spaces of simple Lie groups*, Ann. Math. (2) **101** (1975), 510–520.
- [15] John R. Harper, *The mod 3 homotopy type of F_4* , (1974), 58–67. Lecture Notes in Math., Vol. 418.
- [16] ———, *H-spaces with torsion*, Mem. Amer. Math. Soc. **22** (1979), no. 223, viii+72.
- [17] Peter Hilton, *Homotopy theory and duality*, Gordon and Breach Science Publishers, New York, 1965.
- [18] Stefan Jackowski, James McClure, and Bob Oliver, *Maps between classifying spaces revisited*, The Čech centennial (Boston, MA, 1993), Contemp. Math., vol. 181, Amer. Math. Soc., Providence, RI, 1995, pp. 263–298.
- [19] Stefan Jackowski and James E. McClure, *Homotopy approximations for classifying spaces of compact Lie groups*, Algebraic topology (Arcata, CA, 1986), Springer, Berlin, 1989, pp. 221–234.
- [20] Mamoru Mimura and Hiroshi Toda, *Cohomology operations and homotopy of compact Lie groups. I*, Topology **9** (1970), 317–336.
- [21] J.M. Møller, *F_4 as a 3-compact group*, Preprint (in preparation).
- [22] ———, *Rational isomorphisms of p -compact groups*, Topology **35** (1996), 201–225.
- [23] ———, *The 3-compact group $DI(2)$* , Preprint, July 1997.
- [24] ———, *$PU(p)$ as a p -compact group*, Preprint, April 1997.
- [25] ———, *Toric morphisms of p -compact groups*, Preprint, November 1997.
- [26] J.M. Møller and D. Notbohm, *Centers and finite coverings of finite loop spaces*, J. reine angew. Math. **456** (1994), 99–133.
- [27] Jolanta Słomińska, *Homotopy colimits on $E-I$ -categories*, Algebraic topology Poznań 1989, Lecture Notes in Math., vol. 1474, Springer, Berlin, 1991, pp. 273–294.
- [28] L. Smith and R.M. Switzer, *Realizability and nonrealizability of Dickson algebras*, Proc. Amer. Math. Soc. **89** (1983), 303–313.
- [29] Larry Smith, *The cohomology of stable two stage Postnikov systems*, Illinois J. Math. **11** (1967), 310–329.

- [30] ———, *Polynomial invariants of finite groups*, Research Notes in Mathematics, vol. 6, A K Peters Ltd., Wellesley, MA, 1995.
- [31] Norman Steenrod, *Polynomial algebras over the algebra of cohomology operations*, H-spaces (Actes Réunion Neuchâtel, 1970), Springer, Berlin, 1971, Lecture Notes in Mathematics, Vol. 196, pp. 85–99.
- [32] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.
- [33] Clarence Wilkerson, *A primer on the Dickson invariants*, Proceedings of the North-western Homotopy Theory Conference (Evanston, Ill., 1982) (Providence, R.I.), Amer. Math. Soc., 1983, pp. 421–434.
- [34] Clarence W. Wilkerson, *Integral closure of unstable Steenrod algebra actions*, J. Pure Appl. Algebra **13** (1978), no. 1, 49–55.
- [35] Zdzisław Wojtkowiak, *On maps from holim F to Z* , Algebraic topology, Barcelona, 1986, Lecture Notes in Math., vol. 1298, Springer, Berlin, 1987, pp. 227–236.
- [36] A. Zabrodsky, *On the realization of invariant subgroups of $\pi_*(X)$* , Trans. Amer. Math. Soc. **285** (1984), no. 2, 467–496.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193
BELLATERRA, SPAIN

E-mail address: `broto@manwe.mat.uab.es`

MATEMATISK INSTITUT, UNIVERSITETSPARKEN 5, DK-2100 KØBENHAVN Ø

E-mail address: `moller@math.ku.dk`