

Relative group completions

CARLES CASACUBERTA AND AN DESCHEEMAEKER*

Abstract

We extend arbitrary group completions to the category of pairs (G, N) where G is a group and N is a normal subgroup of G . Relative localizations are defined as special cases. Our construction is a group-theoretical analogue of fibrewise completion and fibrewise localization in homotopy theory, and generalizes earlier work on relative localization of groups at primes. We use our approach to find conditions under which factoring out group radicals preserves exactness, and extend previous results about the effect of plus-constructions on homotopy fibre sequences.

0 Introduction

Relative completions and relative localizations of groups are group-theoretical analogues of fibrewise completions and fibrewise localizations in homotopy theory. The latter were first discussed in the work of Sullivan [24] and Bousfield–Kan [7], and since then by many other authors, primarily focusing on localization at primes; see e.g. [15], [16], [19]. More recently, Bousfield [6, § 4] and Dror Farjoun [13, Ch. 1] have defined fibrewise versions of homotopically idempotent transformations of spaces. The aim of this paper is to provide a similar tool for groups, and give applications within group theory and to homotopy theory.

Our initial goal was to state and prove that, for any group homomorphism f , the f -localization functor in the sense of [8] admits a relative version, that is, can be suitably defined on pairs of groups. However, we will present this result in much greater generality, by proving that every pointed endofunctor which is compatible with conjugation (as defined in Section 1) admits a relative version. All completions are compatible with

*The authors were supported by DGR under grants ACI98–23 and ACI99–34 during the preparation of this article. The first-named author was also supported by DGES under grant PB97–0202 and the second-named author was a fellow of the EC Research Training Network HPRN-CT-1999-00119.

conjugation. By a “completion” we mean the codensity monad of any full embedding (that is, any functorial inverse limit of targets in some class of groups), and a “localization” is an idempotent completion.

Our approach supplies, as a special case, the missing details in the claim made by Bousfield–Kan in [7, IV.5.6] about the existence of fibrewise R -completion of groups, where R is a commutative ring. Our construction also generalizes very naturally Hilton’s construction in [14] and the one described by Descheemaeker–Malfait in [12].

Although localizations are special cases of completions, the relevance of their universal property justifies a distinguished treatment. Thus, if L is any localization in the category of groups, then the relative L -localization of a pair (G, N) is a pair (G', LN) with the same cokernel as (G, N) , together with a morphism of pairs $(G, N) \rightarrow (G', LN)$ sharing the characteristic universal property of idempotent functors. We may describe it by saying that the effect of L on N has been uniformly extended over all cosets of G relative to N .

In the last two sections, we restrict to a special kind of group localizations, namely the ones obtained by dividing out radicals. These are called epireflections. Using their relative version, we analyze the effect of epireflections on group extensions. By doing so, we extend results of Berrick [2] about hypoabelianization of groups and preservation of homotopy fibre sequences under Quillen’s plus-construction. Our results specialize to the case of plus-constructions with respect to homology with mod n coefficients, among other cases. Further generalization should be possible along the lines of [4].

A preliminary version of the present article was useful to Rodríguez and Scevenels in [22, Theorem 6], in order to relate exactness of certain localizations with closure properties of the corresponding image classes. Some of their results are improved here in Section 4.

Acknowledgements Discussions with Jon Berrick, Warren Dicks and Armin Frei were beneficial to us during the preparation of this article.

1 Relativizing pointed endofunctors

Let \mathcal{C} be any category. A functor $L: \mathcal{C} \rightarrow \mathcal{C}$ together with a natural transformation $l: \text{id} \rightarrow L$ is called a *pointed endofunctor* or a *coaugmented functor* in the literature.

If (L, l) is a pointed endofunctor in any \mathcal{C} and G is a group, then (L, l) defines, in a canonical way, a pointed endofunctor in the category of G -objects (that is, objects of \mathcal{C} equipped with an action of G), with G -maps

as morphisms. Indeed, for every G -object N and every element $g \in G$, there is a commutative diagram

$$\begin{array}{ccc} N & \xrightarrow{g} & N \\ \wr \downarrow & & \downarrow \wr \\ LN & \xrightarrow{Lg} & LN \end{array} \quad (1.1)$$

yielding a G -action on LN such that \wr is a G -map.

From now on, we will exclusively work in the categories of groups and relative groups. A *relative group* is a pair (G, N) where G is a group and N is a normal subgroup of G . It is usually more convenient to regard a relative group as a group extension $N \twoheadrightarrow G \twoheadrightarrow Q$. A morphism of relative groups is a commutative diagram

$$\begin{array}{ccccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ \wr \downarrow & & f \downarrow & & \downarrow \\ N' & \twoheadrightarrow & G' & \twoheadrightarrow & Q' \end{array}$$

of group extensions. Note that such a diagram yields an action of G on N' via f , and \wr is automatically a G -map, that is,

$$\wr(gng^{-1}) = f(g)\wr(n)f(g)^{-1}$$

for all $n \in N$ and $g \in G$.

Now let (L, \wr) be any pointed endofunctor in the category of groups. As in [5] or [14], we look for suitable conditions ensuring that the diagram

$$\begin{array}{ccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ \wr \downarrow & & & & \\ LN & & & & \end{array}$$

can be embedded into a morphism of relative groups

$$\begin{array}{ccccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ \wr \downarrow & & f \downarrow & & \text{id} \downarrow \\ LN & \xrightarrow{j} & H & \xrightarrow{p} & Q, \end{array} \quad (1.2)$$

specializing to the standard homomorphism

$$l_*: H^2(Q; N) \rightarrow H^2(Q; LN)$$

whenever N and LN are abelian, and admitting a similar description in terms of cohomology classes of nonabelian 2-cocycles otherwise.

This is the precise group-theoretical analogue of fibrewise functorial transformations in homotopy theory, where “continuity” of the functor is a necessary assumption to ensure that it can be coherently extended over the fibres. Similarly, we have to require that (L, l) be compatible with conjugation, in the sense which we now make precise. Our aim is to guarantee that the action of G on LN via f in (1.2) agrees with the action defined by the functoriality of L as depicted in (1.1).

For a group N , we denote by $\text{Aut}(N)$ the group of automorphisms of N , and by $\text{Out}(N)$ the quotient of $\text{Aut}(N)$ by the inner automorphisms. That is, $\text{Out}(N)$ is the cokernel of the homomorphism $\tau: N \rightarrow \text{Aut}(N)$ given by $\tau_x(y) = xyx^{-1}$.

Definition 1.1 A pointed endofunctor (L, l) in the category of groups is *compatible with conjugation* if the diagram

$$\begin{array}{ccc} N & \xrightarrow{l} & LN \\ \tau \downarrow & & \tau \downarrow \\ \text{Aut}(N) & \xrightarrow{L} & \text{Aut}(LN) \end{array} \quad (1.3)$$

is commutative for every group N .

This property is not automatic. For instance, it fails if L is the identity functor and $l(n) = 1$ for every group N and all $n \in N$. On the other hand, as we prove in Section 2, all group completions are compatible with conjugation.

Note that, if (L, l) is compatible with conjugation, then L induces a group homomorphism

$$\text{Out}(N) \rightarrow \text{Out}(LN),$$

but the converse need not be true (as the same counterexample given above shows). Observe also that the diagram (1.3) trivially commutes if N and LN are abelian.

We next prove that compatibility with conjugation is sufficient to define a relative version of (L, l) . We will give two different, equivalent constructions. The first one is based on earlier work of Hilton [14] and Bokor–Hilton [5]. The second one generalizes the method of Descheemaeker–Malfait published in [12].

First construction Suppose given a group extension $N \twoheadrightarrow G \twoheadrightarrow Q$. Using the technique described in [5], [7, IV.5.6] or [14], we may embed the given extension into a commutative diagram with exact rows

$$\begin{array}{ccccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ \iota \downarrow & & f \downarrow & & \text{id} \downarrow \\ LN & \xrightarrow{j} & H & \xrightarrow{p} & Q, \end{array} \quad (1.4)$$

where we emphasize that the homomorphism j need not be injective if we omit the condition that (L, l) be compatible with conjugation. The group H and the maps in (1.4) are defined as follows; cf. [14]. Let $LN \rtimes G$ denote the semi-direct product with respect to the G -action on LN induced by functoriality from the conjugation action of G on N . Let S be the normal subgroup of $LN \rtimes G$ generated by all the elements of the form $(l(n)^{-1}, n)$ with $n \in N$. For $x \in LN$ and $g \in G$, we denote by $(x, g)S$ the class of (x, g) in the quotient

$$H = (LN \rtimes G)/S.$$

The homomorphism f is defined as $f(g) = (1, g)S$, and the homomorphisms j and p are given respectively by $j(x) = (x, 1)S$ and $p((x, g)S) = gN$. Then the diagram (1.4) commutes, and the kernel of p coincides with the image of j , since, for all $n \in N$ and $x \in LN$, we may write

$$(x, n) = (x l(n), 1)(l(n)^{-1}, n).$$

Now, the assumption that (L, l) is compatible with conjugation implies that j is injective. To see this, it is sufficient to prove that the set of elements of the form $(l(n)^{-1}, n)$ with $n \in N$ is closed under conjugation inside $LN \rtimes G$, and this follows precisely from the commutativity of (1.3) and the fact that $l: N \rightarrow LN$ is a G -map by (1.1).

Note that if L is the identity and $l(n) = 1$ for every N and all $n \in N$, then H is isomorphic to $(N/[N, N]) \rtimes Q$, so in this example j is not injective, unless N is abelian.

Second construction Recall that every extension $N \rtimes G \twoheadrightarrow Q$ of groups determines a group homomorphism $\psi: Q \rightarrow \text{Out}(N)$, which is called an abstract kernel. An extension $N \rtimes G \twoheadrightarrow Q$ is defined to be equivalent to another extension $N \rtimes G' \twoheadrightarrow Q$ if there is an isomorphism $G \cong G'$ inducing the identity on both N and Q . As explained e.g. in [17], there is a one-to-one correspondence between the set $\text{Ext}_\psi(Q, N)$ of equivalence classes of extensions $N \rtimes G \twoheadrightarrow Q$ inducing an abstract kernel $\psi: Q \rightarrow \text{Out}(N)$ and the set $H_\psi^2(Q; N)$ of cohomology classes of pairs (φ, c) , where $\varphi: Q \rightarrow \text{Aut}(N)$ is a function lifting ψ , and $c: Q \times Q \rightarrow N$ is a normalized 2-cocycle, that is,

$$\begin{aligned} \varphi(x) \circ \varphi(y) &= c(x, y) \varphi(xy) c(x, y)^{-1} \\ \varphi(x)(c(y, z)) c(x, yz) &= c(x, y) c(xy, z), \end{aligned}$$

for all $x, y, z \in Q$, and $\varphi(1) = \text{id}$ and $c(x, 1) = c(1, x) = 1$ for all $x \in Q$.

Given a 2-cocycle (φ, c) , the associated extension of N by Q can be explicitly described as the set $N \times Q$ equipped with the multiplication

$$(n, x)(m, y) = (n \varphi(x)(m) c(x, y), xy). \quad (1.5)$$

Now suppose given an extension $N \twoheadrightarrow G \twoheadrightarrow Q$ and a pointed endofunctor (L, l) . Choose a normalized set-theoretical section $\sigma: Q \rightarrow G$ and let (φ, c) be the corresponding 2-cocycle, classifying the given extension. Specifically, φ is the composite of σ with the conjugation homomorphism $G \rightarrow \text{Aut}(N)$, and c is defined as $c(x, y) = \sigma(x)\sigma(y)\sigma(xy)^{-1}$. Then the assumption that (L, l) is compatible with conjugation ensures that $(L \circ \varphi, l \circ c)$ is also a normalized 2-cocycle. Indeed, in order to verify that

$$L(\varphi(x)) \circ L(\varphi(y)) = l(c(x, y)) L(\varphi(xy)) l(c(x, y))^{-1},$$

one needs the functoriality of L and the assumption that (1.3) commutes. The second condition

$$L(\varphi(x))(l(c(y, z))) = l(c(x, y)) l(c(xy, z)) l(c(x, yz))^{-1}$$

only requires that l be a natural transformation.

The 2-cocycle $(L \circ \varphi, l \circ c)$ yields a group extension $LN \twoheadrightarrow H \twoheadrightarrow Q$, which is independent of the choices made, up to equivalence. Moreover, the homomorphism $f: G \rightarrow H$ sending (n, x) to $(l(n), x)$ renders commutative the diagram

$$\begin{array}{ccccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ l \downarrow & & f \downarrow & & \text{id} \downarrow \\ LN & \twoheadrightarrow & H & \twoheadrightarrow & Q, \end{array}$$

in such a way that the action of G on LN defined via f agrees with the action given by conjugation on N and functoriality of L .

We conclude by checking that the bottom extension in this diagram agrees, up to equivalence, with the extension obtained with the first construction. For this, define a section $\sigma': Q \rightarrow (LN \rtimes G)/S$ by $\sigma'(q) = (1, \sigma(q))S$. Then the corresponding 2-cocycle is

$$c'(x, y) = \sigma'(x)\sigma'(y)\sigma'(xy)^{-1} = (1, c(x, y))S = (l(c(x, y)), 1)S,$$

hence $c' = l \circ c$, as desired.

2 Relative completions and localizations

Our main goal in this section is to show that, if the pointed endofunctor (L, l) is a completion functor, then compatibility with conjugation is automatically fulfilled.

We first make precise what we mean by a completion functor. Let \mathcal{D} be any class of objects in a category \mathcal{C} and let $K: \mathcal{D} \rightarrow \mathcal{C}$ denote the full embedding. For any object X in \mathcal{C} , the objects of the comma category $(X \downarrow K)$ are the morphisms $X \rightarrow D$ with D in \mathcal{D} , and morphisms in $(X \downarrow K)$ are commutative triangles.

The \mathcal{D} -completion of an object X is the inverse limit \widehat{X} (if it exists) of the functor from $(X \downarrow K)$ to the category \mathcal{C} sending $X \rightarrow D$ to D . In other words, \mathcal{D} -completion is the codensity monad (or codensity triple) of the full embedding $K: \mathcal{D} \rightarrow \mathcal{C}$, i.e., the right Kan extension of K along itself; see [7, IV.2.2] following Artin–Mazur, and [18, p. 246].

Thus, if \mathcal{D} -completion exists for an object X , then there is a natural morphism $l: X \rightarrow \widehat{X}$, which may be viewed as “the closest approximation to X by an inverse limit of objects in \mathcal{D} ”. Well-known examples in the category of groups include profinite completion (where \mathcal{D} is the class of finite groups) and pronilpotent completion (where \mathcal{D} is the class of nilpotent groups). Each of these admits local versions at primes.

If \mathcal{D} -completion exists for all objects, then it is a pointed endofunctor. The existence of \mathcal{D} -completion is guaranteed for all objects if inverse limits exist in \mathcal{C} and the class \mathcal{D} is a set, but also in other cases, for instance when \mathcal{D} is reflective. If \widehat{X} belongs itself to \mathcal{D} , then \mathcal{D} -completion is idempotent on X . It is in fact idempotent on all objects in many cases, but not always (see [7, IV.2 and IV.5.4]).

Recall that a pointed endofunctor (L, l) is called *idempotent* if $Ll: L \rightarrow LL$ is an isomorphism, and $Ll = lL$; see [8] for a recent survey about this concept. Idempotent pointed endofunctors will be called *localizations*. Such functors L are characterized by the following universal property. For all X, Y and every morphism $\varphi: X \rightarrow LY$, there is a unique morphism $\psi: LX \rightarrow LY$ such that $\psi \circ l = \varphi$.

If L is a localization in any category, then L -local objects are those isomorphic to LX for some X , and L -equivalences are morphisms f such that Lf is an isomorphism. Thus, for every X , the morphism $l: X \rightarrow LX$ is an L -equivalence, and it is the unique L -equivalence from X to an L -local object, up to isomorphism. Furthermore, local objects X and equivalences $f: A \rightarrow B$ are *orthogonal*, in the sense that the induced function

$$f^*: \mathcal{C}(B, X) \rightarrow \mathcal{C}(A, X)$$

is a bijection, where $\mathcal{C}(B, X)$ denotes the set of morphisms $B \rightarrow X$ in \mathcal{C} .

Some examples of localizations in the category of groups are abelianization, hypoabelianization (i.e., dividing out the perfect radical), and localization at primes.

Localizations are special cases of completions. Specifically,

Proposition 2.1 *Every localization (L, l) is isomorphic to \mathcal{D} -completion where \mathcal{D} is the class of L -local groups.*

PROOF. For each object X , the morphism $l: X \rightarrow LX$ is an initial object in the comma category $(X \downarrow K)$, where $K: \mathcal{D} \rightarrow \mathcal{C}$ is the full embedding. \square

We now prove that completions are compatible with conjugation, as defined in Section 1.

Theorem 2.2 *Let \mathcal{D} be a class of groups such that \mathcal{D} -completion exists for all groups. Let N be any group and let $l: N \rightarrow \widehat{N}$ denote its \mathcal{D} -completion. Then the action of N on \widehat{N} induced by conjugation and functoriality satisfies*

$$n \cdot x = l(n) x l(n)^{-1} \quad \text{for all } n \in N \text{ and } x \in \widehat{N}.$$

PROOF. First we need to describe the action of N on \widehat{N} in explicit terms, using the fact that \widehat{N} is an inverse limit. For any element $n \in N$, let $\tau_n: N \rightarrow N$ denote conjugation by n . For each homomorphism $f_D: N \rightarrow D$ with D in \mathcal{D} , we consider the composite

$$N \xrightarrow{\tau_n} N \xrightarrow{f_D} D$$

and denote by $g_{n,D}$ the corresponding homomorphism $\widehat{N} \rightarrow D$ in the limiting cone, satisfying $g_{n,D} \circ l = f_D \circ \tau_n$ (abbreviated to g_D when $n = 1$). Then the automorphism $\gamma_n: \widehat{N} \rightarrow \widehat{N}$ induced by τ_n is uniquely characterized by the property that $g_D \circ \gamma_n = g_{n,D}$ for every $f_D: N \rightarrow D$.

Now fix any $n \in N$. Observe that, for each $f_D: N \rightarrow D$ with D in \mathcal{D} , we have

$$\tau_{f_D(n)} \circ f_D = f_D \circ \tau_n,$$

and this implies that $\tau_{f_D(n)} \circ g_D = g_{n,D}$. Thus, we write $\nu(x) = l(n) x l(n)^{-1}$ for $x \in \widehat{N}$ and check that ν satisfies the property that characterizes the automorphism $\gamma_n: \widehat{N} \rightarrow \widehat{N}$, namely

$$g_D(\nu(x)) = f_D(n) g_D(x) f_D(n)^{-1} = \tau_{f_D(n)}(g_D(x)) = g_{n,D}(x)$$

for all $x \in \widehat{N}$, as needed. \square

As we saw in Section 1, this implies that completion functors admit relative versions. This enlightens and generalizes the construction made by Bousfield–Kan in [7, IV.5.6], dealing with completion with respect to a ring.

Since localizations are special cases of completions, Theorem 2.2 applies to localizations as well. However, the proof of Theorem 2.2 is much simpler

in the case of localizations, since we may then use their universal property to infer the result from the equation

$$n \cdot l(m) = l(n)l(m)l(n)^{-1}$$

for $n, m \in N$, as done in earlier articles about localization at primes.

3 Universal property of relative localizations

In this section we make explicit the universal property of relative group localizations, hence generalizing substantial parts of [5], [9], [12], and [14].

Given a localization L in the category of groups, we say that a relative group $N \twoheadrightarrow G \twoheadrightarrow Q$ is L -local if N is an L -local group. A morphism of relative groups is an L -equivalence if it is orthogonal to all L -local relative groups.

Proposition 3.1 *If a morphism of relative groups*

$$\begin{array}{ccccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow \\ N' & \twoheadrightarrow & G' & \twoheadrightarrow & Q' \end{array}$$

is an L -equivalence, then β is an L -equivalence of groups and γ is an isomorphism.

PROOF. This is proved precisely as in [9, Proposition 1.3]. First use that (α, β, γ) is orthogonal to $D \twoheadrightarrow D \twoheadrightarrow 1$, where D is any L -local group, and then choose $1 \twoheadrightarrow E \twoheadrightarrow E$, where E is any group. \square

Our main result in this section is the following. It includes the statement that the middle map in a relative L -localization is an L -equivalence of groups, which is of great importance for the applications. Note the analogy with fibrewise localization in homotopy theory, by comparing it with [13, Theorem 1.F.3].

Theorem 3.2 *Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be any extension of groups, and let (L, l) be any localization in the category of groups. Then there is a commutative diagram*

$$\begin{array}{ccccc} N & \twoheadrightarrow & G & \twoheadrightarrow & Q \\ l \downarrow & & f \downarrow & & \text{id} \downarrow \\ LN & \twoheadrightarrow & H & \twoheadrightarrow & Q \end{array} \tag{3.1}$$

which is an L -equivalence of relative groups, and where f is an L -equivalence of groups.

PROOF. Theorem 2.2 tells us that localization functors are compatible with conjugation, and this suffices to obtain (3.1) with the property that the action of G on LN given via f agrees with the action given by functoriality of L , as shown in Section 1. Now we want to prove that the morphism (l, f, id) is an L -equivalence of relative groups. Suppose given a group extension $N' \twoheadrightarrow G' \twoheadrightarrow Q'$ where N' is L -local, and a morphism (α, β, γ) from $N \twoheadrightarrow G \twoheadrightarrow Q$ to $N' \twoheadrightarrow G' \twoheadrightarrow Q'$. Since $l: N \rightarrow LN$ is an L -localization of groups, there is a unique $\alpha': LN \rightarrow N'$ such that $\alpha' \circ l = \alpha$.

We first view H as $(LN \rtimes G)/S$, using the notation of Section 1, and define

$$\beta': H \rightarrow G', \quad \beta'((n, g)S) = \alpha'(n) \beta(g).$$

Then β' is a well-defined group homomorphism. In order to check this, the following equality is needed, as in [14, Theorem 1.2]:

$$\alpha'(g \cdot x) = \beta(g) \alpha'(x) \beta(g)^{-1} \quad \text{for all } g \in G \text{ and all } x \in LN. \quad (3.2)$$

This is proved by picking first $x \in \text{im } l$ and then using the universal property of L .

Alternatively, we may view H as an extension of LN by Q classified by $(L \circ \varphi, l \circ c)$, where (φ, c) is a 2-cocycle classifying $N \twoheadrightarrow G \twoheadrightarrow Q$. Then we define a function $\beta': H \rightarrow G'$ as follows, where H is written as a twisted product $LN \times Q$ with the multiplication described in (1.5):

$$\beta'(x, q) = (\alpha'(x), 1) \beta(1, q).$$

In order to show that β' is a group homomorphism, one needs that

$$\alpha'(((L \circ \varphi)(q))(x)) = \beta(1, q) (\alpha'(x), 1) \beta(1, q)^{-1},$$

and this follows again from the universal property of L , by checking it first for $x \in \text{im } l$.

In either case, we find that $(\alpha', \beta', \gamma)$ is a morphism of relative groups, and it satisfies

$$(\alpha', \beta', \gamma) \circ (l, f, \text{id}) = (\alpha, \beta, \gamma).$$

There only remains to prove the uniqueness of $(\alpha', \beta', \gamma)$. Suppose that $(\alpha'', \beta'', \gamma'')$ satisfies

$$(\alpha'', \beta'', \gamma'') \circ (l, f, \text{id}) = (\alpha, \beta, \gamma).$$

Then $\gamma'' = \gamma$, and we also have $\alpha'' = \alpha'$, by the universal property of L . Finally,

$$\beta''((n, g)S) = \beta''((n, 1)S) \beta''((1, g)S) = \alpha''(n) \beta''(f(g)) = \alpha''(n) \beta(g),$$

so $\beta'' = \beta'$. This proves that (l, f, id) is an L -equivalence of relative groups. The fact that f is then an L -equivalence of groups follows from Proposition 3.1. \square

Corollary 3.3 *A morphism (α, β, γ) of relative groups is an L -equivalence if and only if α is an L -equivalence of groups and γ is an isomorphism.*

PROOF. This is seen by applying relative L -localization to both the domain and the codomain of the morphism (α, β, γ) . \square

Note that the converse of Proposition 3.1 is not true in general (although it is true in some special cases, such as in [14, Theorem 2.5]). For example, if we consider the diagram

$$\begin{array}{ccccc} \mathbf{Z}/3 & \twoheadrightarrow & \Sigma_3 & \twoheadrightarrow & \mathbf{Z}/2 \\ \alpha \downarrow & & \beta \downarrow & & \text{id} \downarrow \\ 1 & \twoheadrightarrow & \mathbf{Z}/2 & \twoheadrightarrow & \mathbf{Z}/2, \end{array}$$

where Σ_3 denotes the symmetric group on three elements, and choose L to be localization at the prime 3, then β is an L -equivalence of groups, yet the diagram is not an L -equivalence of relative groups.

4 Dividing out group radicals

In this section we analyze the extent to which dividing out radicals preserves exactness or half-exactness of group extensions. Relative localization is used at a few key places.

We recall from [11] that a *radical* R in the category of groups is a subfunctor of the identity (i.e., a functor assigning to each group G a subgroup RG in such a way that every homomorphism $G \rightarrow K$ induces a homomorphism $RG \rightarrow RK$ by restriction), with the property that RG is normal in G and $R(G/RG) = 1$. The functoriality of R implies that RG is a characteristic subgroup of G for every group G ; therefore, if G embeds as a normal subgroup into some group K , then RG is also normal in K .

A radical R is *idempotent* if $RRG = RG$ for all groups G . To every radical R one can associate an idempotent radical R^∞ as follows; cf. [11, Proposition 1.2]. For each group G , let $R^\infty G$ be the product of all subgroups H of G such that $RH = H$. Then $RR^\infty G = R^\infty G$, and $R^\infty G$ is maximal with this property. We have $R^\infty G \subseteq RG$ for all groups G . In fact, R^∞ is the largest idempotent radical that is a subfunctor of R . It can alternatively be defined as the inverse limit of the following series of radicals: $R^0 = R$; $R^\alpha = RR^{\alpha-1}$ for every successor ordinal α , and R^α is the intersection of R^β

for all $\beta < \alpha$ if α is a limit ordinal. An illuminating example is $RG = [G, G]$, the commutator radical, for which $R^\infty G = \mathcal{P}G$, the perfect radical, which is the intersection of the (transfinite) derived series.

If (L, l) is any localization functor in the category of groups, then we may associate with it a radical, by defining RG to be the kernel of the localizing homomorphism $l: G \rightarrow LG$. However, such radicals need not be idempotent in general; for instance, $[G, G]$ is the kernel of the abelianization homomorphism.

A localization functor (L, l) is called an *epireflection* if, for every group G , the natural map $l: G \rightarrow LG$ is an epimorphism. If R is any radical in the category of groups, then $LG = G/RG$ defines an epireflection. Thus, there is a bijective correspondence between radicals and epireflections. An epireflection is called a *reduction* if the corresponding radical is idempotent.

Among localization functors, it is possible to characterize epireflections and reductions in terms of closure properties of the class of L -local groups. The following result, whose proof uses relative localization, generalizes Theorem 6 in [22].

Theorem 4.1 *Let (L, l) be a localization in the category of groups. Then L is an epireflection if and only if the class of L -local groups is closed under subgroups, and L is a reduction if and only if the class of L -local groups is closed under subgroups and formation of extensions.*

PROOF. Suppose that L is an epireflection. Let G be an L -local group, and S a subgroup of G . Then, by assumption, $l: S \rightarrow LS$ is surjective. Since the inclusion $S \hookrightarrow G$ factors through LS , $l: S \rightarrow LS$ is also injective, so S is L -local. Conversely, suppose that the class of L -local groups is closed under subgroups. Let G be any group and $l: G \rightarrow LG$ its localization. We may factor l as $G \twoheadrightarrow \text{im } l \hookrightarrow LG$ and, by assumption, $\text{im } l$ is L -local. Since every homomorphism from G to an L -local group factors uniquely through $\text{im } l$, the arrow $G \twoheadrightarrow \text{im } l$ is an L -equivalence. Hence, the inclusion $\text{im } l \hookrightarrow LG$ is also an L -equivalence and therefore it is an isomorphism. This shows that $l: G \rightarrow LG$ is surjective.

Now suppose that L is a reduction. This means that $LG = G/RG$ where $RRG = RG$ for all groups G . As this is an epireflection, the class of L -local groups is closed under subgroups. Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be a group extension where N and Q are L -local. Thus, $RN = 1$ and $RQ = 1$. Since RG maps into RQ , we have $RG \subseteq N$. Then $RG = RRG \subseteq RN$ implies that $RG = 1$, so G is L -local as well. Finally, suppose that the class of L -local groups is closed under subgroups and formation of extensions. Then L is an epireflection and hence $LG = G/RG$ for some radical R and all groups G . Hence, it is enough to show that RRG equals RG for all G . This is the

same as proving that $L(RG) = 1$ for all G . Take any group G and consider the relative localization

$$\begin{array}{ccccc} RG & \twoheadrightarrow & G & \twoheadrightarrow & LG \\ \wr \downarrow & & f \downarrow & & \text{id} \downarrow \\ L(RG) & \twoheadrightarrow & H & \twoheadrightarrow & LG, \end{array}$$

where, by Theorem 3.2, the map f is an L -equivalence. Since the composite of f with the surjection $H \twoheadrightarrow LG$ is an L -equivalence, we may infer that the latter is also an L -equivalence. Since we are assuming that the class of L -local groups is closed under extensions, the group H is L -local. Therefore, the surjection $H \twoheadrightarrow LG$ is an isomorphism, and this implies that $L(RG) = 1$, as needed. \square

Let R be any radical in the category of groups, and consider the epireflection given by $LG = G/RG$ for all G . In this situation, relative L -localization of any group extension $N \twoheadrightarrow G \twoheadrightarrow Q$ yields a short exact sequence

$$N/RN \twoheadrightarrow G/RN \twoheadrightarrow Q, \quad (4.1)$$

since RN is normal in G . Examples include

$$N/[N, N] \twoheadrightarrow G/[N, N] \twoheadrightarrow Q \quad \text{and} \quad N/IN \twoheadrightarrow G/IN \twoheadrightarrow Q,$$

where IN is the *isolator subgroup* of N , that is, the smallest normal subgroup of N such that N/IN is torsion-free.

Furthermore, we can consider the commutative diagram

$$\begin{array}{ccccccc} RN & \xrightarrow{Ri} & RG & \xrightarrow{Rp} & RQ \\ \downarrow & & \downarrow & & \downarrow \\ N & \xrightarrow{i} & G & \xrightarrow{p} & Q \\ \downarrow & & \downarrow & & \downarrow \\ LN & \xrightarrow{Li} & LG & \xrightarrow{Lp} & LQ \end{array}$$

where the upper and lower rows need not be exact in general, although the restriction Ri is necessarily injective, and Lp is necessarily surjective. It is also clear that the composites $Rp \circ Ri$ and $Lp \circ Li$ are trivial, by functoriality.

Proposition 4.2 *Let $N \xrightarrow{i} G \xrightarrow{p} Q$ be any group extension. Let R be any radical, and let $LG = G/RG$ be the associated epireflection. Then the following assertions hold.*

- (i) $\text{im } Ri = \ker Rp$ if and only if Li is injective.

- (ii) $\text{im } Li = \ker Lp$ if and only if Rp is surjective.
- (iii) The sequence $1 \rightarrow RN \rightarrow RG \rightarrow RQ \rightarrow 1$ is exact if and only if the sequence $1 \rightarrow LN \rightarrow LG \rightarrow LQ \rightarrow 1$ is exact.

PROOF. This is a standard diagram-chase argument. In fact, the Nine-Lemma, as stated e.g. in [23, p. 98], holds in the category of groups as well. \square

In other words, an epireflection L preserves exactness of a group extension if and only if the associated radical does. Now the following result extends [22, Proposition 5].

Theorem 4.3 *Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be any group extension and let (L, l) be an epireflection. If Q is L -local, then the sequence $LN \rightarrow LG \rightarrow Q \rightarrow 1$ is exact. Moreover, if L is a reduction, then the sequence $1 \rightarrow LN \rightarrow LG \rightarrow Q \rightarrow 1$ is exact.*

PROOF. The assumption that Q is L -local tells us that $RQ = 1$, hence the first claim, by Proposition 4.2. To prove the second statement, apply relative L -localization to the extension $N \twoheadrightarrow G \twoheadrightarrow Q$ and recall from Theorem 4.1 that the class of L -local groups for any reduction is closed under extensions. \square

5 Word radicals and plus-constructions

Let \mathcal{V} be any variety of groups [21], and denote by RG the verbal subgroup of a group G , i.e., the kernel of the projection of G onto \mathcal{V} . Then R is a radical. Radicals of this kind are called *word radicals*. Examples are $RG = [G, G]$ (the commutator subgroup); $RG = \Gamma^i G$ (the i th term of the lower central series); $RG = D^i G$ (the i th term of the derived series); or $RG = G^n$ (the subgroup generated by all n th powers). In fact, every variety can be defined by a set of words $\{x^n, c_1, c_2, c_3, \dots\}$ where each c_i is a commutator word and n is a nonnegative integer, which is called the *exponent* of the variety.

We emphasize that, if R is a word radical and $G \twoheadrightarrow Q$ is any group epimorphism, then the restriction $RG \rightarrow RQ$ is necessarily surjective. Therefore, we may use Proposition 4.2 to conclude that, for any word radical R , every group extension $N \twoheadrightarrow G \twoheadrightarrow Q$ yields a four-term exact sequence

$$1 \rightarrow (N \cap RG)/RN \rightarrow N/RN \rightarrow G/RG \rightarrow Q/RQ \rightarrow 1.$$

The perfect radical \mathcal{P} is not a word radical. For the perfect radical, a group extension $N \twoheadrightarrow G \twoheadrightarrow Q$ in which the restriction $\mathcal{P}G \rightarrow \mathcal{P}Q$ is

surjective was called an “extension preserving perfect radicals” in [2]. Thus, by Proposition 4.2, an extension $N \twoheadrightarrow G \twoheadrightarrow Q$ preserves perfect radicals if and only if the sequence

$$1 \rightarrow (N \cap \mathcal{P}G)/\mathcal{P}N \rightarrow N/\mathcal{P}N \rightarrow G/\mathcal{P}G \rightarrow Q/\mathcal{P}Q \rightarrow 1$$

is exact.

As we said in Section 4, the perfect radical may be viewed as R^∞ where R is the commutator radical. The next proposition gives sufficient conditions under which, for a radical R and a group epimorphism $G \twoheadrightarrow Q$, the restriction $R^\infty G \twoheadrightarrow R^\infty Q$ is surjective. This generalizes Proposition 2.3 from [2].

Proposition 5.1 *Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be any group extension and R a radical. Then the sequence*

$$1 \rightarrow (N \cap R^\infty G)/R^\infty N \rightarrow N/R^\infty N \rightarrow G/R^\infty G \rightarrow Q/R^\infty Q \rightarrow 1$$

is exact in each of the following cases:

- (i) *The extension $N \twoheadrightarrow G \twoheadrightarrow Q$ is split.*
- (ii) *R is a word radical and $R^m G \subseteq N \cdot R^\infty G$ for some finite m .*
- (iii) *R is a word radical and $R^n N \subseteq R^\infty G$ for some finite n .*

PROOF. Part (i) and part (ii) follow as in the proof of [2, Proposition 2.3]. In the course of the proof of (ii), one uses the fact that the restriction $R^m G \twoheadrightarrow R^m Q$ is surjective for all m , since R is a word radical. Part (iii) uses the fact that, if J is the inverse image of $R^\infty Q$ in G , then $R^n J = R^{n+1} J$, where n is the integer given by the assumption made in (iii). The details are as in [2, Lemma 2.1]. \square

Certain results about Quillen’s plus-construction remain valid if one replaces the commutator radical by any word radical R . The plus-construction with respect to a word radical was defined in [11] as follows.

For every word radical R there is a locally free group Φ_R with the property that, given any group G , the subgroup $R^\infty G$ is generated by the images of all homomorphisms $\Phi_R \rightarrow G$. We say that Φ_R *generates* or *sweeps* the radical R^∞ , as in [3]. The group Φ_R was constructed in [11, Theorem 3.3], by generalizing the technique used in [3, Example 5.3] with the perfect radical.

For each connected space X , define X_R^+ as the localization of X , in the sense of Farjoun [13], with respect to the map from an Eilenberg–Mac Lane

space $K(\Phi_R, 1)$ to a point. Thus, there is a natural map $l: X \rightarrow X_R^+$ which is a homotopy equivalence precisely when the based mapping space

$$\text{map}_*(K(\Phi_R, 1), X)$$

is weakly contractible (that is, it is path-connected and all its homotopy groups are trivial). One refers to this property by saying that X is $K(\Phi_R, 1)$ -null.

If R is the commutator radical, then X_R^+ is homotopy equivalent to Quillen's plus-construction, as shown in [3]. If R is any word radical, then the map $l: X \rightarrow X_R^+$ induces an epimorphism $\pi_1(X) \twoheadrightarrow \pi_1(X_R^+)$ factoring out precisely the radical $R^\infty \pi_1(X)$, and induces an isomorphism in homology with \mathbf{Z}/n coefficients if the variety associated with R has exponent $n > 1$; see [11, Theorem 4.1]. In particular, if R corresponds to the words x^n and $[x, y]$, then X_R^+ is a plus-construction for homology with mod n coefficients, in the sense of [20], or a partial \mathbf{Z}/n -completion as in [7, VII.6].

Proposition 5.2 *Let $N \twoheadrightarrow G \twoheadrightarrow Q$ be any group extension and R any radical. If $R^\infty Q = 1$, then the sequence $1 \rightarrow N/R^\infty N \rightarrow G/R^\infty G \rightarrow Q \rightarrow 1$ is exact.*

PROOF. This is a direct consequence of Theorem 4.3 and the fact that dividing out R^∞ is a reduction because R^∞ is idempotent. \square

The homotopy-theoretical version of Proposition 5.2 extends [1, Theorem 1.1] to arbitrary word radicals. This theorem is derived from results in [13]; see also [4].

Theorem 5.3 *Let $F \twoheadrightarrow E \twoheadrightarrow B$ be a homotopy fibre sequence of connected spaces, and let R be any word radical. If $B_R^+ \simeq B$, then $F_R^+ \twoheadrightarrow E_R^+ \twoheadrightarrow B_R^+$ is a homotopy fibre sequence.*

PROOF. Apply the plus-construction fibrewise

$$\begin{array}{ccccc} F & \longrightarrow & E & \longrightarrow & B \\ \downarrow l & & \downarrow f & & \downarrow \text{id} \\ F_R^+ & \longrightarrow & X & \longrightarrow & B \end{array}$$

and recall from [13, Theorem 1.F.3] that f induces a homotopy equivalence $E_R^+ \simeq X$. Now, if Φ_R is a locally free group sweeping the radical R^∞ , we have that B and F_R^+ are both $K(\Phi_R, 1)$ -null, so X is also $K(\Phi_R, 1)$ -null and this implies that $X_R^+ \simeq X$. \square

We end the article by showing that, if the radical R corresponds to a variety of exponent zero, then the condition that $B_R^+ \simeq B$ is equivalent to the

condition that $R^\infty \pi_1(B)$ be trivial. For varieties of nonzero exponent, this is not true, as we prove by means of a counterexample. Before displaying it, we need to analyze more closely the effect of plus-construction with respect to any radical R . Recall from [7] that an abelian group A is called Ext- p -complete, where p is a prime, if $\text{Hom}(\mathbf{Z}[1/p], A) = 0$ and $\text{Ext}(\mathbf{Z}[1/p], A) = 0$.

Theorem 5.4 *Let Φ be the direct limit of any sequence of free groups indexed by the first infinite ordinal. Then a connected space X is $K(\Phi, 1)$ -null if and only if the following conditions hold:*

- $\text{Hom}(\Phi, \pi_1(X))$ is trivial.
- $\text{Hom}(H_1(\Phi), \pi_i(X)) = 0$ and $\text{Ext}(H_1(\Phi), \pi_i(X)) = 0$ for $i \geq 2$.

PROOF. Write Φ as the direct limit of

$$F_1 \xrightarrow{\varphi_1} F_2 \xrightarrow{\varphi_2} F_3 \xrightarrow{\varphi_3} \dots \quad (5.1)$$

where each F_i is free on a set $\{x_i^j\}$ of free generators. Let F_g be the free product of all the groups F_i , and let F_r be another free group with a set of generators r_i^j corresponding bijectively with the generators x_i^j of F_g . Consider the homomorphism $\psi: F_r \rightarrow F_g$ defined as $\psi(r_i^j) = (x_i^j)^{-1} \varphi_i(x_i^j)$ for all i, j . Thus, the normal closure N of the image of ψ in F_g is the kernel of a free presentation of Φ ,

$$N \twoheadrightarrow F_g \twoheadrightarrow \Phi,$$

yielding, by abelianization, a sequence

$$F_r/[F_r, F_r] \twoheadrightarrow F_g/[F_g, F_g] \twoheadrightarrow \Phi/[\Phi, \Phi] \quad (5.2)$$

which is a free abelian presentation of the abelianization of Φ , hence exact.

In homotopy-theoretical terms, we may consider a wedge of circles W_i with fundamental group F_i , for every i , and consider the homotopy colimit of the sequence of maps corresponding to (5.1). This space is a 2-dimensional $K(\Phi, 1)$, since it is an ascending union of 2-dimensional $K(F_i, 1)$ spaces. (In the case of sequences indexed by ordinals bigger than the first infinite ordinal, the homotopy colimit could fail to be 2-dimensional.) The cell decomposition of this homotopy colimit is described by a homotopy cofibre sequence

$$W_r \rightarrow W_g \rightarrow K(\Phi, 1), \quad (5.3)$$

inducing precisely (5.2) on H_1 . The spaces W_r and W_g are wedges of circles with fundamental groups F_r and F_g , respectively.

Now a connected space X is $K(\Phi, 1)$ -null if and only if the sets of pointed homotopy classes of maps $[\Sigma^n K(\Phi, 1), X]$ are trivial for all $n \geq 0$. Hence, the long exact sequence

$$\begin{aligned} \cdots \rightarrow [\Sigma K(\Phi, 1), X] &\rightarrow [\Sigma W_g, X] \rightarrow [\Sigma W_r, X] \\ &\rightarrow [K(\Phi, 1), X] \rightarrow [W_g, X] \rightarrow [W_r, X], \end{aligned}$$

obtained from (5.3) proves the statement, since

$$[\Sigma^n W_g, X] \cong [W_g, \Omega^n X] \cong \text{Hom}(F_g, \pi_{n+1}(X))$$

for all n , and similarly with W_r . \square

Corollary 5.5 *Let R be a word radical associated with a variety \mathcal{V} , and let X be a connected space.*

- (a) *If \mathcal{V} has exponent zero, then $X_R^+ \simeq X$ if and only if $R^\infty \pi_1(X)$ is trivial.*
- (b) *If \mathcal{V} has a prime exponent p , then $X_R^+ \simeq X$ if and only if $R^\infty \pi_1(X)$ is trivial and, for each $i \geq 2$, the group $\pi_i(X)$ is Ext- p -complete.*

PROOF. Let Φ_R be a locally free group sweeping R^∞ , constructed as in [11, Theorem 3.3]. We recall that Φ_R is a free product of countable locally free groups, which is perfect if \mathcal{V} has exponent zero, and

$$H_1(\Phi_R) \cong \bigoplus_{i \in I} \mathbf{Z}[1/p] \tag{5.4}$$

for some set of indices I if \mathcal{V} has exponent p . Since $X_R^+ \simeq X$ if and only if X is $K(\Phi_R, 1)$ -null, both statements (a) and (b) are consequences of Theorem 5.4. \square

Corollary 5.6 *Let R be a word radical corresponding to a variety of exponent p . Let X be a connected space such that $\pi_1(X)$ is abelian and p -divisible. Then X_R^+ is homotopy equivalent to the p -completion of X .*

PROOF. It follows from Theorem 5.4 and the isomorphism (5.4) that, if $\pi_1(X)$ is abelian, then X is $K(\Phi_R, 1)$ -null if and only if it is $K(\mathbf{Z}[1/p], 1)$ -null, since $\mathbf{Z}[1/p]$ is also locally free. Hence, X_R^+ is homotopy equivalent to the localization of X with respect to the map $K(\mathbf{Z}[1/p], 1) \rightarrow *$. The assumption that $\pi_1(X)$ is p -divisible is equivalent to imposing that $\mathbf{Z}[1/p]$

sweeps $\pi_1(X)$, and this guarantees that X_R^+ is homotopy equivalent to the p -completion of X , by [10, Theorem 4.4]. \square

Using this result, we show that the assumption that $B_R^+ \simeq B$ cannot be replaced with the assumption that $R^\infty\pi_1(B) = 1$ in Theorem 5.3 if the variety associated with R has nonzero exponent. The homotopy fibre sequence

$$K(\mathbf{Z}, 1) \longrightarrow K(\mathbf{Z}/2, 1) \longrightarrow K(\mathbf{Z}, 2)$$

corresponding to a nonzero element in $H^2(\mathbf{Z}/2; \mathbf{Z})$ is transformed by the plus-construction with respect to any word radical of exponent p , where p is any odd prime, into the sequence

$$K(\mathbf{Z}, 1) \longrightarrow * \longrightarrow K(\widehat{\mathbf{Z}}_p, 2),$$

where $\widehat{\mathbf{Z}}_p$ denotes the p -adic integers. This is of course not a homotopy fibre sequence.

References

- [1] A. J. Berrick, The plus-construction and fibrations, *Quart. J. Math. Oxford* (2) **33** (1982), 149–157.
- [2] A. J. Berrick, Group epimorphisms preserving perfect radicals, and the plus-construction, in: *Algebraic K-Theory, Lecture Notes in Math.* vol. 1046, Springer-Verlag, Berlin Heidelberg New York, 1984, 1–12.
- [3] A. J. Berrick and C. Casacuberta, A universal space for plus-constructions, *Topology* **38** (1999), 467–477.
- [4] A. J. Berrick and E. Dror Farjoun, A characterization of fibrations preserved by certain localizations, preprint.
- [5] I. Bokor and P. Hilton, A push-out construction for groups with operators, *Exposition. Math.* **17** (1999), 143–159.
- [6] A. K. Bousfield, Localization and periodicity in unstable homotopy theory, *J. Amer. Math. Soc.* **7** (1994), 831–873.
- [7] A. K. Bousfield and D. M. Kan, *Homotopy Limits, Completions and Localizations*, *Lecture Notes in Math.* vol. 304, Springer-Verlag, Berlin Heidelberg New York, 1972.

- [8] C. Casacuberta, On structures preserved by idempotent transformations of groups and homotopy types, in: Crystallographic Groups and Their Generalizations, Contemp. Math. vol. 262, Amer. Math. Soc., Providence, 2000, 39–68.
- [9] C. Casacuberta and M. Castellet, Localization methods in the study of the homology of virtually nilpotent groups, Math. Proc. Cambridge Philos. Soc. **112** (1992), 551–564.
- [10] C. Casacuberta and J. L. Rodríguez, On towers approximating homological localizations, J. London Math Soc. (2) **56** (1997), 645–656.
- [11] C. Casacuberta, J. L. Rodríguez and D. Scevenels, Singly generated radicals associated with varieties of groups, in: Groups St Andrews 1997 in Bath I, London Math. Soc. Lecture Note Ser. vol. 260, Cambridge University Press, Cambridge, 1999, 202–210.
- [12] A. Descheemaeker and W. Malfait, P -localization of relative groups, J. Pure Appl. Algebra **159** (2001), 43–56.
- [13] E. Dror Farjoun, Cellular Spaces, Null Spaces and Homotopy Localization, Lecture Notes in Math. vol. 1622, Springer-Verlag, Berlin Heidelberg New York, 1996.
- [14] P. Hilton, Relative nilpotent groups, in: Categorical Aspects of Topology and Analysis, Lecture Notes in Math. vol. 915, Springer-Verlag, Berlin Heidelberg New York, 1982, 136–147.
- [15] P. Hilton, Localization of crossed modules, in: Séminaire d’Algèbre Paul Dubreil et Marie-Paule Malliavin, Lecture Notes in Math. vol. 1029, Springer-Verlag, Berlin Heidelberg New York, 1983, 311–324.
- [16] I. Llerena, Localization of fibrations with nilpotent fibre, Math. Z. **188** (1985), 397–410.
- [17] S. Mac Lane, Homology, Grund. math. Wiss. Einz. Band 114, Springer-Verlag, Berlin Heidelberg New York, 1975.
- [18] S. Mac Lane, Categories for the Working Mathematician, Graduate Texts in Math. vol. 5, Springer-Verlag, Berlin Heidelberg New York, 1975.
- [19] J. P. May, Fibrewise localization and completion, Trans. Amer. Math. Soc. **258** (1980), 127–146.
- [20] W. Meier and R. Strebel, Homotopy groups of acyclic spaces, Quart. J. Math. Oxford Ser. (2) **32** (1981), 81–95.
- [21] H. Neumann, Varieties of Groups, Ergeb. Math. Grenzgeb. Band 37, Springer-Verlag, Berlin Heidelberg New York, 1967.

- [22] J. L. Rodríguez and D. Scevenels, Universal epimorphic equivalences for group localizations, *J. Pure Appl. Algebra* **148** (2000), 309–316.
- [23] J. R. Strooker, *Introduction to Categories, Homological Algebra and Sheaf Cohomology*, Cambridge University Press, Cambridge, 1978.
- [24] D. Sullivan, Genetics of homotopy theory and the Adams conjecture, *Ann. of Math.* **100** (1974), 1–79.

Departament d'Àlgebra i Geometria, Universitat de Barcelona,
Gran Via de les Corts Catalanes, 585; E-08007 Barcelona, Spain
e-mail: casac@mat.ub.es

Centre de Recerca Matemàtica, Institut d'Estudis Catalans
Apartat 50; E-08193 Bellaterra, Spain
e-mail: adesche@crm.es

and

Department of Mathematics, Katholieke Universiteit Leuven Campus Kor-
trijk
Universitaire Campus; B-8500 Kortrijk, Belgium
e-mail: An.Descheemaeker@kulak.ac.be