

Fixed subgroups are compressed in free groups

A. Martino

Centre de Recerca Matemàtica
Barcelona, Spain
A.Martino@crm.es

E. Ventura

Dept. Mat. Apl. III,
Universitat Politècnica de Catalunya,
Barcelona, Spain
enric.ventura@upc.es

Abstract

In this paper we prove that the fixed subgroup of an arbitrary family of endomorphisms ψ_i , $i \in I$, of a finitely generated free group F , is F -super-compressed. In particular, $r(\cap_{i \in I} \text{Fix } \psi_i) \leq r(M)$ for every subgroup $M \leq F$ containing $\cap_{i \in I} \text{Fix } \psi_i$. This provides new evidence towards the inertia conjecture for fixed subgroups of free groups. As a corollary, we show that, in the free group of rank n , every strictly ascending chain of fixed subgroups has length at most $2n$.

1 Introduction

Let F be a finitely generated free group, and let endomorphisms of F act on the right, $x \mapsto x\psi$.

Given an endomorphism ψ of F , its *fixed subgroup*, denoted $\text{Fix } \psi$, is the subgroup of elements in F fixed by ψ , $\text{Fix } \psi = \{x \in F \mid x\psi = x\}$. If $\Psi = \{\psi_i \mid i \in I\}$ is a family of endomorphisms of F , we denote by $\text{Fix } \Psi$ the subgroup of elements simultaneously fixed by all ψ_i , $\text{Fix } \Psi = \cap_{i \in I} \text{Fix } \psi_i$.

Following the terminology introduced in [11], we say that a subgroup $H \leq F$ is *1-endo-fixed* if $H = \text{Fix } \psi$ for some endomorphism $\psi \in \text{End}(F)$. And we say that H is *endo-fixed* if $H = \text{Fix } \Psi$ for some family of endomor-

phisms $\Psi \subseteq \text{End}(F)$. The concepts of *1-auto-fixed*, *1-mono-fixed*, *auto-fixed* and *mono-fixed* subgroups are analogously defined.

Bestvina-Handel [2] proved that, if $\psi: F \rightarrow F$ is an automorphism, then $r(\text{Fix } \psi) \leq r(F)$, improving the previously celebrated result of Gersten [7] about finite generation of such fixed subgroups. Immediately after Bestvina and Handel announced their result, Imrich-Turner published [8], where they extended it to arbitrary endomorphisms. Some years latter, Dicks-Ventura [5] introduced the concept of inertia and showed that the fixed subgroup of an injective endomorphism of F is F -inert. A subgroup $H \leq F$ is called F -inert when $r(H \cap K) \leq r(K)$ for every $K \leq F$. This result was another extension of the Bestvina-Handel Theorem. Since the family of F -inert subgroups is closed under intersections, an easy corollary was that the rank of an arbitrary mono-fixed subgroup of F is also bounded above by the rank of F . However, in [5] problem 2, it was asked if fixed subgroups of arbitrary endomorphisms of F are necessarily F -inert, and in [18] it was conjectured that, in fact, they are. This is still open and we will refer to it as the “inertia conjecture”.

Using inertia, Bergman [1] showed that the rank of an arbitrary endo-fixed subgroup of F is always bounded above by the rank of F , even without the injectivity hypothesis on the involved endomorphisms. This was the first evidence in favour of the inertia conjecture.

In [5] the concept of compression of a subgroup of F was introduced, being a necessary condition for its inertia. We re-estate it here along with some technical variations.

Let $F^{\text{ab}} = F/F'$ be the abelianization of F and let $\pi: F \rightarrow F^{\text{ab}}$ be the corresponding projection. For any given subgroup $H \leq F$, the *abelian rank of H with respect to F* , denoted $r^{\text{ab}}(H; F)$, is defined as the rank of the free abelian group $H\pi = HF'/F'$, that is, the rank (as an abelian group) of the image of H under the global abelianization π . Note that, in general, this is not the same as $r(H^{\text{ab}})$.

Definition 1.1 Let F be a free group and $H \leq F$.

We say that H is *F -compressed* if $r(H) \leq r(K)$ for every $H \leq K \leq F$.

Similarly, H is called *F -strictly-compressed* when $r(H) < r(K)$ for every $H < K \leq F$.

Finally, H is called *F -super-compressed* if for any subgroup $H < K \leq F$, one has both $r(H) \leq r(K)$ and $r(H) + r^{\text{ab}}(H; F) < r(K) + r^{\text{ab}}(K; F)$. Equivalently, H is *F -super-compressed* if H is F -compressed and, for every $K \leq F$ strictly containing H but having the same rank, $r^{\text{ab}}(H; F) < r^{\text{ab}}(K; F)$ is satisfied.

Clearly, every F -strictly compressed and every F -super-compressed subgroups of F are F -compressed. Also, every F -inert subgroup of F is F -compressed, but it is not known if the converse is true. This was stated in [5] as Problem 1, and in [18] as the “compressed-inert conjecture”.

The main result in the present paper is Theorem 3.3, where a simple argument (making use of theorems of Bergman, Takahasi, Martino-Ventura, Bestvina-Handel and Dyer-Scott) is given to prove that arbitrary endo-fixed subgroups of F are F -super-compressed. In particular, they are F -compressed, providing new evidence in support of the inertia conjecture.

For this purpose, we will also make use of the concept of algebraic extension. Following [9], a pair of subgroups $H \leq K$ of F is called an *algebraic extension* if H is not contained in any proper free factor of K . Note that, in general, if $\psi: F \rightarrow F$ is an endomorphism and $H \leq K$ is an algebraic extension, then $H\psi \leq K\psi$ need not be algebraic (while in fact it is, if ψ is an automorphism). A result of Takahasi [15] states that any finitely generated subgroup H of F has a finite number of algebraic extensions, i.e. there are only finitely many subgroups $K \leq F$ such that H is contained in K but not in any of its proper free factors. Simpler arguments for this result were recently given independently by Ventura [17], Margolis-Sapir-Weil [10] and Kapovich-Myasnikov [9]. This result will be crucial in the proof of our main result.

The structure of the present paper is the following. In section 2 we prove some general properties of the three concepts of compression for subgroups of free groups. In section 3 the main result (Theorem 3.3), namely the F -super-compression of endo-fixed subgroups of F , is proven. Finally, in section 4 we apply the results to better understanding ascending chains of endo-fixed subgroups of free groups.

2 Properties of compression

In this section we prove that the three concepts of compression behave well with respect to free products.

Lemma 2.1 *Let U, V be finitely generated subgroups of a free abelian group. Then,*

$$r(U + V) = r(U) + r(V) - r(U \cap V).$$

Proof. Noting that $r(U) = \dim_{\mathbb{Q}}(U \otimes_{\mathbb{Z}} \mathbb{Q})$, the formula follows from the corresponding result in linear algebra. \square

A subgroup $R \leq F$ is called a *retract* of F when the identity $Id: R \rightarrow R$ extends to a homomorphism $\rho: F \rightarrow R$, called a *retraction*. For example, free factors of F are retracts of F .

Lemma 2.2 *Let F be a free group and let $R \leq F$ be a retract of F . For every $H \leq R$, $r^{\text{ab}}(H; R) = r^{\text{ab}}(H; F)$.*

Proof. The equality between the abelian ranks with respect to R and F is clear if we show $F' \cap R = R'$. One of the inclusions is obvious. To show the other, let $\rho: F \rightarrow R$ be a retraction and note that if the commutator of $x, y \in F$ lies in R then $[x, y] = [x, y]\rho = [x\rho, y\rho] \in R'$, since $x\rho, y\rho \in R$. Thus, $F' \cap R \leq R'$. \square

Proposition 2.3 *Let F be a free group and $H = A * B$ a finitely generated subgroup.*

- i) If H is F -compressed then A is also F -compressed.*
- ii) If H is F -strictly-compressed then A is also F -strictly-compressed.*
- iii) If H is F -super-compressed then A is also F -super-compressed.*

Proof. Let L be an arbitrary subgroup of F containing A . Consider $K = \langle L, B \rangle$, which is a subgroup of F containing H , and having rank $r(K) \leq r(L) + r(B)$.

Suppose that H is F -compressed. Then,

$$r(A) + r(B) = r(H) \leq r(K) \leq r(L) + r(B).$$

So, $r(A) \leq r(L)$. This proves (i).

Now, suppose that H is F -strictly-compressed. By (i), A is F -compressed and hence, to see (ii), it only remains to show that $r(A) < r(L)$ whenever $A < L$. Suppose then that $A < L$. If $A < L \leq A * B$ then A is a proper free factor of L and we are done. Otherwise, there exists $x \in L$, $x \notin A * B$. Thus $H < K$ and, by hypothesis,

$$r(A) + r(B) = r(H) < r(K) \leq r(L) + r(B).$$

So, $r(A) < r(L)$. This completes (ii).

Finally, suppose that H is F -super-compressed. By (i), A is F -compressed and hence, to see (iii), it only remains to show that $r^{\text{ab}}(A; F) < r^{\text{ab}}(L; F)$ whenever $A < L$ and $r(A) = r(L)$. So, suppose that $A < L$ and $r(A) = r(L)$.

Note that, by the equality of ranks, L cannot be contained in $A * B$ and so, $H < K$. Also,

$$r(H) = r(A) + r(B) = r(L) + r(B) \geq r(K).$$

Hence, by the hypothesis, $r(H) = r(K)$ and $r^{\text{ab}}(H; F) < r^{\text{ab}}(K; F)$. Now, using Lemma 2.1,

$$r(K\pi) = r(L\pi) + r(H\pi) - r(L\pi \cap H\pi),$$

where $\pi: F \rightarrow F^{\text{ab}}$ denotes the abelianization map. But $A \leq L \cap H$, so

$$\begin{aligned} r^{\text{ab}}(K; F) &\leq r^{\text{ab}}(L; F) + r^{\text{ab}}(H; F) - r^{\text{ab}}(A; F) \\ &< r^{\text{ab}}(L; F) + r^{\text{ab}}(K; F) - r^{\text{ab}}(A; F). \end{aligned}$$

Hence, $r^{\text{ab}}(A; F) < r^{\text{ab}}(L; F)$. This completes (iii). \square

Proposition 2.4 *Let F be a free group, and $A \leq H \leq F$, $B \leq K \leq F$ and $M \leq F$ be subgroups such that $F = H * K * M$.*

- i) If A is H -compressed and B is K -compressed then $A * B$ is F -compressed.*
- ii) If A is H -strictly-compressed and B is K -strictly-compressed then $A * B$ is F -strictly-compressed.*
- iii) If A is H -super-compressed and B is K -super-compressed then $A * B$ is F -super-compressed.*

Proof. Let $A * B \leq L \leq F = H * K * M$. Writing $A \leq L_A = L \cap H \leq H$ and $B \leq L_B = L \cap K \leq K$, the Kurosh Subgroup Theorem ensures us the existence of a subgroup $L' \leq F$ such that $L = L_A * L_B * L'$.

Assume the hypothesis in (i). Then, $r(A) \leq r(L_A)$, $r(B) \leq r(L_B)$ and so,

$$r(A * B) = r(A) + r(B) \leq r(L_A) + r(L_B) + r(L') = r(L).$$

This proves (i).

The same argument works to prove (ii), with the additional observation that if $A * B < L$ then either $A < L_A$, or $B < L_B$ or $L' \neq 1$. Thus, by the hypothesis in (ii), either $r(A) < r(L_A)$, or $r(B) < r(L_B)$ or $L' \neq 1$. In this situation, the inequality in the above computation is strict, that is, $r(A * B) < r(L)$.

Finally, assume the hypothesis in (iii). Using (i) it only remains to prove that if $A * B < L$ and $r(A * B) = r(L)$ then $r^{\text{ab}}(A * B; F) < r^{\text{ab}}(L; F)$.

In this situation, $r(A) = r(L_A)$, $r(B) = r(L_B)$ and $L' = 1$. But either $A < L_A$, or $B < L_B$ so, by the hypothesis, either $r^{\text{ab}}(A; H) < r^{\text{ab}}(L_A; H)$ or $r^{\text{ab}}(B; K) < r^{\text{ab}}(L_B; K)$. Thus, using Lemma 2.2,

$$\begin{aligned}
r^{\text{ab}}(A * B; F) &= r^{\text{ab}}(A; F) + r^{\text{ab}}(B; F) \\
&= r^{\text{ab}}(A; H) + r^{\text{ab}}(B; K) \\
&< r^{\text{ab}}(L_A; H) + r^{\text{ab}}(L_B; K) \\
&= r^{\text{ab}}(L_A; F) + r^{\text{ab}}(L_B; F) \\
&= r^{\text{ab}}(L_A * L_B; F) \\
&= r^{\text{ab}}(L; F). \quad \square
\end{aligned}$$

3 Compression of fixed subgroups

In [12], Martino-Ventura gave an explicit description of 1-auto-fixed subgroups of finitely generated free groups. We state it here for later use.

Theorem 3.1 (Martino-Ventura, [12]) *Let F be a non-trivial finitely generated free group and let $\psi \in \text{Aut}(F)$ with $\text{Fix } \psi \neq 1$. Then, there exist integers $r \geq 1$, $s \geq 0$, ψ -invariant non-trivial subgroups $K_1, \dots, K_r \leq F$, primitive elements $y_1, \dots, y_s \in F$, a subgroup $L \leq F$, and elements $1 \neq h'_j \in H_j = K_1 * \dots * K_r * \langle y_1, \dots, y_j \rangle$, $j = 0, \dots, s-1$, such that*

$$F = K_1 * \dots * K_r * \langle y_1, \dots, y_s \rangle * L$$

and $y_j \psi = h'_{j-1} y_j$ for $j = 1, \dots, s$; moreover,

$$\text{Fix } \psi = \langle w_1, \dots, w_r, y_1^{-1} h_0 y_1, \dots, y_s^{-1} h_{s-1} y_s \rangle$$

for some non-proper powers $1 \neq w_i \in K_i$ and $1 \neq h_j \in H_j$ such that $h_j \psi = h'_j h_j h'^{-1}_j$, $i = 1, \dots, r$, $j = 0, \dots, s-1$. \square

Using Theorem 3.1 and the standard covering theory for graphs (see [14], [17] or [9]), we can extend Theorem 2.2 in [17] to arbitrary 1-auto-fixed subgroups of F . Namely, we will show that for any given automorphism $\psi: F \rightarrow F$, every subgroup of F strictly containing $\text{Fix } \psi$ has either bigger rank or bigger abelian rank than those of $\text{Fix } \psi$. After this, we will have all the ingredients to prove our main result, stating the F -super-compression of any endo-fixed subgroup of F .

Proposition 3.2 *Let F be a finitely generated free group and let $\psi: F \rightarrow F$ be an automorphism. Then, for every K with $\text{Fix } \psi < K \leq F$, either $r(\text{Fix } \psi) < r(K)$ or $r^{\text{ab}}(\text{Fix } \psi; F) < r^{\text{ab}}(K; F)$.*




Figura 1.

Proof. The result is clear if $\text{Fix } \psi = 1$. So, assume $\text{Fix } \psi \neq 1$.

Consider the description of $\text{Fix } \psi$ given in Theorem 3.1. Now, the argument works exactly as the proof given for Theorem 2.2 in [17]. The core graph Z corresponding to $\text{Fix } \psi$, with base point u and the corresponding labels on edges and circles, is depicted in Fig. 1. Let E' be the set of edges in Z whose deletion disconnects the graph; for $j = 1, \dots, s$ denote by e_j the edge in E' with label y_j . Observe that, since $h_{j-1} \in K_1 * \dots * K_r * \langle y_1, \dots, y_{j-1} \rangle$, none of the edges in the component of $Z - E'$ containing e_j has label y_j .

By Theorem 1.7 in [17], it is enough to prove that any proper extension $\text{Fix } \psi < K$ given by a quotient \bar{Z} of Z has either bigger rank or bigger abelian rank than those of $\text{Fix } \psi$. Fix one such K (and the corresponding $\bar{Z} \neq Z$), and let J be the set of those j such that some vertex in the component of $Z - E'$ containing e_j gets identified with some vertex of a different component of $Z - E'$.

Suppose $J \neq \emptyset$, and let j_0 be the largest element of J . Clearly, there is a closed path in the quotient graph \bar{Z} based at \bar{u} which crosses exactly one edge labelled y_{j_0} , and only once. This closed path determines an element in KF'/F' belonging to a complement of $(\text{Fix } \psi)F'/F'$ in F^{ab} . Thus, $r^{\text{ab}}(\text{Fix } \psi; F) < r^{\text{ab}}(K; F)$.

This leaves the case where $J = \emptyset$, that is, where vertices in different components of $Z - E'$ remain unidentified in $\bar{Z} \neq Z$. In this case, \bar{Z} is the same as Z with every component of $Z - E'$ replaced by some quotient of itself (with at least one of those quotients being proper). For every $i = 1, \dots, r$ and $j = 0, \dots, s - 1$, the elements w_i and h_j are not proper powers, so the subgroups $\langle w_i \rangle$ and $\langle h_j \rangle$ are F -strictly compressed. Thus, by

Proposition 2.4 (ii), $\langle w_1, \dots, w_r \rangle$ is F -strictly compressed too. We deduce that $r(\text{Fix } \psi) < r(K)$. \square

Note that Proposition 3.2 is not saying that 1-auto-fixed subgroups of F are F -compressed, since the statement leaves the possibility of the existence of some subgroup $\text{Fix } \psi < K \leq F$ with bigger abelian rank but smaller rank than those of $\text{Fix } \psi$ (in fact, in the case $J \neq \emptyset$ we have no control on the rank of the graph \overline{Z}). In the next theorem we give an argument showing, in particular, this compression.

Theorem 3.3 *Let F be a finitely generated free group. Any endo-fixed subgroup of F is F -super-compressed.*

Proof. Let $\Psi \subseteq \text{End}(F)$. First, we will prove that $\text{Fix } \Psi$ is F -compressed. Then, using Proposition 3.2, we will extend this to show that it is F -super-compressed.

By [7], $\text{Fix } \Psi$ is finitely generated. Let r be the minimum among the ranks of all those subgroups of F containing $\text{Fix } \Psi$. Note that $r \leq r(\text{Fix } \Psi)$, and the equality holds if and only if $\text{Fix } \Psi$ is F -compressed.

Consider $\mathcal{M} = \{M \leq F \mid \text{Fix } \Psi \leq M, r(M) = r\} \neq \emptyset$. Observe that, by the minimality of r , every $M \in \mathcal{M}$ is an algebraic extension of $\text{Fix } \Psi$. Hence, by [9], [10], [15] or [17], $|\mathcal{M}| < \infty$.

By [11] Corollary 3.4, there exists φ (in the submonoid of $\text{End}(F)$ generated by Ψ) such that $\text{Fix } \Psi$ is a free factor of $\text{Fix } \varphi$. Furthermore, it is easy to see that, for every positive integer s , φ restricts to a finite order automorphism of $\text{Fix } \varphi^s$, whose fixed subgroup is $\text{Fix } \varphi$ itself. So, by a result of Dyer-Scott [6], $\text{Fix } \varphi$ is a free factor of $\text{Fix } \varphi^s$. Hence, $\text{Fix } \Psi$ is a free factor of $\text{Fix } \varphi^s$ for every $s \geq 1$.

Now choose an arbitrary $M \in \mathcal{M}$. Since the rank of a subgroup never increases when taking images, it is clear that $M_k = M\varphi^k \in \mathcal{M}$, for every $k \geq 0$. By the finiteness of \mathcal{M} , there exists an integer $k \geq 0$ and a positive integer $s \geq 1$ such that $M_{k+s} = M_k\varphi^s = M_k$. Since $r(M_{k+s}) = r(M_k)$, φ^s restricts to an automorphism of M_k , say $\varphi_{M_k}^s \in \text{Aut}(M_k)$. Now, using Bestvina-Handel Theorem,

$$r(M_k \cap \text{Fix } \varphi^s) = r(\text{Fix } \varphi_{M_k}^s) \leq r(M_k) = r(M) = r.$$

But we noted above that $\text{Fix } \Psi$ is a free factor of $\text{Fix } \varphi^s$. Hence, it is also a free factor of $M_k \cap \text{Fix } \varphi^s$. Thus, $r(\text{Fix } \Psi) \leq r$ that is, $r(\text{Fix } \Psi) = r$. This means that $\text{Fix } \Psi$ is F -compressed.

It remains to show $r^{\text{ab}}(\text{Fix } \Psi; F) < r^{\text{ab}}(M; F)$, assuming $M \neq \text{Fix } \Psi$.

In [16], Turner showed that the *stable image* of φ^s ,

$$R = \bigcap_{i \geq 1} F(\varphi^s)^i = \bigcap_{i \geq 1} F\varphi^i,$$

is a retract of F where φ^s restricts to an automorphism. Let $\varphi_R^s \in \text{Aut}(R)$ denote the restriction of φ^s to R . Observe that $\text{Fix } \varphi_R^s = \text{Fix } \varphi^s$ since $\text{Fix } \varphi^s \leq R$.

Having proved that endo-fixed subgroups of F are F -compressed, and using Proposition 3.2, we see that $\text{Fix } \varphi^s = \text{Fix } \varphi_R^s$ is R -super-compressed. Recall that $\text{Fix } \Psi$ is a free factor of $\text{Fix } \varphi^s$ and hence, by Proposition 2.3 (iii), $\text{Fix } \Psi$ is also R -super-compressed.

Now, observe that M_k cannot be equal to $\text{Fix } \Psi$. For, if it were, then as φ acts as the identity on $\text{Fix } \Psi$ which is a proper subgroup of M , the map $\varphi^k : M \rightarrow M_k$ would be a surjective map with non-trivial kernel between two free groups of the same rank. As finitely generated free groups are Hopfian, this cannot be the case and hence $\text{Fix } \Psi < M_k$.

But, since $r(\text{Fix } \Psi) = r = r(M_k)$ and $M_k \leq R$, the R -super-compression of $\text{Fix } \Psi$ implies that

$$r^{\text{ab}}(\text{Fix } \Psi; R) < r^{\text{ab}}(M_k; R)$$

and, by Lemma 2.2,

$$r^{\text{ab}}(\text{Fix } \Psi; F) < r^{\text{ab}}(M_k; F).$$

Finally, since $\varphi^k : M \rightarrow M_k$ induces a surjective homomorphism from MF'/F' to M_kF'/F' , we have

$$r^{\text{ab}}(\text{Fix } \Psi; F) < r^{\text{ab}}(M_k; F) \leq r^{\text{ab}}(M; F).$$

This completes the proof. \square

Corollary 3.4 *Let F be a finitely generated free group. Any endo-fixed subgroup of F is F -compressed.*

4 Ascending chains of fixed subgroups

In this last section, we use the information obtained before to analyze strictly ascending chains of endo-fixed subgroups of a finitely generated free group.

Theorem 4.1 *In a free group F of rank n , every strictly ascending chain of endo-fixed subgroups has length at most $2n$.*

Furthermore, there exist such chains of length $2n - 1$, even using only 1-auto-fixed subgroups.

Proof. By the above Theorem 3.3, the function $r(-) + r^{\text{ab}}(-; F)$ strictly increases in any step of such an ascending chain. And, clearly, its minimum and maximum values among endo-fixed subgroups of F , are 0 (for the trivial subgroup) and $2n$ (for F itself), respectively. So, strictly ascending chains of endo-fixed subgroups of F have length at most $2n$.

Now, we need to construct such a chain with length $2n - 1$, and using only 1-auto-fixed subgroups. Let $\{x_1, \dots, x_n\}$ be a basis of F . Given two integers $1 \leq p \leq q \leq n$, let $r, s, t \geq 0$ be such that $p = r$, $q = r + s$ and $n = r + s + t$. Define $\psi_{p,q}$ to be the automorphism of F given by

$$\begin{aligned} \psi_{p,q}: F &\mapsto F \\ x_i &\mapsto x_i, & i = 1, \dots, r \\ x_j &\mapsto x_1^j x_j, & j = r + 1, \dots, r + s \\ x_k &\mapsto x_k^{-1}, & k = r + s + 1, \dots, r + s + t. \end{aligned}$$

It is not difficult to see that

$$\text{Fix } \psi_{p,q} = \langle x_1, \dots, x_r, x_{r+1}^{-1} x_1 x_{r+1}, \dots, x_{r+s}^{-1} x_1 x_{r+s} \rangle.$$

We have $r^{\text{ab}}(\text{Fix } \psi_{p,q}; F) = r = p$ and $r(\text{Fix } \psi_{p,q}) = r + s = q$. The following is a strictly ascending chain of 1-auto-fixed subgroups with length $2n - 1$:

$$1 < \text{Fix } \psi_{1,1} < \dots < \text{Fix } \psi_{1,n} < \text{Fix } \psi_{2,n} < \dots < \text{Fix } \psi_{n,n} = F. \quad \square$$

The previous result leaves the four questions whether the length of the longest strictly ascending chains of 1-auto-fixed, 1-endo-fixed, auto-fixed or endo-fixed subgroups of the free group of rank n , is either $2n - 1$ or $2n$. For $n = 2$ the four questions coincide since the four families of subgroups do coincide (see Theorem 3.9 in [17]). For $n \geq 3$, the families of 1-endo-fixed and 1-auto-fixed subgroups are known to be different (see [13]), while in [11] the families of 1-auto-fixed and auto-fixed subgroups are conjectured to coincide. So, in general, the four questions are not the same.

In the following proposition we show that, in the cases $n = 2$ and $n = 3$, and for 1-auto-fixed subgroups, the exact maximum length is $2n - 1$. In the remark below we point out a reason why it seems difficult to extrapolate the arguments given to the general case.

Let F be a free group of rank $n \geq 1$.

Let $0 \leq p \leq q \leq n$ be two integers. A subgroup $H \leq F$ is said to be of *type* (p, q) if $r^{\text{ab}}(H; F) = p$ and $r(H) = q$. And, given also $0 \leq p' \leq q' \leq n$, an inclusion $H < K$ of subgroups of F is said to be of *type* “ $(p, q) < (p', q')$ ” when H is of type (p, q) and K is of type (p', q') . Note that, by Theorem 3.3,

any strict inclusion of endo-fixed subgroups is of type “ $(p, q) < (p', q')$ ”, where $p \leq p'$, $q \leq q'$, and at least one of these two inequalities is strict.

By Corollary 3.4 in [11], Corollary 2 in [16], and Proposition 1 in [3], any endo-fixed subgroup of F with rank n is automatically 1-auto-fixed, and contains a primitive element. Thus, no endo-fixed subgroup of F is of type $(0, n)$. Hence, the inclusion types “ $(0, n-1) < (0, n)$ ” and “ $(0, n) < (1, n)$ ” do not appear in any ascending chain of endo-fixed subgroups. Clearly, the type “ $(0, 1) < (1, 1)$ ” is also impossible there, because any endomorphism fixing a power of an element, fixes the element itself. We will see below that all the other possible inclusion types can be realized by 1-auto-fixed subgroups.

Proposition 4.2 *For $n = 2$ and $n = 3$, the length of the longest strictly ascending chains of 1-auto-fixed subgroups of the free group of rank n , is $2n - 1$.*

Proof. Let F be a free group of rank n . We already know the existence of strictly ascending chains of 1-auto-fixed subgroups of F with length $2n - 1$. Assume now that $H_0 < H_1 < \dots < H_{2n}$ is such a chain with length $2n$ and we will reach a contradiction when $n = 2$ or $n = 3$.

Since in any inclusion the value of $r(-) + r^{\text{ab}}(-; F)$ increases, the length of the chain forces H_0 to be of type $(0, 0)$, H_{2n} of type (n, n) , and every inclusion in the chain of type either “ $(p, q) < (p+1, q)$ ” or “ $(p, q) < (p, q+1)$ ” for some p, q .

Suppose $n = 2$. The impossibility of the inclusion types “ $(0, 1) < (1, 1)$ ”, “ $(0, 1) < (0, 2)$ ” and “ $(0, 2) < (1, 2)$ ” leads immediately to a contradiction.

Suppose $n = 3$. The impossibility of the inclusion types “ $(0, 1) < (1, 1)$ ”, “ $(0, 2) < (0, 3)$ ” and “ $(0, 3) < (1, 3)$ ” implies that H_0, H_1, H_2, H_3, H_5 and H_6 are of type $(0, 0), (0, 1), (0, 2), (1, 2), (2, 3)$ and $(3, 3)$, respectively. Furthermore, the type of H_4 is either $(2, 2)$ or $(1, 3)$.

By Theorem 3.1, there exists a basis $\{a, b, c\}$ of F and two non-proper powers $w, h \in \langle a, b \rangle'$ such that $H_2 = \langle w, c^{-1}hc \rangle$. Note that, since $w \neq 1$ has trivial abelianization, $K = \langle a, b \rangle$ is the smallest free factor of F containing w . Consider the core graph X_2 corresponding to H_2 , which consists of two circles labelled w and h , and an edge e with label c from a vertex ιe in the circle with label h to a vertex τe in the circle with label w (and with τe being the base point). Since $r(H_2) = r(H_3) = 2$, the inclusion $H_2 < H_3$ is algebraic and so, the core graph X_3 corresponding to H_3 is a quotient of X_2 . In particular, the image of e , say e' , is the unique edge in X_3 with label c . If $X_3 - \{e'\}$ were disconnected then $H_2 = H_3$ (because w and h are not proper powers), which is not the case. Thus, $X_3 - \{e'\}$ is connected

and hence, it is a circle with two (possibly trivial) hairs going to $\iota e'$ and $\tau e'$. Now, let $u \in K$ be the label of a (possibly trivial) path in $X_3 - \{e'\}$ from $\tau e'$ to $\iota e'$. It is clear that $H_3 = \langle w, uc \rangle$.

Let α be the automorphism of F defined by $a \mapsto a$, $b \mapsto b$, $c \mapsto u^{-1}c$. By looking at the corresponding core graph, it is clear that $H_3\alpha = \langle w, c \rangle$, is F -strictly-compressed (again using the fact that $w \in K$ is not a proper power). Hence, H_3 is also F -strictly-compressed, which implies that H_4 (and so $H_4\alpha$) is of type $(1, 3)$.

Let $\psi \in \text{Aut}(F)$ be such that $\text{Fix } \psi = H_4 \geq H_3 = \langle w, uc \rangle$, and let $\psi' = \alpha^{-1}\psi\alpha$. A simple computation shows that $\text{Fix } \psi' = H_4\alpha \geq \langle w, c \rangle$. We noted above that K is the smallest free factor of F containing $w = w\psi'$, so K is ψ' -invariant. Then (and using also the fact that ψ' fixes c) it is easy to see that $H_4\alpha = \text{Fix } \psi' = \text{Fix } \psi'_K * \langle c \rangle$. But $H_4\alpha$ is of type $(1, 3)$, hence $\text{Fix } \psi'_K \leq K$ is of type $(0, 2)$. This is a contradiction with the fact that $\text{Fix } \psi'_K$ is a 1-auto-fixed subgroup of the free group K , of rank 2. \square

Remark 4.3 Note that the argument given in the previous observation for the case $n = 3$ is global in the sense that, assuming the existence of a chain of length $2n$, we get a contradiction by analyzing more than one consecutive inclusion. It is impossible to do this locally, that is, looking only at some particular inclusion in the chain, because, as we see below, all possible types of inclusions except “ $(0, n - 1) < (0, n)$ ”, “ $(0, n) < (1, n)$ ” and “ $(0, 1) < (1, 1)$ ” can be realized by automorphisms.

Let $\{x_1, \dots, x_n\}$ be a basis of F .

Let $1 \leq p \leq q \leq n$ be two integers and choose $r, s, t \geq 0$ and $\psi_{p,q}$ as in the proof of Theorem 4.1.

Suppose that $p + 1 \leq q$. Then, $s \geq 1$ and changing the $\psi_{p,q}$ -image of x_{r+1} from $x_1^{r+1}x_{r+1}$ to x_{r+1} we obtain another automorphism ψ of F such that

$$\text{Fix } \psi = \langle x_1, \dots, x_r, x_{r+1}, x_{r+2}^{-1}x_1x_{r+2}, \dots, x_{r+s}^{-1}x_1x_{r+s} \rangle.$$

Hence, $\text{Fix } \psi_{p,q} < \text{Fix } \psi$ is a strict inclusion of 1-auto-fixed subgroups of type “ $(p, q) < (p + 1, q)$ ”.

Suppose that $q + 1 \leq n$. Then, $t \geq 1$ and changing the $\psi_{p,q}$ -image of x_{r+s+1} from x_{r+s+1}^{-1} to $x_1^{r+s+1}x_{r+s+1}$ we obtain another automorphism ψ of F such that

$$\text{Fix } \psi = \langle x_1, \dots, x_r, x_{r+1}^{-1}x_1x_{r+1}, \dots, x_{r+s}^{-1}x_1x_{r+s}, x_{r+s+1}^{-1}x_1x_{r+s+1} \rangle.$$

Hence, $\text{Fix } \psi_{p,q} < \text{Fix } \psi$ is a strict inclusion of 1-auto-fixed subgroups of type “ $(p, q) < (p, q + 1)$ ”.

Now, let $0 = p \leq q \leq n - 1$ be two integers with $1 \leq q$. Choose $r, s, t \geq 0$ such that $r = 2$, $q + 1 = r + s$ and $n = r + s + t$. Define $\psi_{0,q}$ to be the automorphism of F given by

$$\begin{aligned} \psi_{0,q}: F &\mapsto F \\ x_1 &\mapsto [x_1, x_2]^{-1} x_1 [x_1, x_2] \\ x_2 &\mapsto [x_1, x_2]^{-1} x_2 [x_1, x_2] \\ x_j &\mapsto [x_1, x_2]^j x_j, & j = 3, \dots, r + s \\ x_k &\mapsto x_k^{-1}, & k = r + s + 1, \dots, r + s + t. \end{aligned}$$

It is not difficult to see that

$$\text{Fix } \psi_{0,q} = \langle [x_1, x_2], x_3^{-1} [x_1, x_2] x_3, \dots, x_{r+s}^{-1} [x_1, x_2] x_{r+s} \rangle,$$

which is a 1-auto-fixed subgroup of type $(0, 1 + s)$, that is, (p, q) .

Suppose $2 \leq q$. Then, $s \geq 1$ and changing the $\psi_{0,q}$ -image of x_3 from $[x_1, x_2]^3 x_3$ to x_3 we obtain another automorphism ψ of F such that

$$\text{Fix } \psi = \langle [x_1, x_2], x_3, x_4^{-1} [x_1, x_2] x_4, \dots, x_{r+s}^{-1} [x_1, x_2] x_{r+s} \rangle.$$

Hence, $\text{Fix } \psi_{0,q} < \text{Fix } \psi$ is a strict inclusion of 1-auto-fixed subgroups of type “ $(0, q) < (1, q)$ ”.

Suppose $q \leq n - 2$. Then $t \geq 1$ and changing the $\psi_{0,q}$ -image of x_{r+s+1} from x_{r+s+1}^{-1} to $[x_1, x_2]^{r+s+1} x_{r+s+1}$ we obtain another automorphism ψ of F such that

$$\text{Fix } \psi = \langle [x_1, x_2], x_3^{-1} [x_1, x_2] x_3, \dots, x_{r+s}^{-1} [x_1, x_2] x_{r+s}, x_{r+s+1}^{-1} [x_1, x_2] x_{r+s+1} \rangle.$$

Hence, $\text{Fix } \psi_{0,q} < \text{Fix } \psi$ is a strict inclusion of 1-auto-fixed subgroups of type “ $(0, q) < (0, q + 1)$ ”.

Finally, the only remaining case is $p = q = 0$. And it is obvious that there are inclusions of fixed subgroups of type “ $(0, 0) < (0, 1)$ ”. \square

Acknowledgments

Both authors thank Warren Dicks for interesting comments and suggestions and for improving some aspects of the paper. The first named author gratefully acknowledges the postdoctoral grant SB2001-0128 funded by the Spanish government, and thanks the CRM for its hospitality during the last period of this research. The second named author gratefully acknowledges partial support by DGI (Spain) through grant BFM2000-0354.

References

- [1] G.M. Bergman, Supports of derivations, free factorizations and ranks of fixed subgroups in free groups, *Trans. Amer. Math. Soc.*, **351** (1999), 1531–1550.
- [2] M. Bestvina, M. Handel, Train tracks and automorphisms of free groups, *Ann. of Math.*, **135** (1992), 1–51.
- [3] D.J. Collins, E.C. Turner, An automorphism of a free group of finite rank with maximal rank fixed point subgroup fixes a primitive element, *J. Pure Appl. Algebra*, **88** (1993), 43–49.
- [4] D.J. Collins, E.C. Turner, All automorphisms of free groups with maximal rank fixed subgroups, *Math. Proc. Cambridge Philos. Soc.*, **119** (1996), 615–630.
- [5] W. Dicks, E. Ventura, The group fixed by a family of injective endomorphism of a free group, *Contemp. Math.*, **195** (1996), 1–81.
- [6] J.L. Dyer, G.P. Scott, Periodic automorphisms of free groups, *Comm. Alg.*, **3** (1975), 195–201.
- [7] S.M. Gersten, Fixed points of automorphisms of free groups, *Adv. Math.*, **64** (1987), 51–85.
- [8] W. Imrich, E.C. Turner, Endomorphisms of free groups and their fixed points, *Math. Proc. Cambridge Philos. Soc.*, **105** (1989), 421–422.
- [9] I. Kapovich and A. Myasnikov, Stallings Foldings and Subgroups of Free Groups, *J. Algebra*, **248**, 2 (2002), 608–668.
- [10] S. Margolis, M. Sapir and P. Weil, Closed subgroups in pro-V topologies and the extension problems for inverse automata, *Internat. J. Algebra Comput.* **11**, 4 (2001), 405–445.
- [11] A. Martino, E. Ventura, On automorphism-fixed subgroups of a free group, *J. Algebra*, **230** (2000), 596–607.
- [12] A. Martino, E. Ventura, A description of auto-fixed subgroups in the free group, preprint, <http://www-eupm.upc.es/~ventura/>.
- [13] A. Martino, E. Ventura, Examples of retracts in free groups that are not the fixed subgroup of any automorphism, to appear in *J. Algebra*.
- [14] J.R. Stallings, Topology of finite graphs, *Invent. Math.*, **71** (1983), 551–565.

- [15] M. Takahasi, Note on chain conditions in free groups, *Osaka Math. Journal* **3**, 2 (1951), 221–225.
- [16] E.C. Turner, Test words for automorphisms of free groups, *Bull. London Math. Soc.*, **28** (1996), 255–263.
- [17] E. Ventura, On fixed subgroups of maximal rank, *Comm. Algebra*, **25** (1997), 3361–3375.
- [18] E. Ventura, Fixed subgroups in free groups: a survey, *Contemp. Math.*, **296** (2002), 231–255.