

ANALYTIC K-SPACES

S. TODORČEVIĆ AND C. UZCÁTEGUI

ABSTRACT. We study sequential convergence in spaces with analytic topologies avoiding thus a number of standard pathologies. For example, we identify bisequentiality of an analytic space as the Fréchet property of its square. We show that a countable Fréchet group is metrizable if and only if its topology is analytic. We also investigate the diagonal sequence properties and show their productiveness in the class of analytic spaces.

1. INTRODUCTION

This is a continuation of our paper [23] where we study effective versions of some standard topological problems and results. Thus we restrict ourselves to (regular) countable topological spaces X with the property that the family τ_X of all open subsets of X is in some sense effective. For example, if τ_X as a subset of the Cantor cube 2^X is a continuous image of the irrationals then we call any such X an *analytic space*. Many of the standard examples of countable spaces are analytic. For example, the *Arens space* [1], the *Arhangel'ski-Franklin space* [3], and the countable *Sequential fan* [14] are all analytic spaces. On the other hand, many topological applications to the study of, say, weak topologies of Banach spaces require results about countable analytic spaces (see e.g., [2]).

Recall, that X is said to be a *k-space* if and only if an arbitrary subset of X is closed just in case its intersection with an arbitrary compact subset of X is closed (see, e.g., [17]). In our context this reduces to the more familiar class of *sequential spaces*. Recall that X is said to be sequential if for every non closed $A \subseteq X$ there is a sequence of elements of A converging to a point outside of A . If we require a sequence of elements of A to converge to an arbitrary point of the closure of A we get the considerably more restrictive class of *Fréchet spaces*. It is usually in relation to these classes of spaces that one considers various ways to obtain a converging sequence out of a sequence of converging sequences. Recall that the *diagonal-sequence property* states that if $\{x_{nk}\}$ is a double-indexed sequence of members of X such that for some $x \in X$ and all n , $x_{nk} \rightarrow_k x$ then for each n we

can choose $k(n)$ such that $x_{nk(n)} \rightarrow_n x$. If we require that some infinite subsequence of $\{x_{nk(n)}\}$ converges to x rather than the sequence itself, we get the *weak diagonal sequence property*. Note that the diagonal sequence property and the weak diagonal sequence property are formally incomparable with the Fréchet property. Consider, for example, Arens space and the Sequential fan. The former has the diagonal sequence property but it is not Fréchet while the later is Fréchet but it fails even the weak diagonal sequence property. It turns out that in the context of analytic spaces the diagonal sequence property is as restrictive as first countability (metrizability) (see [20] and [23]). We give here a variation of this result by showing that an analytic sequential space with the diagonal sequence property is weakly first countable. Recall that we say that X is *weakly first countable* if for every $x \in X$ we can find a decreasing sequence of sets $B(x, n) \ni x$ such that a set V is open iff for all $x \in V$, there is m with $B(x, m) \subseteq V$. Note that the topology of an arbitrary countable weakly first countable space is analytic (in fact, $F_{\sigma\delta}$). For example, the Arens space is a typical example of a weakly first countable space. Note that every weakly first countable space has the diagonal sequence property and that every Fréchet weakly first countable space is in fact first countable.

Recall now that X is said to be *bisequential* if for every ultrafilter \mathcal{U} over X converging to some point x there is a sequence $A_n \in \mathcal{U}$ converging to x . Clearly every bisequential space is Fréchet but not vice versa. Consider, for example, the Sequential fan. Note also that every bisequential space has the weak diagonal sequence property but not vice versa. Consider, for example, the Arens space. We show however that the two properties jointly characterize bisequentiality in the class of analytic space. Thus we show that every analytic Fréchet space with the weak diagonal sequence property is bisequential. We give some application of this result to the study of products as well as to the study of countable topological groups. For example, we show that the square of an analytic Fréchet space X is Fréchet if and only if X contains no copy of the sequential fan $S(\omega)$. As another application we show that analytic Fréchet groups are metrizable solving thus the effective version of the well known problem of Malyhin (see, e.g., [14]). The preservation of the weak diagonal sequence property in products of analytic spaces seems curiously related to the problem whether the Sequential fan is a test space for the failure of this property in the class of analytic spaces. This can be seen from the fact which we show here which says that the sequential fan does not embed into the product of two analytic spaces with the weak diagonal sequence property. This can be regarded as a proof of the effective version of a conjecture of Nogura[11]. The proof of

the unrestricted version of Nogura's conjecture is given in [22] and our proof here can be regarded as its effective version.

2. TEST SPACES

In this section we work in the class of countable regular spaces and we introduce some critical examples of analytic spaces that are quite useful in testing the convergence properties introduced above. The first such a space is the Arens space [1], denoted by S_2 , the space on $\omega^{\leq 2}$ with the following topology. Each sequence of length 2 is isolated, a basic nbhd of the sequence $\langle n \rangle$ consists of all sequences of the form $\langle n, m \rangle$ for all but finitely many m 's and, finally, a set U is a basic nbhd of the empty sequence \emptyset if there is a function $f : \mathbb{N} \rightarrow \mathbb{N}$ and an integer n with the property that $\langle m \rangle, \langle m, k \rangle \in U$ for all $m \geq n$ and all $k \geq f(m)$. The following well known fact gives a clear indication of the importance of the Arens space for this area (see, e.g., [16]).

Proposition 2.1. *A sequential space is Fréchet if and only if it contains no homeomorphic copy of the Arens space S_2 .*

The sequential fan, denoted by $S(\omega)$, is the space defined over $\mathbb{N} \times \mathbb{N} \cup \{\infty\}$ where all points in $\mathbb{N} \times \mathbb{N}$ are isolated and the nbhd filter of ∞ is generated by the sets of the form $U_f = \{(n, m) \in \mathbb{N} \times \mathbb{N} : m \geq f(n)\} \cup \{\infty\}$ for $f \in \mathbb{N}^{\mathbb{N}}$. The sequential fan $S(\omega)$ is a typical Fréchet space without the weak diagonal sequence property. The following well known fact shows that inside the class of Fréchet spaces $S(\omega)$ is a test space for the weak diagonal sequence property (see, e.g., [16]).

Proposition 2.2. *A Fréchet space has the weak diagonal sequence property if and only if it contains no homeomorphic copy of $S(\omega)$.*

The following result of Nyikos[12] shows that inside the class of topological groups the two spaces perform the same task of testing

Proposition 2.3. *A topological group contains a copy of the Arens space if and only if it contains a copy of the sequential fan.*

Corollary 2.4. *A sequential topological group is Fréchet if and only if it has the weak diagonal sequence property.*

Note that we are working inside the class of countable spaces X , so if such an X contains a copy of S_2 or $S(\omega)$ then it also contains closed subspaces homeomorphic to S_2 or $S(\omega)$. This will be implicitly used below. A natural generalization of the Arens space is the Arkhangel'skiĭ-Franklin space S_ω [3]. It is defined in [3] as the direct limit of the sequence S_n ($n \in \omega$), where S_{n+1} is related to S_n the same way the Arens space S_2 is related to the converging sequence S_1 . We shall, however, not work with

this description of the space S_ω as we shall prefer to view it as an instance of a parametrized class of spaces to be considered below in Section 6. We should note, however, that the space S_ω is another test space, especially when one considers high sequential orders. For example, to show that a given sequential space X has sequential order ω_1 one typically shows this by embedding S_ω in X (see [3], [7], [23]).

3. SELECTIVE POINTS

A point $x \in X$ is called a *Fréchet point*, if for every $A \subseteq X$ with $x \in \overline{A}$ there is a sequence $x_n \in A$ converging to x (in short, X is Fréchet at x). Analogously, we define the notion of a *bisequential point*. We will say that x is a q^+ -point, if for every A with $x \in \overline{A}$ and every partition $A = \bigcup_n F_n$ of A into finite sets, there is a subset B of A such that $x \in \overline{B}$ and $|B \cap F_n| \leq 1$ for all n . We say that x is a p^+ -point if given any decreasing sequence P_n of subset of $X \setminus x$ such that $x \in \overline{P_n}$ for all n , there is a set $P \setminus x$ such that $x \in \overline{P}$ and $P \subseteq^* P_n$ (i.e. $P \setminus P_n$ is finite) for all n . We note that a point which is at the same time a Fréchet point and a p^+ -point is also called in the literature a *countably bisequential point*, or a *strongly Fréchet point* [10, 16]. We will say that x is a *selective point* if it is both a p^+ and q^+ -point. Finally, we say that a space X is Fréchet (respectively, bisequential, or selective) if every point of X is a Fréchet (respectively, bisequential, or selective) point in X .

Recall that $A^{(1)}$ is the set of all limits of convergent sequences in A . Let $A^{(0)} = A$, $A^{(\alpha+1)} = [A^{(\alpha)}]^{(1)}$ and $A^{(\beta)} = \bigcup_{\alpha < \beta} A^{(\alpha)}$ for β a limit ordinal. For $x \in \overline{A}$, the *sequential rank* of x in A , denoted by $rk(x, A)$, is the least ordinal α such that $x \in A^{(\alpha)}$. The *sequential closure* of A is the set $A^{(\omega_1)}$. Thus X is sequential iff $\overline{A} = A^{(\omega_1)}$. We show first a general fact about the sequential rank.

Lemma 3.1. *Suppose X is a countable sequential space. Then for every $A \subseteq X$ and every $x \in \overline{A}$ there is $D \subseteq A$ such that $x \in \overline{D}$ and $rk(y, E) \leq rk(x, A)$ for every $E \subseteq D$ and every $y \in \overline{E}$.*

Proof. By induction on the sequential rank. If $rk(x, A) = 0$, then take $D = \{x\}$. Suppose that $rk(x, A) > 0$ and that the conclusion is true for all countable $B \subseteq X$ and all $y \in \overline{B}$ such that $rk(y, B) < rk(x, A)$. Let $x_n \in \overline{A}$ be a sequence converging to x such that $rk(x_n, A) < rk(x, A)$. Choose a sequence V_n of clopen subsets of X such that $x_n \in V_n$ and such V_n is a relatively discrete sequence in $X \setminus \{x\}$. By the inductive hypothesis applied to x_n and $A \cap V_n$ we can find $D_n \subseteq V_n \cap A$ such that $x_n \in \overline{D_n}$ and $rk(y, E) \leq rk(x_n, V_n \cap A) (= rk(x_n, A))$ for every $E \subseteq D_n$ and every $y \in \overline{E}$. Let $D = \bigcup_n D_n$. Then D satisfies the conclusion of the Lemma. \square

We shall also need the following fact about sequential spaces.

Lemma 3.2. *Suppose X is a sequential space. Let $x \in X$ be an accumulation point of some set $A \subset X$ and let E_n be a partition of A into pairwise disjoint finite subsets. Then there is another such partition F_k of A which is coarser than E_n and which has the property that $\bigcup_{k \in M} F_k$ accumulates to x for every infinite $M \subseteq \mathbb{N}$.*

Proof. The proof is by induction on $rk(x, A)$. The case $rk(x, A) = 1$ is immediate, so let us assume that $rk(x, A) > 1$ and that the conclusion of the lemma is true for smaller ranks. Choose a sequence x_j converging to x such that $rk(x_j, A) < rk(x, A)$ for all j . By the inductive hypothesis, for each j we can choose a partition E_k^j of A into finite pairwise disjoint sets that is coarser than the given partition E_n and such that $\bigcup_{k \in M} E_k^j$ accumulates to x_j for every infinite $M \subseteq \mathbb{N}$. Choose a partition F_l of A into pairwise disjoint finite sets that is coarser than E_n and which has the property that for every j and all but finitely many l , the set F_l contains at least one of the sets of the form E_k^j for some k . It is clear that such a partition F_l satisfies the conclusion of the Lemma. \square

Proposition 3.3. *Every point of a countable sequential space is a q^+ -point.*

Proof. The proposition will be proved by induction on $rk(x, A)$. Let F_m be a partition of A into finite pieces. Let $x \in \overline{A}$, we will show that there is $B \subseteq A$ such that $|B \cap F_m| \leq 1$ for all m and $x \in \overline{B}$. The result is obvious for $x \in A^{(1)}$. Suppose we have proved the proposition for every y and B such that $rk(y, B) < rk(x, A)$. Let $x_n \in \overline{A} \setminus A$ be a sequence converging to x such that $1 \leq rk(x_n, A) < rk(x, A)$ for all n . We may assume to have also a sequence V_n pairwise disjoint open subsets of X such that $x_n \in V_n$ for all n . Applying Lemma 3.1 to the points x_n and sets $A \cap V_n$, and possibly shrinking A , we may assume that $rk(x_n, B) \leq rk(x_n, A)$ for all n and all $B \subseteq A \cap V_n$ (equivalently, for all $B \subseteq A$) such that $x_n \in \overline{B}$. By (the proof of) Lemma 3.2, going to a coarser partition than F_m , we may assume that for all n and all infinite $M \subseteq \mathbb{N}$, the union $F_M = \bigcup_{m \in M} F_m$ accumulates to x_n . Choose a sequence M_n of pairwise disjoint infinite subsets of \mathbb{N} . Then for all n , $rk(x_n, F_{M_n}) \leq rk(x_n, A) < rk(x, A)$, so the inductive hypothesis applies giving us $B_n \subseteq F_{M_n}$ such that $x_n \in \overline{B_n}$ and $|B_n \cap F_m| \leq 1$ for all $m \in M_n$. Let $B = \bigcup_{n \in \mathbb{N}} B_n$. Then $|B \cap F_m| \leq 1$ for all m and $x \in \overline{B}$, as required. \square

Proposition 3.4. *A Fréchet point $x \in X$ is a p^+ -point iff it has the weak diagonal sequence property.*

Proof. Let x be a Fréchet point. First, suppose x is a p^+ -point. Let x_{nk} be a double-indexed sequence in X such that for all n , $x_{nk} \rightarrow_k x$. Let

P_n be $\{x_{mk} : m \geq n, k \geq 1\}$. Then P_n is a decreasing sequence of sets with $x \in \overline{P_n}$. Since x is a \mathfrak{p}^+ -point, there is $P \subseteq^* P_n$ such that $x \in \overline{P}$. Since x is Fréchet, there is a sequence $\{y_m\}$ in P converging to x . Pick a subsequence y_{m_j} and an increasing sequence of integer $n(j)$ such that $y_{m_j} \in \{x_{n(j),k} : k \geq 1\}$.

Suppose now that x has the weak diagonal sequence property. To see that x is a \mathfrak{p}^+ -point, let P_n be a decreasing sequence of sets with $x \in \overline{P_n}$. Since x is Fréchet, there is $x_{nk} \in P_n$ such that $x_{nk} \rightarrow_k x$ for all n . Let $n(j)$ and $k(j)$ two sequences of integers such that $n(j)$ is increasing and $x_{n(j),k(j)} \rightarrow_j x$. Let P be the range of this sequence. Then $P \subseteq^* P_n$ for all n . \square

Corollary 3.5. *A countable Fréchet space is selective if and only if it has the weak diagonal sequence property.*

We finish this section by mentioning two results from [21] the second of which will play an important role in sections to come. First of all note the following reformulation of the result from Chapter 1, Section 12 of [21].

Theorem 3.6. *Every sequential space with a countably tight compactification is selective.*

Thus, in particular, under PFA, every space with a countably tight compactification is selective. The second result appears in [21] as Exercise 3 on page 53 and it is the result that will play a key role in the rest of the paper when analyzing bisequentiality in the class of countable analytic spaces.

Theorem 3.7. *A point in a countable analytic space is selective if and only if it is bisequential.*

Corollary 3.8. *Every countable analytic space with a countably tight compactification is bisequential.*

Recall that a typical example of an analytic topology on a countable set is the topology of pointwise convergence restricted to some countable set X of Borel functions defined on some Polish space P . By a result of Rosenthal [13], such an X will satisfy the hypothesis of Corollary 3.8 whenever its pointwise closure in $[-\infty, +\infty]^P$ contains no copy of $\beta\mathbb{N}$. It follows that the well-known result of Bourgain, Fremlin, and Talagrand [4] is an immediate consequence of Corollary 3.8.

4. DIAGONAL SEQUENCE PROPERTIES

It turns out that the diagonal sequence property is quite strong in the context of analytic Fréchet spaces. An interpretation of the analytic gap

theorem of [20] shows that a Fréchet analytic space has the diagonal sequence property iff it is first countable (see also [23, theorem 6.6]). We now show that a similar interpretation of the analytic gap theorem of [20] gives an analogous result for the class of analytic sequential rather than Fréchet spaces.

Theorem 4.1. *An analytic sequential space X is weakly first countable if and only if X has the diagonal sequence property.*

Proof. Suppose X has the diagonal sequence property. Let $C_x = \{A \subseteq X : A \rightarrow x\}$ and $D_x = \{B \subseteq X : x \notin \overline{B}\}$. Note that C_x and D_x are two orthogonal families of subsets of ω , i.e., the intersection of an arbitrary member of C_x with an arbitrary member of D_x is finite. Note also that D_x is an analytic family of subsets of ω so the analytic gap theorem of [20] applies giving us the following two alternatives written in the terminology of [20]:

- (1) There is a sequence $A(x, n)$ of members of C_x^\perp such that for all $B \in D_x$ there is m such that $B \subseteq A(x, m)$.
- (2) There is a C_x -tree all of whose branches belong to D_x .

An application of the diagonal sequence property of X easily eliminates the alternative (2). It follows that the alternative (1) holds. Without loss of generality, we may assume that the $A(x, n)$'s are increasing for each x and $x \notin A(x, n)$ for all n . Let $B(x, m) = X \setminus A(x, m)$. We claim that the $B(x, m)$'s form a weak base. Let V be an open set and $x \in V$. Suppose that $B(x, n) \not\subseteq V$ for all n . Then pick $x_n \in B(x, n) \setminus V$ and let $B = \{x_n : n \in \mathbb{N}\}$. Since $B \cap V$ is empty, then $x \notin \overline{B}$. Thus by (1) there is m such that $B \cap B(x, m)$ is finite. But this is a contradiction, since the $B(x, n)$'s are decreasing.

Now suppose that a subset V of X satisfies that for all $x \in V$, there is m with $B(x, m) \subseteq V$. We will show that V is sequentially open, and thus open. Let $x \in V$ and $A \in C_x$. By hypothesis there is m such that $B(x, m) \subseteq V$. By (1), $A \cap A(x, m)$ is finite, then $A \subseteq^* B(x, m) \subseteq V$.

Suppose now that X is weakly first countable and $B(x, m)$ is a weak base. Let $d(x, y)$ be $1/m$ if $y \in B(x, m) \setminus B(x, m+1)$ and $d(x, y) = 2$ otherwise. To see that the diagonal sequence property holds it clearly suffices to show that a sequence x_n converges to x iff $d(x, x_n) \rightarrow 0$. To show this claim, suppose $x_n \rightarrow x$ but $d(x, x_n) \not\rightarrow 0$. Let $k > 0$ be such that $A = \{x_n : d(x, x_n) > 1/k\}$ is infinite. Then $A \cup \{x\}$ is closed. Hence for all $y \notin A \cup \{x\}$ there is n_y such that $B(y, n_y) \subseteq X \setminus (A \cup \{x\})$. Since $B(x, k) \subseteq X \setminus A$. It follows that $X \setminus A$ is open. This contradicts that x_n converges to x . The other implication is straightforward. \square

Corollary 4.2. *The topology of every analytic sequential space with the diagonal sequence property is $F_{\sigma\delta}$.*

Remark 4.3. Note that the typical analytic sequential spaces S_2 and S_ω are weakly first countable and that in some sense this is the way their topologies are given.

Let us now return to analytic Fréchet spaces. We have already noted that in this context the diagonal sequence property reduces to first countability. The following result gives a characterization of the weak diagonal sequence property in this context.

Theorem 4.4. *An analytic Fréchet space is bisquential if and only if it has the weak diagonal sequence property.*

Proof. Only the implication from the weak diagonal sequence property towards the bisquentiality is not obvious. Suppose X is Fréchet and has the weak diagonal sequence property. From propositions 3.3 and 3.4 we have that every point of X is selective. Then by theorem 3.7 X is bisquential. \square

Corollary 4.5. *A Fréchet analytic space is bisquential iff it contains no closed copy of $S(\omega)$.*

Remark 4.6. The assumption that X is Fréchet is essential here. For example, the result does not extend to the wider class of sequential spaces which can be seen by noting that Arens space S_2 contains no copy of $S(\omega)$.

Corollary 4.7. *A countable space with an F_σ basis is bisquential if and only if it is sequential.*

Proof. Suppose X is sequential. Since neither Arens space nor the sequential fan admits a F_σ basis, then X contains no closed copy of either of them. Thus X is Fréchet and has the weak diagonal sequence property. Hence X is bisquential. \square

Remark 4.8. There are spaces with a F_σ basis which are not sequential and spaces with F_σ basis which are sequential but not metrizable. For example, it is well known that the space $CO(2^{\mathbb{N}})$ of all clopen subsets of $2^{\mathbb{N}}$, as a subspace of $\{0, 1\}^{2^{\mathbb{N}}}$ with the product topology, it is not sequential but as it is easily checked it has an F_σ basis. On the other hand, the subspace $BCO(2^{\mathbb{N}})$ of $CO(2^{\mathbb{N}})$ consisting only of basic clopen sets (including the empty set, of course) is (bi)sequential (see Example 5.6 of [23]).

5. PRODUCTS

The sequential fan $S(\omega)$ is a typical Fréchet space whose square is not Fréchet (consider the following subset of $S(\omega)^2$: $Z = \{(m, n), (0, m)\}$:

$m, n \in \mathbb{N}$ and note that $(\infty, \infty) \in \overline{Z}$ while no sequence $((m_k, n_k), (0, m_k))$ in Z converges to (∞, ∞) .) We start this section with a result which shows that in the context of countable analytic space, $S(\omega)$ is a test space for this phenomenon.

Theorem 5.1. *An analytic space X is bisquential if and only if its square X^2 is Fréchet .*

Proof. Suppose X^2 is Fréchet . Since $S(\omega)^2$ is not Fréchet , then X cannot contain a closed copy of $S(\omega)$. Hence by theorem 4.5 X is bisquential. \square

Corollary 5.2. *The square of an analytic Fréchet space X is Fréchet if and only if X contains no closed copy of the sequential fan $S(\omega)$.*

Remark 5.3. Another example of two Fréchet analytic spaces whose product is not sequential is the following. Let \mathcal{F} be the dual filter of $\text{FIN} \times \emptyset$, that is to say, the filter on $\mathbb{N} \times \mathbb{N}$ given by $A \in \mathcal{F}$ iff there is n such that $A \cap (\{m \in \mathbb{N} : m < n\} \times \mathbb{N}) = \emptyset$. Let Y be the space $\mathbb{N} \times \mathbb{N} \cup \{\infty\}$ with the topology where every element of $\mathbb{N} \times \mathbb{N}$ is isolated and \mathcal{F} is the nbhd filter of ∞ . Then Y is Fréchet (in fact, metrizable) and $S(\omega) \times Y$ is not sequential. To see this, consider the diagonal $D \subseteq (\mathbb{N} \times \mathbb{N})^2$ which is sequentially discrete but $(\infty, \infty) \in \overline{D}$. It follows that if X is an analytic Fréchet space such that $X \times \mathbb{Q}$ is sequential, then X is bisquential. To see this, suppose towards a contradiction that $S(\omega)$ embeds in X . Let Y be the metrizable space given above. Then Y is homeomorphic to a closed subspace of \mathbb{Q} . Thus $X \times \mathbb{Q}$ contains a closed copy of $S(\omega) \times Y$. But this is a contradiction, since $S(\omega) \times Y$ is not sequential.

Let us now turn to the productiveness of the weak diagonal sequence property. In [11] Nogura shows that there exist two countable Fréchet spaces X and Y with the weak diagonal sequence property such that $X \times Y$ is neither Fréchet nor it has the weak diagonal sequence property and asks whether for such X and Y the product $X \times Y$ would have one of these two properties if it has the other. In [22] the first author showed that the Open Coloring Axiom implies the preservation of the weak diagonal sequence property whenever one takes products that are Fréchet. However, this still leaves the following question unanswered.

Question 5.4. *Is the weak diagonal sequence property productive in the class of analytic spaces?*

One can similarly ask whether the assumption of OCA can be removed from the proof of [22] provided we are working in the class of countable spaces with analytic topologies. We shall now see that this is indeed so.

Theorem 5.5. *Suppose X and Y are two analytic spaces with the weak diagonal sequence property. Then their product $X \times Y$ will have the weak diagonal sequence property whenever it is Fréchet.*

Proof. The proof is really just an effective version of the proof of Nogura's conjecture given in [22], so we are assuming the reader has a copy of that proof at hand. By way to a contradiction, we assume that we have a Fréchet product $X \times Y$ which fails to have the weak diagonal sequence property. By Proposition 2.2 the sequential fan $S(\omega)$ embeds into $X \times Y$. From this one easily constructs two analytic topologies τ_X and τ_Y on $\omega^2 \cup \{\infty\}$ with ∞ as the only non isolated point such that τ_X and τ_Y both have the weak diagonal sequence property while the topology of $S(\omega)$ is generated by $\tau_X \cup \tau_Y$ as subbasis. For $n \in \omega$, set

$$C_n = \{n\} \times \omega$$

Note that each C_n is a converging sequence in both topologies τ_X and τ_Y . Let

$$\mathcal{A} = \{A \subseteq \omega^2 : A \rightarrow_{\tau_X} \infty \text{ and } A \cap C_n \text{ is finite for all } n\},$$

$$\mathcal{B} = \{B \subseteq \omega^2 : B \rightarrow_{\tau_Y} \infty \text{ and } B \cap C_n \text{ is finite for all } n\}.$$

Thus, \mathcal{A} (respectively, \mathcal{B}) is the family of all sequences that converge to ∞ relative to τ_X (respectively, relatively to τ_Y) and which are orthogonal to each member of the sequence $\{C_n\}$ of converging sequences. Let

$$\mathcal{X} = \{(A, B) \in \mathcal{A} \times \mathcal{B} : A \cap B = \emptyset\}.$$

We endow \mathcal{X} with the standard separable metric topology induced from 2^{ω^2} . Consider the following subset of the set $\mathcal{X}^{[2]}$ of all unordered pairs of elements of \mathcal{X} :

$$\mathcal{K} = \{(A, B), (A', B')\} \in \mathcal{X}^{[2]} : (A \cap B') \cup (A' \cap B) \neq \emptyset\}.$$

Note that \mathcal{K} is an open subset of $\mathcal{X}^{[2]}$. Recall the following alternative of the Open Coloring Axiom:

Case 2: There is a decomposition $\mathcal{X} = \bigcup_{n=0}^{\infty} \mathcal{X}_n$ such that $(\mathcal{X}_n)^{[2]} \cap \mathcal{K} = \emptyset$ for all n .

The proof of [22] shows that this alternative of OCA is not possible. The proof given in Chapter 8 of [18] that PFA implies OCA produces a forcing extension of the universe which contains an uncountable $\mathcal{Y} \subseteq \mathcal{X}$ such that $\mathcal{Y}^{[2]} \subseteq \mathcal{K}$ and for some $f \in \omega^\omega$,

$$A \cup B \subseteq \Gamma_f \text{ for all } (A, B) \in \mathcal{Y}.$$

Note that while \mathcal{Y} consists of ordered pairs of the ground model sets the set Γ_f might be new so the ground model sets from $\tau_X \cup \tau_Y$ may not be

sufficient to separate it from the point ∞ of $S(\omega)$. However if in the forcing extension we let $\dot{\tau}_X$ and $\dot{\tau}_Y$ be the images of the new set of irrationals under the same two continuous maps that define these two topologies in the ground model, the Shoenfield absoluteness theorem gives us that $\dot{\tau}_X \cup \dot{\tau}_Y$ is still a subbasis of the sequential fan $S(\omega)$. It follows that the possibly new set Γ_f does not accumulate to ∞ in $S(\omega)$ relative to the new subbasis, and therefore, working in the forcing extension, we can find $U \in \dot{\tau}_X$ and $V \in \dot{\tau}_Y$ such that $\infty \in U \cap V$ and

$$U \cap V \cap \Gamma_f = \emptyset.$$

Working still in the forcing extension, for $D \subseteq \omega \times \omega$ and $n \in \omega$, we set

$$D/n = D \setminus (n \times \omega) \text{ and } D \upharpoonright n = D \cap (n \times \omega).$$

Note that for every $(A, B) \in \mathcal{Y}$ there is n such that $A/n \subseteq U$ and $B/n \subseteq V$. So we can find $n \in \omega$ and uncountable subset \mathcal{Y}_0 of \mathcal{Y} such that:

- (a) $A/n \subseteq U$ and $B/n \subseteq V$ for all $(A, B) \in \mathcal{Y}_0$,
- (b) $A \upharpoonright n = A' \upharpoonright n$ and $B \upharpoonright n = B' \upharpoonright n$ for all $(A, B), (A', B') \in \mathcal{Y}_0$.

Pick two distinct elements (A, B) and (A', B') of \mathcal{Y}_0 . Then the unordered pair $\{(A, B), (A', B')\}$ belongs to \mathcal{K} , and therefore,

$$U \cap V \cap \Gamma_f \supseteq (A \cap B') \cup (A' \cap B) \neq \emptyset,$$

a contradiction. This finishes the proof. \square

Remark 5.6. Note that the argument just given is a copy of the corresponding part of the proof in [22] which also shows that the following alternative of the effective OCA fails: *Case 2: There is a perfect set $\mathcal{Y} \subseteq \mathcal{X}$ such that $\mathcal{Y}^{[2]} \subseteq \mathcal{K}$.* This would give us another way to prove Theorem 5.5 if the effective OCA can indeed be applied. Unfortunately, assuming no additional set-theoretic axioms, the effective OCA applies only to analytic base sets \mathcal{X} which is not so in our case here. It turns out that there exist analytic subfamilies \mathcal{A}^* and \mathcal{B}^* of \mathcal{A} and \mathcal{B} , respectively, which are large enough in the sense that they contain all the results of the applications of the weak diagonal sequence property needed for in the argument of [22] in order to eliminate the Case 2 for the corresponding analytic family $\mathcal{X}^* = \{(A, B) \in \mathcal{A}^* \times \mathcal{B}^* : A \cap B = \emptyset\}$. For a given subset C of ω , set $C^{[2]} = \{(m, n) \in C^2 : m < n\}$. For $\mathcal{V} \in \{\mathcal{A}, \mathcal{B}\}$, set

$$\mathcal{R}_{\mathcal{V}} = \{(C, A) \in \mathcal{P}(\omega) \times \mathcal{V} : A \subseteq^* C^{[2]}\}.$$

The assumption that τ_X and τ_Y have the weak diagonal sequence property yields that $\mathcal{R}_{\mathcal{A}}$ and $\mathcal{R}_{\mathcal{B}}$ are two coanalytic relations whose projections on the first coordinate include the family $[\omega]^\omega$ of all infinite subsets of ω . It turns out that these two relations admit local uniformizations by continuous

functions. In other words, there is $D^* \in [\omega]^\omega$ and continuous functions $f : [D^*]^\omega \rightarrow \mathcal{P}(\omega^2)$ and $g : [D^*]^\omega \rightarrow \mathcal{P}(\omega^2)$ such that $\mathcal{R}_{\mathcal{A}}(C, f(C))$ and $\mathcal{R}_{\mathcal{B}}(C, g(C))$ hold for all $C \in [D^*]^\omega$. Let \mathcal{A}^* and \mathcal{B}^* be the families obtained from the ranges of f and g , respectively, by closing under finite changes. Note that so obtained families \mathcal{A}^* and \mathcal{B}^* are analytic. The reader is also left to check that \mathcal{A}^* and \mathcal{B}^* are indeed large enough to allow the proof from [22] be applied in showing that \mathcal{X}^* cannot be covered by countably many sets whose symmetric squares avoid \mathcal{K} .

6. HOMOGENEOUS SPACES

We start by recalling a well-known construction. Let $\vec{\mathcal{F}} = \{\mathcal{F}_s : s \in \omega^{<\omega}\}$ be a collection of filters over \mathbb{N} each of which contains every cofinite sets. Define a topology $\tau_{\vec{\mathcal{F}}}$ over $\omega^{<\omega}$ by letting a subset U of $\omega^{<\omega}$ be open if and only if

$$\{n \in \mathbb{N} : \widehat{s}n \in U\} \in \mathcal{F}_s \text{ for all } s \in U$$

It is clear that $\tau_{\vec{\mathcal{F}}}$ is T_2 , zero dimensional and has no isolated points. If all filters \mathcal{F}_s are equal to a filter \mathcal{F} we will denote by $\tau_{\mathcal{F}}$ the corresponding topology. A particular important case is when \mathcal{F} is the filter CF of all cofinite sets. It is not difficult to show that the corresponding space is homeomorphic to S_ω . To analyze this class of spaces we need the following notion. An $\vec{\mathcal{F}}$ -tree with root $s \in \omega^{<\omega}$ is a subset T of $\omega^{<\omega}$ with s as its minimal node such that

- (i) If $t \in T$, then $t|m \in T$ for all $lh(s) \leq m \leq lh(t)$.
- (ii) $\{n : \widehat{t}n \in T\} \in \mathcal{F}_t$, for all $t \in T$.

Proposition 6.1. *The collection of all $\vec{\mathcal{F}}$ -trees with root s forms a local basis at s for the topology $\tau_{\vec{\mathcal{F}}}$.*

Proof. It is clear that every $\vec{\mathcal{F}}$ -tree is a $\tau_{\vec{\mathcal{F}}}$ -open set. So, it suffices to show that for every $\tau_{\vec{\mathcal{F}}}$ -open set O and any $s \in O$ there is a $\vec{\mathcal{F}}$ -tree T with stem s contained in O . We will define by induction a sequence $T_n \subseteq \omega^{lh(s)+n}$ such that $T_n \subseteq O$ for all n . For $t \in \omega^{<\omega}$, let $A_t = \{n : \widehat{t}n \in O\}$. Since O is open, $A_t \in \mathcal{F}_t$ for every $t \in O$. Let $T_0 = \{s\}$. Suppose we have defined T_k and let

$$T_{k+1} = \bigcup_{t \in T_k} \{\widehat{t}n : n \in A_t\}$$

By the inductive hypothesis, $T_{k+1} \subseteq O$ and therefore $A_t \in \mathcal{F}_t$ for every $t \in T_{k+1}$. Let $T = \bigcup_k T_k$. Then T is a $\vec{\mathcal{F}}$ -tree with stem s and by construction $T \subseteq O$. \square

A particular striking class of examples is obtained when one takes $\mathcal{F}_s = \mathcal{U}$ for all s , where \mathcal{U} is a fixed non-principal ultrafilter on ω . The corresponding space $(\omega^{<\omega}, \tau_{\mathcal{U}})$ resembles the space of Sirota [15] though in this exact form

it appears in the paper of Louveau [9] who showed that in this case $\tau_{\mathcal{U}}$ is an extremely disconnected topology on $\omega^{<\omega}$. Note that so extremal choice of the parameter is not going to give us anything new in the realm of analytic spaces since every analytic extremely disconnected space must be discrete. To see this, suppose X is an extremely disconnected non discrete analytic space. Let $x \in X$ be a non isolated point. Let $\{O_i\}_i \in I$ be a maximal family of pairwise disjoint open sets such that $x \notin \overline{O_i}$ for all $i \in I$. Define an ideal \mathcal{I} over I by letting $A \in \mathcal{I}$ if and only if $x \notin \overline{\bigcup_{i \in A} O_i}$. Then \mathcal{I} is a non principal analytic ideal on I . It is also easy to check that it is maximal. However, it is well-known that nonprincipal maximal ideals are never analytic.

It is easy to see that $\tau_{\mathcal{F}}$ is never a Fréchet topology. For instance, observe that \emptyset is an accumulation point of ω^2 , but no sequence in ω^2 converges to it. Recall that a filter \mathcal{F} over \mathbb{N} is said to be *Fréchet* if for all $A \subseteq \mathbb{N}$ that has nonempty intersection with every member of \mathcal{F} , there is an infinite $B \subseteq A$ such that $B \setminus C$ is finite for all $C \in \mathcal{F}$ (or equivalently, the space $\mathbb{N} \cup \{\infty\}$ with the topology where each $n \in \mathbb{N}$ is isolated and \mathcal{F} is the nbhd filter of ∞ is a Fréchet space). The next proposition characterizes when $\tau_{\mathcal{F}}$ is sequential.

Proposition 6.2. *The topology $\tau_{\mathcal{F}}$ is sequential if and only if \mathcal{F}_s is Fréchet for every $s \in \omega^{<\omega}$.*

Proof. Suppose that $\tau_{\mathcal{F}}$ is sequential. Fix $s \in \omega^{<\omega}$. It is clear that $X_s = \{s\} \cup \{\widehat{s}n : n \in \mathbb{N}\}$ is a closed subspace of $\omega^{<\omega}$ and thus it is itself sequential. Since s is the only non isolated point of X_s , then it is Fréchet. Thus \mathcal{F}_s is Fréchet as it is the nbhd filter of s in X_s . Conversely, suppose that each \mathcal{F}_s is Fréchet and let $A \subseteq \omega^{<\omega}$. We will show that the sequential closure $[A]_{\text{seq}}$ of A is $\tau_{\mathcal{F}}$ -closed. Let $t \notin [A]_{\text{seq}}$. We claim that the set $D = \{n \in \mathbb{N} : \widehat{t}n \notin [A]_{\text{seq}}\}$ belongs to \mathcal{F}_t . Otherwise, if $D \notin \mathcal{F}_t$, then $E = \mathbb{N} \setminus D$ satisfies that $E \cap C \neq \emptyset$ for every $C \in \mathcal{F}_t$. Therefore, as \mathcal{F}_t is Fréchet, there is $B \subseteq E$ such that $B \setminus C$ is finite for all $C \in \mathcal{F}_t$. This says that $\{\widehat{t}n : n \in B\}$ is a sequence in $[A]_{\text{seq}}$ converging to t , which contradicts our assumption. Thus D belongs to \mathcal{F}_t and therefore $[A]_{\text{seq}}$ is closed. \square

Corollary 6.3. *If $(\omega^{<\omega}, \tau_{\mathcal{F}})$ is sequential, then S_{ω} embeds into it as a closed subspace.*

Question 6.4. *Is there a homogeneous analytic sequential space of sequential order ω_1 containing no copy of S_{ω} ?*

Remark 6.5. In [24] it is shown that a closed subspace X of S_{ω} contains a copy of S_{ω} iff the sequential rank of X is ω_1 . Moreover, it is also shown that S_{ω} contains subspaces with Borel topology of arbitrarily high Borel rank.

Therefore if there is a space as in 6.4, then there must be a uniform bound for the Borel complexity of its subspaces (with Borel topology).

The following simple fact shows that many of the spaces from the class are indeed homogeneous.

Proposition 6.6. *For every filter \mathcal{F} , the space $(\omega^{<\omega}, \tau_{\mathcal{F}})$ is homogeneous.*

Proof. Since $(\omega^{<\omega}, \tau_{\mathcal{F}})$ is a regular space without isolated points, it suffices to show (see [5]) that for every $s, t \in \omega^{<\omega}$ there are two $\tau_{\mathcal{F}}$ -clopen nbhds U and V , respectively, of s and t , and a homeomorphism $h : U \rightarrow V$ with $h(s) = t$. Let U and V be the \mathcal{F} -trees with stem s , respectively t , $\{u \in \omega^{<\omega} : s \leq u\}$ and $\{u \in \omega^{<\omega} : t \leq u\}$. Define h by $h(\widehat{s}u) = \widehat{t}u$. It is easy to check that h is an homeomorphism. \square

For two filters \mathcal{F} and \mathcal{G} on ω we write $\mathcal{F} \leq \mathcal{G}$ if there is A in \mathcal{F} and B in \mathcal{G}^+ such that the restriction of \mathcal{F} on A and the restriction of \mathcal{G} on B are isomorphic filters. Note that if $(\omega^{<\omega}, \tau_{\mathcal{F}})$ is homeomorphic to a subspace of $(\omega^{<\omega}, \tau_{\mathcal{G}})$ then $\mathcal{F} \leq \mathcal{G}$.

Proposition 6.7. *There is a family \mathcal{F}_i ($i \in I$) of size bigger than the continuum of Fréchet filters on ω such that $\mathcal{F}_i \not\leq \mathcal{F}_j$ whenever $i \neq j$.*

Proof. We construct the filters on $2^{<\omega}$ rather than on ω identifying the Cantor set 2^ω with the set of branches of the binary tree $2^{<\omega}$. For a subset C of the complete binary tree $2^{<\omega}$, let $[C] = \{x \in 2^\omega : x \cap C \text{ is infinite}\}$. For a subset X of the Cantor set 2^ω , let \mathcal{F}_X be the filter on $2^{<\omega}$ generated by the complements of branches from X as well as complements of finite sets. Note that \mathcal{F}_X is always a Fréchet filter on $2^{<\omega}$. Suppose $\phi : A \rightarrow B$ is a bijection witnessing $\mathcal{F}_X \leq \mathcal{F}_Y$ for some subsets X and Y of the Cantor set. Define $f_\phi : [A] \rightarrow \mathcal{P}([B])$ and $g_\phi : [B] \rightarrow \mathcal{P}([A])$ as follows

$$\begin{aligned} f_\phi(x) &= \{y \in 2^\omega : \phi''x \cap y \text{ is infinite}\}, \\ g_\phi(y) &= \{x \in 2^\omega : \phi^{-1}y \cap x \text{ is infinite}\}. \end{aligned}$$

Note that $f_\phi(x)$ is a finite subset of Y for all $x \in X$ and that $g_\phi(y)$ is a finite subset of X for all $y \in Y$. Also note that $y \in f_\phi(x)$ iff $x \in g_\phi(y)$. Pick a well-ordering of 2^ω of the minimal possible length. Using a standard diagonalisation argument over $<_w$, we can construct a subset Z of 2^ω of size continuum such that for every bijection $\phi : A \rightarrow B$ between two subsets of $2^{<\omega}$ there is $z_0 \in Z$ such that, if for some $x >_w z_0$ in Z a set of the form $f_{\phi(x)}$ of $g_{\phi(x)}$ is finite, then it contains no elements of Z that are $>_w x$. Pick a family \mathcal{X} of size bigger than the continuum of subsets of Z such that $X \setminus Y$ has size continuum for every pair X and Y of distinct elements of \mathcal{X} . Then \mathcal{F}_X ($X \in \mathcal{X}$) satisfies the conclusion of the Proposition. \square

Corollary 6.8. *There is a family X_i ($i \in I$) of size bigger than the continuum of sequential homogeneous spaces of sequential order ω_1 such that the space X_i is not homeomorphic to a subspace of X_j whenever $i \neq j$.*

Note that Corollary 6.8 gives a very generous positive answer to a question from [3] (see page 319). The question has been actually answered long ago by Kannan[8] though he was able to construct only two new sequential homogeneous spaces. It is interesting, however, that Kannan's spaces are both analytic so one may ask for an analytic analogue of Corollary 6.8.

Question 6.9. *Is there is an uncountable family of pairwise nonhomeomorphic analytic sequential spaces of sequential order ω_1 .*

Applying the result of van Douwen [5] to homogeneous space $(\omega^{<\omega}, \tau_{\mathcal{F}})$, we conclude that for every \mathcal{F} and every countable and infinite group G there is a topology τ on G such that (G, τ) is homeomorphic to $(\omega^{<\omega}, \tau_{\mathcal{F}})$ and such that the multiplication of G is left-continuous with respect to the topology τ . So, it is natural to ask whether one can find such a topology τ for which the group operations are actually continuous. It turns out that there is no group structure on $\omega^{<\omega}$ compatible with $\tau_{\mathcal{F}}$ for \mathcal{F} Fréchet . In fact, one can say even more.

Theorem 6.10. *If a topological group is homeomorphic to a space of the form $(\omega^{<\omega}, \tau_{\mathcal{F}})$, then it has no non trivial convergent sequences.*

Proof. Let $*$ and $^{-1}$ be some group operations on $\omega^{<\omega}$ continuous relative to some topology on $\omega^{<\omega}$ of the form $\tau_{\mathcal{F}}$ for a filter \mathcal{F} on \mathbb{N} . We can assume w.l.o.g that \emptyset is the identity element. Suppose $x_k \rightarrow x$ is a non trivial convergent sequence in $(\omega^{<\omega}, \tau_{\mathcal{F}})$. Also w.l.o.g. assume that $x = \emptyset$. Observe that any sequence $\tau_{\mathcal{F}}$ -convergent to \emptyset is eventually of the form $\langle n_k \rangle$ for some strictly increasing sequence $n_k \in \mathbb{N}$. Thus we can assume that the x_k 's are of that form. Now for each fixed k , we have that $\langle n_j \rangle * \langle n_k \rangle \rightarrow \langle n_k \rangle$. The only non trivial sequences converging to $\langle n \rangle$ relative to $\tau_{\mathcal{F}}$ are eventually of the form $\langle n, m_k \rangle$ for some strictly increasing sequence $m_k \in \mathbb{N}$. Therefore we can find a strictly increasing sequence j_k of integers such that $y_k = \langle n_{j_k} \rangle * \langle n_k \rangle$ belongs to ω^2 for all k . By the joint continuity of the operations y_k converges to \emptyset . This is a contradiction, since no sequence in ω^2 converges to \emptyset . \square

Remark 6.11. It has been shown by Louveau [9] (see also Sirota[15]) that if \mathcal{U} is a selective ultrafilter, then $([\omega]^{<\omega}, \tau_{\mathcal{U}})$ is a topological group with the operation of symmetric difference, so the hypothesis of Theorem 6.10 is apriori realizable.

7. ANALYTIC GROUPS

It is known that there exist analytic homogeneous sequential spaces with an arbitrary sequential order $\leq \omega_1$ (see [6]). It turns out that for topological groups the situation is much less clear. While there exists a countable (analytic) sequential group of sequential order ω_1 (the free group of the converging sequence), the existence of such group of sequential order $< \omega_1$ depends presently on the Continuum Hypothesis (see, [14]). It is therefore rather natural to investigate the following Question.

Question 7.1. *What are the possible sequential orders of analytic sequential groups?*

This can also be considered as a variation on the following well known open problem about countable topological groups (see, e.g., [14]).

Question 7.2. *(Malyhin) Is every countable Fréchet topological group metrizable?*

The countability requirement is quite essential in Malyhin's problem. Consider, for example, the subgroup of the Haar group $\{0, 1\}^{\omega_1}$ consisting of countably supported members of $\{0, 1\}^{\omega_1}$. On the other hand, it is well known that the full Haar group $\{0, 1\}^{\omega_1}$ may be a Fréchet space under various additional set-theoretic assumptions, so in such a situation any countable dense subgroup of $\{0, 1\}^{\omega_1}$ gives a negative answer to Malyhin's problem (see [14]). Perhaps less known is the fact that only Martin's axiom is sufficient to produce not only negative answer to Malyhin's problem but also to the problem of productiveness of the Fréchet property in the realm of countable topological groups. This can easily be deduced from the fact (first pointed out in the Remark on p.150 of [19]) that under MA there exist two sets of reals X and Y such that $C_p(X)$ and $C_p(Y)$ are Fréchet spaces but their product is not. To see this, pick a countable subset D of the product which accumulates to 0 but no sequence of elements of D converges to 0, and let G and H be the subgroups of $C_p(X)$ and $C_p(Y)$, respectively, generated by the projections of D . It is easily seen that the topologies of such subgroups G and H can never be analytic. Similarly, no countable dense subgroup of $\{0, 1\}^{\omega_1}$ can have analytic topology. So, it is natural to consider the status of Malyhin's problem in the class of analytic spaces. As an application of theorem 4.4, we give a positive answer to the effective version of Malyhin's problem.

Theorem 7.3. *A countable Fréchet topological group is metrizable iff its topology is analytic.*

Proof. Only the reverse implication needs a proof. Let G be a Fréchet topological group with an analytic topology. It suffices to show that G is

first countable. By theorems 2.4 and 4.4, G is bisquential. It is known that bisquential groups are first countable (see [14]). To see this, let \mathcal{U} be an ultrafilter extending the nbhd filter of the identity element e of G and moreover assume that \mathcal{U} contains no nowhere dense sets. Let A_n be a sequence of elements of \mathcal{U} converging to e . Let $B_n = \text{int}(\overline{A_n})$. Note that B_n is nonempty for all n . Moreover, note that $B_n \cdot (B_n)^{-1}$ form a countable nbhd base of e . \square

Remark 7.4. There are examples of sequential analytic topological groups without the Fréchet property. For example, the free topological group over the convergent sequence is sequential but not Fréchet and its topology is analytic (see [14]).

REFERENCES

- [1] R. Arens, Note on convergence in topology, *Math. Mag.*, 23:229–234, 1950.
- [2] S. Argyros, G. Godefroy and H. P. Rosenthal, Descriptive Set Theory and Banach Spaces, in *Handbook of Banach Spaces*, vol 2, , Elsevier 2002.
- [3] A. V. Arkhangel'skiĭ and S. P. Franklin, Ordinal invariants for topological spaces, *Mich. Math. J.*, 15:313–320, 1968.
- [4] J. Bourgain, D.H. Fremlin and M. Talagrand, Pointwise compact sets of Baire-measurable functions. *Amer. J. Math.* 100:845–886, 1978 .
- [5] E. K. van Douwen, Countable homogenous spaces and countable groups, in Z. Frolik et al., editor, *Proc. Sixth Prague Topological Symposium 1986*, pages 135–154, Helderman Verlag, 1988.
- [6] L. Foged, Sequential order of homogeneous spaces and product spaces, *Topology Proc.*, 6:287-298, 1981.
- [7] D. H. Fremlin, Sequential convergence in $C_p(X)$, *Comment. Math. Univ. Carolin.*, 35:371-382, 1994.
- [8] V. Kannan, An extension that nowhere has the Fréchet property, *Michigan Math. J.*, 20:225-234, 1973.
- [9] A. Louveau, Sur un article de S.Sirota, *Bull. Sci. Math.*, 96: 3-7 1972.
- [10] E. Michael, A quintuple quotient quest, *Top. and its Appl.*, 2:91–138, 1972.
- [11] T. Nogura, *Products of $\langle \alpha_i \rangle$ -spaces*, *Topology and its Appl.*, 21, 251-259, 1985.
- [12] P. Nyikos, Metrizable and the Frechet-Urysohn property in topological groups, *Proc. Amer. Math. Soc.*, 83(4):793–801, 1981.
- [13] H.P. Rosenthal, Pointwise compact subsets of the first Baire class. *Amer. J. Math.* 99:362–378, 1977.
- [14] D. Shakhmatov, Convergence in the presence of algebraic structure, in M. Husek and J. van Mill (editors), *Recent Progress in Topology II*, Elsevier 2002.
- [15] S. Sirota, A product of topological groups and extremal disconnectedness, *Mat. Sb.(N.S.)*, 97(121):179-192, 1969.
- [16] F. Siwiec, Generalizations of first axiom of countability, *Rocky Mountain J. Math.*, 5:1–60, 1975.
- [17] N. Steenrod, A convenient category of spaces, *Mich. Math. J.*, 15:133-152, 1967.
- [18] S. Todorčević, *Partition Problems in Topology*, , American Mathematical Society, Providence 1989.

- [19] S. Todorčević, Some applications of S and L combinatorics, *Ann. New York Acad. Sci.*, 705, 130-167, 1993.
- [20] S. Todorčević, Analytic gaps, *Fund. Math.*, 150(1):55–66, 1996.
- [21] S. Todorčević, *Topics in Topology*, LNM 1652. Springer, 1997.
- [22] S. Todorčević, A proof of Nogura's conjecture, *Proc. Amer. Math. Soc.*, 131: 3919-3923, 2003.
- [23] S. Todorčević and C. Uzcátegui, Analytic topologies over countable sets, *Top. and its Appl.*, 111(3):299–326, 2001.
- [24] C. Uzcátegui, On the complexity of the subspaces of S_ω , *Fund. Math.*, 176: 1–16, 2003.

Université Paris VII-C.N.R.S., UMR 7056, 2 Place Jussieu, 75251 Paris Cedex 05, France.

E-mail: `stevo@math.jussieu.fr`

Departamento de Matemáticas, Facultad de Ciencias, Universidad de Los Andes, Mérida 5101, Venezuela.

E-mail: `uzca@ciens.ula.ve`