

## LARGE CARDINALS AND $L$ -LIKE UNIVERSES

SY D. FRIEDMAN

There are many different ways to extend the axioms of ZFC. One way is to adjoin the axiom  $V = L$ , asserting that every set is constructible. This axiom has many attractive consequences, such as the generalised continuum hypothesis (GCH), the existence of a definable wellordering of the class of all sets, as well as strong combinatorial principles, such as  $\diamond$ ,  $\square$  and the existence of morasses.

However  $V = L$  adds no consistency strength to ZFC. As many interesting set-theoretic statements have consistency strength beyond ZFC, it is common in set theory to assume at least the existence of inner models of  $V$  which contain large cardinals.

Can we simultaneously have the advantages of both the axiom of constructibility and the existence of large cardinals? Unfortunately even rather modest large cardinal hypotheses, such as the existence of a measurable cardinal, refute  $V = L$ . We can however hope for the following compromise:

*$V$  is an “ $L$ -like” model containing large cardinals.*

In this article we explore the possibilities for this assertion, for various notions of “ $L$ -like” and for various types of large cardinals.

There are two approaches to this problem. The first approach is via the *Inner model program*. Show that any universe with large cardinals has an  $L$ -like inner model with large cardinals.

The inner model program, through use of fine structure theory and the theory of iterated ultrapowers, has succeeded in producing very  $L$ -like inner models containing many Woodin cardinals.

An alternative approach is given by the

---

The author wishes to thank the Austrian Science Fund (FWF) for its generous support and the Centre de Recerca Matemàtica (CRM), where this article was written in the month of September 2005.

*Outer model program.* Show that any universe with large cardinals has an  $L$ -like outer model with large cardinals.

We will show that  $L$ -like outer models with extremely large cardinals can be obtained using the method of iterated forcing.

### *Large cardinals*

A cardinal  $\kappa$  is *inaccessible* iff it is uncountable, regular and larger than the power set of any smaller cardinal. It is *measurable* iff there is a  $\kappa$ -complete, nonprincipal ultrafilter on  $\kappa$ .

Measurability is equivalent to a property expressed in terms of *embeddings*, and stronger large cardinal properties are also expressed in this way. As usual,  $V$  denotes the universe of all sets. Let  $M$  be an inner model, i.e., a transitive proper class that satisfies the axioms of ZFC. A class function  $j : V \rightarrow M$  is an *embedding* iff it preserves the truth of formulas with parameters in the language of set theory and is not the identity. If  $j$  is an embedding then there is a least ordinal  $\kappa$  such that  $j(\kappa) \neq \kappa$ , called the *critical point* of  $j$ , which is a measurable cardinal.

For an ordinal  $\alpha$ ,  $j : V \rightarrow M$  is  $\alpha$ -*strong* iff  $V_\alpha$  is contained in  $M$ . A cardinal  $\kappa$  is  $\alpha$ -strong iff there is an  $\alpha$ -strong embedding with critical point  $\kappa$ . *Strong* means  $\alpha$ -strong for all  $\alpha$ .

Kunen ([7]) showed that no embedding is strong. However a cardinal can be strong, as embeddings witnessing its  $\alpha$ -strength can vary with  $\alpha$ . Stronger properties are obtained by requiring  $j : V \rightarrow M$  to have strength depending on the image under  $j$  of its critical point. For example,  $\kappa$  is *superstrong* iff there is a nontrivial elementary embedding  $j : V \rightarrow M$  with critical point  $\kappa$  which is  $j(\kappa)$ -strong. An important weakening of superstrength is the property that for each  $f : \kappa \rightarrow \kappa$  there is a  $\bar{\kappa} < \kappa$  closed under  $f$  and a nontrivial elementary embedding  $j : V \rightarrow M$  with critical point  $\bar{\kappa}$  which is  $j(f)(\bar{\kappa})$ -strong; such  $\kappa$  are known as *Woodin cardinals*. The consistency strength of the existence of a Woodin cardinal is strictly between that of a strong cardinal and a superstrong cardinal.

We can demand more than superstrength. A cardinal  $\kappa$  is *hyperstrong* iff it is the critical point of an embedding  $j : V \rightarrow M$  which is  $j(\kappa) + 1$ -strong. For a finite  $n > 0$ ,  $n$ -*superstrength* is obtained by requiring  $j$  to be  $j^n(\kappa)$ -strong, where  $j^1 = j$ ,  $j^{k+1} = j \circ j^k$ . Finally,  $\kappa$  is  $\omega$ -*superstrong* iff it is the critical point of an embedding  $j : V \rightarrow M$  which is  $n$ -superstrong for

all  $n$ . Kunen's result [7] shows that no embedding  $j$  with critical point  $\kappa$  is  $j^\omega(\kappa) + 1$ -strong, where  $j^\omega(\kappa)$  is the supremum of the  $j^n(\kappa)$  for finite  $n$ .

*The inner model program*

If  $\kappa$  is inaccessible, then  $\kappa$  is also inaccessible in  $L$ , the most  $L$ -like model of all. This is not the case for measurability, however if  $\kappa$  is measurable then  $\kappa$  is measurable in an inner model  $L[U]$ , where  $U$  is an ultrafilter on  $\kappa$ , which has a definable wellordering and in which GCH,  $\diamond$ ,  $\square$  hold and gap 1 morasses exist. For a strong cardinal  $\kappa$  there is a similarly  $L$ -like inner model  $L[E]$  in which  $\kappa$  is strong, where  $E$  now is not a single ultrafilter, but rather a sequence of generalised ultrafilters, called *extenders*. More recent work yields similar results for Woodin cardinals, and even for Woodin limits of Woodin cardinals (see [9]).

However, progress beyond that has been impeded by the so-called *iterability problem*.

*The outer model program*

How can we obtain  $L$ -like outer models with large cardinals? For inaccessibles one has the following result of Jensen (see [1]):

**Theorem 1.** ( *$L$ -coding*) *There is a generic extension  $V[G]$  of  $V$  such that*

- a. ZFC holds in  $V[G]$ .*
- b.  $V[G] = L[R]$  for some real  $R$ .*
- c. Every inaccessible cardinal of  $V$  remains inaccessible in  $V[G]$ .*

There are similar  $L[U]$  and  $L[E]$  coding theorems (see [4] for the former), providing outer models of the form  $L[U][R]$  and  $L[E][R]$ ,  $R$  a real, which are just as  $L$ -like as  $L[U]$  and  $L[E]$ , preserving measurability and strength, respectively.

However the approach via coding is limited in its use. Obtaining  $L$ -like outer models via coding depends on the existence of  $L$ -like inner models, such as  $L[U]$  or  $L[E]$ , which, as we have observed, are not known to exist beyond Woodin limits of Woodin cardinals. And there are problems with the coding method itself which arise already just past a strong cardinal.

A more promising approach is to use iterated forcing. To illustrate this, consider the problem of making the GCH true in an outer model. Begin with an arbitrary model  $V$  of ZFC. Using forcing, we can add a function from  $\aleph_1$  onto  $2^{\aleph_0}$  without adding reals, thereby making CH true. By forcing again, we add a function from (the possibly new)  $\aleph_2$  onto (the possibly new)

$2^{\aleph_1}$  without adding subsets of  $\aleph_1$ , thereby obtaining  $2^{\aleph_1} = \aleph_2$ . Continue this indefinitely (via a reverse Easton iteration) and the result is a model of the GCH.

Do we preserve large cardinal properties if we make GCH true in this way? The answer is Yes.

**Theorem 2.** *(Large cardinals and the GCH) If  $\kappa$  is superstrong then there is an outer model in which  $\kappa$  is still superstrong and the GCH holds. The same holds for hyperstrong,  $n$ -superstrong for finite  $n$  and  $\omega$ -superstrong.*

*Proof.* First we describe in more detail the above iteration to make GCH true. By induction on  $\alpha$  we define the iteration  $P_\alpha$  of length  $\alpha$ :  $P_0$  is the trivial forcing. For limit  $\lambda$ ,  $P_\lambda$  is the inverse limit of the  $P_\alpha$ ,  $\alpha < \lambda$ , if  $\lambda$  is singular and is the direct limit of the  $P_\alpha$ ,  $\alpha < \lambda$ , if  $\lambda$  is regular. For successor  $\alpha + 1$ ,  $P_{\alpha+1} = P_\alpha * Q_\alpha$ , where  $Q_\alpha$  is the forcing that collapses  $2^{\aleph_\alpha}$  to  $\aleph_{\alpha+1}$  using conditions of size at most  $\aleph_\alpha$ . For any cardinal  $\kappa$  of the form  $\beth_{\alpha+1}$ , the entire iteration  $P$  can be factored as  $P_\kappa * P^\kappa$ , where  $P_\kappa$  has a dense subset of size  $\kappa$  and  $P^\kappa$  is  $\kappa^+$ -closed. In particular, strongly inaccessible cardinals remain strongly inaccessible after forcing with  $P$ .

Now suppose that  $\kappa$  is superstrong, witnessed by the embedding  $j : V \rightarrow M$ , and that  $G$  is  $P$ -generic. Let  $P^*$  denote  $M$ 's version of  $P$ . To show that  $\kappa$  is superstrong in  $V[G]$ , it suffices to find a  $P^*$ -generic  $G^*$  which contains  $j[G]$ , the pointwise image of  $G$  under  $j$ , as a subclass.

Now  $P_\alpha^*$  is the same as  $P_\alpha$  for  $\alpha < j(\kappa)$ , as  $j$  is a superstrong embedding. The first difference between  $P^*$  and  $P$  is at  $j(\kappa)$ :  $P_{j(\kappa)}^*$  is the direct limit of the  $P_\alpha$ ,  $\alpha < j(\kappa)$ , as  $j(\kappa)$  is inaccessible in  $M$ ; but  $j(\kappa)$  is not necessarily regular in  $V$  and therefore it is possible that  $P_{j(\kappa)}^*$  is the *inverse* limit of the  $P_\alpha$ ,  $\alpha < j(\kappa)$ . So we cannot simply choose  $G_{j(\kappa)}^*$  to be  $G_{j(\kappa)}$ , as the latter is generic for the wrong forcing.

But this problem is easily fixed: As  $j(\kappa)$  is in fact Mahlo in  $M$ , it follows that  $P_{j(\kappa)}^*$  has the  $j(\kappa)$ -cc in  $M$ . So any  $G_{j(\kappa)}^*$  contained in  $P_{j(\kappa)}^*$  whose intersection with each  $P_\alpha$ ,  $\alpha < j(\kappa)$ , is  $P_\alpha$ -generic must also be  $P_{j(\kappa)}^*$ -generic. It follows that we can take  $G_{j(\kappa)}^*$  to simply be the intersection of  $G_{j(\kappa)}$  with  $P_{j(\kappa)}^*$ . Notice that  $G_{j(\kappa)}^*$  trivially contains the pointwise image of  $G_\kappa$  under  $j$  as  $j$  is the identity below  $\kappa$ .

Finally we must define a generic  $G^*, j(\kappa)$  for the ‘‘upper part’’  $P^*, j(\kappa)$  of the  $P^*$  iteration, which starts at  $j(\kappa)$  and is defined in the ground model  $M[G_{j(\kappa)}^*]$ . In addition,  $G^*, j(\kappa)$  must contain the pointwise image of  $G^\kappa$

under  $j^*$ , where  $j^*$  is the lifting of  $j$  to  $V[G_\kappa]$  and  $G^\kappa$  is generic for  $P^\kappa$ , an iteration starting at  $\kappa$  defined over the ground model  $V[G_\kappa]$ .

In fact this latter requirement completely determines  $G^*, j^{(\kappa)}$ :

**Lemma 3.**  $j^*[G^\kappa]$  generates a  $P^*, j^{(\kappa)}$ -generic over  $M[G_{j^{(\kappa)}}^*]$ , i.e., each predense subclass of  $P^*, j^{(\kappa)}$  which is definable over  $M[G_{j^{(\kappa)}}^*]$  has an element which is extended by a condition in  $j^*[G^\kappa]$ .

*Proof.* We only consider predense subsets of  $P^*, j^{(\kappa)}$  in  $M[G_{j^{(\kappa)}}^*]$ ; a similar argument works for predense subclasses.

We can assume that  $j : V \rightarrow M$  is given as an extender ultrapower embedding. This means that each element of  $M$  is of the form  $j(f)(a)$ , where  $a$  belongs to  $V_{j^{(\kappa)}}^M = V_{j^{(\kappa)}}$  and  $f$  is a function (in  $V$ ) with domain  $V_\kappa$ . In particular,  $D$  is of the form  $\sigma^{G_{j^{(\kappa)}}^*}$  where the name  $\sigma$  can be written as  $j(f)(a)$  with  $f$  and  $a$  as above. Now using the  $\kappa^+$ -closure of  $P^\kappa$ , choose a condition  $p$  in  $G^\kappa$  which extends an element of  $f(\bar{a})$  whenever  $\bar{a}$  belongs to  $V_\kappa$  and  $f(\bar{a})^{G^\kappa}$  is predense on  $P^*, \kappa$ . Then  $j^*(p)$  obviously belongs to  $j^*[G^\kappa]$  and extends an element of  $j(f)(a)^{G_{j^{(\kappa)}}^*} = \sigma^{G_{j^{(\kappa)}}^*} = D$ , as desired.  $\square$  (Lemma 3)

This completes the construction of  $G^*$  and therefore the proof that  $P$  preserves superstrong cardinals.

Now suppose that  $\kappa$  is hyperstrong. Again we need to find a  $P^*$ -generic  $G^*$  containing  $j[G]$  as a subclass. The forcings  $P_{j^{(\kappa)+1}} = P_{j^{(\kappa)}} * Q_{j^{(\kappa)}}$  and  $P_{j^{(\kappa)+1}}^*$  agree as  $j^{(\kappa)}$  is regular in  $V$  and  $M$  contains  $V_{j^{(\kappa)+1}}$ . Also,  $j^*[g_\kappa]$ , where  $j^*$  is the lifting of  $j$  to  $V[G_\kappa]$  and  $g_\kappa$  is the  $Q_\kappa^{G_\kappa}$ -generic chosen by  $G$  at stage  $\kappa$  of the iteration, is a set of conditions in  $Q_{j^{(\kappa)}}^{G_{j^{(\kappa)}}}$  which belongs to  $M[G_{j^{(\kappa)}}]$  and has size  $2^\kappa$  there; therefore  $j^*[g_\kappa]$  has a lower bound in  $Q_{j^{(\kappa)}}^{G_{j^{(\kappa)}}}$ . By choosing our generic  $G$  so that  $g_{j^{(\kappa)}}$  includes this lower bound, we can succeed in lifting  $j$  to  $V[G_{\kappa+1}]$ . We may assume that  $j : V \rightarrow M$  is given by a hyperextender; this means that each element of  $M$  is of the form  $j(f)(a)$  where  $f$  is a function in  $V$  with domain  $V_{\kappa+1}$  and  $a$  is an element of  $V_{j^{(\kappa)+1}}$ . Then we can use the argument of Lemma 3 to generate the entire generic  $G^*$  containing  $j[G]$ .

The case of  $n$ -superstrongs raises a new difficulty. We first treat the case  $n = 2$ . As in the superstrong case,  $P$  and  $P^*$  may take different limits at  $j^2(\kappa)$ , as the latter may be singular in  $V$ . As in that case, we can obtain a  $P_{j^2(\kappa)}^*$ -generic by intersecting  $G_{j^2(\kappa)}$  with  $P_{j^2(\kappa)}^*$ . However we must also

ensure that  $G_{j^2(\kappa)}^*$  contain  $j[G_{j(\kappa)}]$  as a subset. Write  $P_{j(\kappa)}$  as  $P_\kappa * P_{j(\kappa)}^\kappa$ ; it suffices to have that  $G_{j^2(\kappa)}^*$  contains  $j^*[G_{j(\kappa)}^\kappa]$  as a subset, where  $j^*$  is the lifting of  $j$  to  $V[G_\kappa]$  and  $G_{j(\kappa)}^\kappa$  is  $P_{j(\kappa)}^\kappa$ -generic over  $V[G_\kappa]$ . For the latter we need only know that  $j^2(\kappa)$  has cofinality unequal to  $j(\kappa)$ , for then  $j^*[G_{j(\kappa)}^\kappa]$  is a set of conditions bounded in  $j^2(\kappa)$  and therefore by 2-superstrength belongs to  $M$  and has a lower bound.

**Lemma 4.** *Assume that  $j$  has been chosen so that  $j^2(\kappa)$  is as small as possible. Then  $j^2(\kappa)$  has cofinality  $\kappa^+$ .*

*Proof.* We may assume that the given 2-superstrong embedding  $j : V \rightarrow M$  has the property that each element of  $M$  is of the form  $j(f)(a)$  where  $f$  is a function in  $V$  with domain  $V_{j(\kappa)}$  and  $a$  belongs to  $V_{j^2(\kappa)}$ .

Now let  $H$  consist of all elements of  $M$  of the form  $j(f)(a)$  where  $f$  is a function in  $V$  with domain  $V_\kappa$  and  $a$  belongs to  $V_{j(\kappa)}$ . Then after transitive collapse, the inclusions  $\text{Range } f \subseteq H \subseteq M$  give rise to embeddings  $k : V \rightarrow N$  and  $l : N \rightarrow M$ . The first embedding  $k$  is a superstrong embedding with critical point  $\kappa$  (the ‘‘superstrong embedding derived from  $j$ ’’) and the second embedding  $l$  is an ‘‘external’’ superstrong embedding with critical point  $j(\kappa)$  (‘‘external’’ in the sense that it is not definable in  $N$ ). Also  $k(\kappa) = j(\kappa)$  and  $l(j(\kappa)) = j^2(\kappa)$ .

Now we argue as follows. Let  $h$  be generic over  $N$  for the collapse of  $2^{j(\kappa)}$  to  $j(\kappa)^+$  (using conditions of size at most  $j(\kappa)$ ). Then  $h$  adds a function  $g$  from  $(j(\kappa)^+)^N$  onto  $\{f \in N \mid f : j(\kappa) \rightarrow j(\kappa)\}$  such that  $g \upharpoonright \alpha$  belongs to  $N$  for each  $\alpha < (j(\kappa)^+)^N$ . Now for each  $\alpha < (j(\kappa)^+)^N$  let  $\kappa_\alpha$  be the least cardinal less than  $j^2(\kappa)$  closed under all functions  $l(f)$  for  $f$  in  $g[\alpha]$ . Then as  $j^2(\kappa)$  is regular in  $M$ , each  $\kappa_\alpha$  is less than  $j^2(\kappa)$  and there is an ‘‘external’’ superstrong embedding  $l^*$  with critical point  $j(\kappa)$  such that  $l^*(j(\kappa))$  equals  $\kappa^*$ , the supremum of the  $\kappa_\alpha$ ’s. But then  $(l^*)^{-1} \circ j = j^*$  is a 2-superstrong embedding with critical point  $\kappa$  such that  $(j^*)^2(\kappa) = \kappa^*$ . By the minimality of  $j^2(\kappa)$  it follows that  $\kappa^* = j^2(\kappa)$ , so  $j^2(\kappa)$  has cofinality  $(j(\kappa)^+)^N$  in  $N[h]$  and therefore cofinality  $(j(\kappa)^+)^N$  in  $N$ . As  $N$  is the ultrapower of  $V$  via an extender with critical point  $\kappa$ , it follows that  $(j(\kappa)^+)^N$  has cofinality  $\kappa^+$ , as desired.  $\square$  (Lemma 4)

The previous lemma allows us to lift  $j$  to  $V[G_{j(\kappa)}]$ . Then the method of Lemma 3 can be used to generate the entire generic  $G^*$ .

For the case  $n > 2$  the argument is similar; the general version of Lemma 4 states that  $j^n(\kappa)$  has cofinality  $(j^{n-2}(\kappa))^+$  when it is chosen minimally for an  $n$ -superstrong  $\kappa$ .

Finally we consider  $\omega$ -superstrength. Again we must choose  $G^*$  to be  $P^*$ -generic over  $M$  and to contain the pointwise image of  $G$  under  $j$ . Let  $j^\omega(\kappa)$  denote the supremum of the  $j^n(\kappa)$ ,  $n \in \omega$ . As in Lemma 3 it suffices to find  $G_{j^\omega(\kappa)}^*$  which is  $P_{j^\omega(\kappa)}^*$ -generic and contains  $j[G_{j^\omega(\kappa)}]$  as a subset. Note that  $j[G_\kappa] = G_\kappa$  is trivially contained in  $G_{j^\omega(\kappa)}$  and  $j^*[G_{j^\omega(\kappa)}^\kappa]$  has a lower bound in  $P_{j^\omega(\kappa)}^\kappa$  (as defined in  $V[G_\kappa]$ ); by choosing  $G_{j^\omega(\kappa)}$  to contain this lower bound we may arrange that  $G_{j^\omega(\kappa)}$  contain  $j[G_{j^\omega(\kappa)}]$ . It remains to show:

**Lemma 5.**  $G_{j^\omega(\kappa)} \cap P_{j^\omega(\kappa)}^*$  is  $P_{j^\omega(\kappa)}^*$ -generic over  $M$ .

*Proof.* Suppose that  $D \in M$  is dense on  $P_{j^\omega(\kappa)}^*$  and write  $D$  as  $j(f)(a)$  where  $f$  has domain  $V_{j^\omega(\kappa)}$  and  $a$  belongs to  $V_{j^{n+1}(\kappa)}$  for some  $n$ . (We may assume that every element of  $M$  is of this form.) Choose  $p$  in  $G_{j^\omega(\kappa)}$  such that  $p$  reduces  $f(\bar{a})$  below  $j^n(\kappa)$  whenever  $\bar{a}$  belongs to  $V_{j^n(\kappa)}$  and  $f(\bar{a})$  is open dense on  $P_{j^\omega(\kappa)}$ , in the sense that if  $q$  extends  $p$  then  $q$  can be further extended into  $f(\bar{a})$  without changing  $q$  at or above  $j^n(\kappa)$ . Such a  $p$  exists using the  $j^n(\kappa)^+$ -closure of  $P_{j^\omega(\kappa)}^{j^n(\kappa)}$  in  $V[G_{j^n(\kappa)}]$ . Then  $j(p)$  belongs to  $j[G_{j^\omega(\kappa)}]$  and reduces  $D$  below  $j^{n+1}(\kappa)$ . As  $G_{j^{n+1}(\kappa)}$  is  $P_{j^{n+1}(\kappa)}$ -generic and  $P, P^*$  agree below  $j^{n+1}(\kappa)$ , it follows that  $G_{j^\omega(\kappa)} \cap P_{j^\omega(\kappa)}^*$  intersects  $D$ , as desired.  $\square$  (Lemma 5)

This completes the proof of Theorem 2.  $\square$

Another important property of  $L$  is the existence of a definable wellordering of the universe.

**Theorem 6.** (Large cardinals and definable wellorderings) *If  $\kappa$  is superstrong then there is an outer model in which  $\kappa$  is still superstrong and there is a definable wellordering of the universe. The same holds for hyperstrong,  $n$ -superstrong for finite  $n$  and  $\omega$ -superstrong.*

*Proof.* By Theorem 2 we may assume the GCH. Let  $\kappa$  have one of the large cardinal properties mentioned in the theorem, as witnessed by the embedding  $j : V \rightarrow M$ . Choose  $\lambda$  to be a cardinal greater than  $j^\omega(\kappa)$ . By the method of  $L$ -coding (see Theorem 1), we can enlarge  $V$  without adding subsets of  $\lambda$  to be of the form  $L[A]$ ,  $A$  a subset of  $\lambda^+$ . By the argument of Lemma 3 the embedding  $j$  lifts to  $L[A]$  and therefore  $\kappa$  retains its large cardinal properties.

Now we introduce a definable wellordering. Perform a reverse Easton iteration of length  $\lambda^+$ , indexed by successor cardinals greater than  $\lambda^+$ , where at the  $i$ -th successor cardinal, an  $i^+$ -Cohen set is added iff  $i$  belongs to  $A$ .

The result is that  $i$  belongs to  $A$  iff not every subset of the successor of the  $i$ -th successor cardinal is constructible from a subset of the  $i$ -th successor cardinal. Now the result of this iteration is a model of the form  $L[B]$  where  $B$  is a subset of  $\lambda^{(\lambda^+)}$ . Repeat this to code  $B$  using the next interval of successor cardinals. Continuing this indefinitely yields a model with a wellordering definable from the parameter  $\lambda$ .

To eliminate the parameter  $\lambda$ , use a pairing function  $f : \text{Ord} \times \text{Ord} \rightarrow \text{Ord}$  on the ordinals and arrange that the universe is of the form  $L[C]$  where  $C$  is a class of ordinals and for any  $i$ ,  $i$  is in  $C$  iff some subset of the successor to the  $f(i, j)$ -th successor cardinal is not constructible from a subset of the  $f(i, j)$ -th successor cardinal, for all sufficiently large  $j$ .  $\square$

Jensen's (global)  $\square$  principle asserts the existence of a sequence  $\langle C_\alpha \mid \alpha \text{ singular} \rangle$  such that  $C_\alpha$  has ordertype less than  $\alpha$  for each  $\alpha$  and  $C_{\bar{\alpha}} = C_\alpha \cap \bar{\alpha}$  whenever  $\bar{\alpha} \in \text{Lim } C_\alpha$ . The following strengthens a result of Doug Burke [2]:

**Theorem 7.** (*Superstrong cardinals and  $\square$* ) *If  $\kappa$  is superstrong then there is an outer model in which  $\kappa$  is still superstrong and  $\square$  holds.*

*Proof.* By Theorem 2 we may assume the GCH. Consider now the reverse Easton iteration  $P$  where at the regular stage  $\alpha$ ,  $Q_\alpha$  is a  $P_\alpha$ -name for the forcing which adds a  $\square$ -sequence on the singular limit ordinals less than  $\alpha$ . A condition in  $Q_\alpha$  is a sequence  $\langle C_\beta \mid \beta \leq \gamma, \beta \text{ singular} \rangle$ ,  $\gamma < \alpha$ , such that  $C_\beta$  has ordertype less than  $\beta$  for each  $\beta$  and  $C_{\bar{\beta}} = C_\beta \cap \bar{\beta}$  whenever  $\bar{\beta}$  belongs to  $\text{Lim } C_\beta$ .

Using the fact that  $P_\alpha$  forces  $\square$ -sequences of any regular length less than  $\alpha$ , it is easy to verify by induction that any condition in  $Q_\alpha$  can be extended to have arbitrarily large length less than  $\alpha$ . Also  $Q_\alpha$ , and indeed the entire iteration from stage  $\alpha$  on, is  $\alpha$ -distributive.

Let  $P^*$  denote  $M$ 's version of  $P$ . We want to construct  $G^*$  to be  $P^*$ -generic over  $M$  and to contain  $j[G]$  as a subclass. As in earlier arguments,  $P$  and  $P^*$  agree strictly below  $j(\kappa)$  but not necessarily at  $j(\kappa)$ , which is regular in  $M$  but may be singular in  $V$ ; as before we take  $G_{j(\kappa)}^*$  to be  $G_{j(\kappa)} \cap P_{j(\kappa)}^*$ . Our new task is to define a  $Q_{j(\kappa)}$ -generic  $g$  over  $M[G_{j(\kappa)}^*]$ .

Assume that  $j : V \rightarrow M$  is chosen with  $j(\kappa)$  minimal. Then by the argument of Lemma 4,  $j(\kappa)$  has cofinality  $\kappa^+$ . We can assume that  $j$  is given by an ultrapower, and therefore that  $(j(\kappa)^+)^M$  also has cofinality  $\kappa^+$ . Now we can build  $g$  in  $\kappa^+$  steps, using  $j(\kappa)$ -distributivity to meet fewer than

$j(\kappa)$  open dense sets at each step. We must also ensure that  $g$  extend  $g_\kappa$ ; but this is easy to arrange as the latter is a condition in the forcing  $Q_{j(\kappa)}$ .

Finally the rest of  $G^*$  can be generated from  $j[G]$  as in Lemma 3.  $\square$

The proof of the previous theorem does not work for hyperstrong  $\kappa$ , and there is a good reason for this.  $\kappa$  is *subcompact* iff for any  $B \subseteq H_{\kappa^+}$  there are  $\mu < \kappa$ ,  $A \subseteq H_{\mu^+}$  and an elementary embedding  $j : (H_{\mu^+}, A) \rightarrow (H_{\kappa^+}, B)$  with critical point  $\mu$ . (Note that by elementarity,  $j$  must send  $\mu$  to  $\kappa$ .)

**Proposition 8.** (a) *If  $\kappa$  is hyperstrong then  $\kappa$  is subcompact.* (b) (Jensen) *If there is a subcompact cardinal then  $\square$  (even when restricted to ordinals between  $\kappa$  and  $\kappa^+$ ) fails.*

*Proof.* (a) Suppose that  $j : V \rightarrow M$  witnesses hyperstrength. Then for all subsets  $B$  of  $j(\kappa)^+$  in the range of  $j$ ,  $j$  gives an elementary embedding of  $(H_{\kappa^+}, A)$  into  $(H_{j(\kappa)^+}, B)$ , where  $j(A) = B$ ; moreover this embedding belongs to  $M$  as  $j$  is hyperstrong and  $j \upharpoonright H_{\kappa^+}$  belongs to  $H_{j(\kappa)^+}$ . As the range of  $j$  is an elementary submodel of  $M$ , it follows that there is an elementary embedding of some  $(H_{\mu^+}, A)$  into  $(H_{j(\kappa)^+}, B)$  (sending  $\mu$  to  $j(\kappa)$ ) which belongs to the range of  $j$ . So  $j(\kappa)$  is subcompact in  $\text{Range } j$  and therefore by elementarity subcompact in  $M$ . As  $j$  is elementary,  $\kappa$  is subcompact in  $V$ .

(b) Suppose that  $\kappa$  is subcompact and  $\vec{C} = \langle C_\alpha \mid \kappa < \alpha < \kappa^+, \alpha \text{ singular} \rangle$  has the properties of a  $\square$ -sequence. By thinning out the  $C_\alpha$ 's we can ensure that each has ordertype at most  $\kappa$ . Let  $j$  be an embedding from  $(H_{\mu^+}, \vec{C})$  to  $(H_{\kappa^+}, \vec{C})$ , sending  $\mu$  to  $\kappa$ . Let  $\alpha$  be the supremum of the ordinals in the range of  $j$ . Then  $\alpha$  has cofinality  $\mu^+$ . The ordinals in the range of  $j$  form a  $< \mu$ -closed and therefore  $\omega$ -closed unbounded subset of  $\alpha$ . And  $\text{Lim } C_\alpha$  is a closed unbounded subset of  $\alpha$ . Therefore the intersection  $D$  of these two sets is unbounded in  $\alpha$ . By the coherence property of  $\vec{C}$ , the ordertype of  $C_\beta$  for sufficiently large  $\beta$  in  $D$  is at least  $\mu$ . But as the ordertype of  $C_\alpha$  is at most  $\kappa$  (in fact less than  $\kappa$ ), the ordertype of  $C_\beta$  for all  $\beta$  in  $D$  is strictly less than  $\kappa$ . Thus there are  $\beta$  in  $D \subseteq \text{Range } j$  with  $C_\beta$  of ordertype not in  $\text{Range } j$ , contradicting the elementarity of  $j$ .  $\square$

For uncountable, regular  $\kappa$ ,  $\diamond_\kappa$  says that there exists  $\langle D_\alpha \mid \alpha < \kappa \rangle$  such that  $D_\alpha$  is a subset of  $\alpha$  for each  $\alpha$  and for every subset  $D$  of  $\kappa$ ,  $\{\alpha < \kappa \mid D_\alpha = D \cap \alpha\}$  is stationary in  $\kappa$ .  $\diamond$  asserts that  $\diamond_\kappa$  holds for every uncountable, regular  $\kappa$ .

**Theorem 9.** (Large cardinals and  $\diamond$ ) *If  $\kappa$  is superstrong then there is an outer model in which  $\kappa$  is still superstrong and  $\diamond$  holds. The same holds for hyperstrong,  $n$ -superstrong for finite  $n$  and  $\omega$ -superstrong.*

*Proof.* The proof combines the proofs of Theorems 2 and 7. As in the latter proof, we use a reverse Easton iteration  $P$  where at each regular stage  $\alpha$ ,  $Q_\alpha$  is  $\alpha$ -distributive (and in fact in the present context, the entire iteration starting with  $\alpha$  is  $\alpha$ -closed). In this case, a condition in  $Q_\alpha$  is a sequence  $\langle D_\beta \mid \beta < \gamma \rangle$ ,  $\gamma < \alpha$ , such that  $D_\beta$  is a subset of  $\beta$  for each  $\beta < \gamma$ . It is easy to show that a  $Q_\alpha$ -generic yields a  $\diamond_\alpha$ -sequence, using the  $\alpha$ -closure of  $Q_\alpha$ .

The proof in the superstrong case is just as in Theorem 7. For hyperstrong  $\kappa$  (witnessed by  $j : V \rightarrow M$ ), we need only observe that if  $g_{\kappa^+}$  is  $Q_{\kappa^+}$ -generic then  $j[g_{\kappa^+}]$  has a lower bound in the forcing  $Q_{j(\kappa)^+}$ . (This is where the argument with  $\square$  breaks down.)

For  $n$ -superstrongs,  $2 < n$  finite, we face the problem of building a  $Q_{j^n(\kappa)}$ -generic containing the image of  $g_{j^{n-1}(\kappa)}$  under (the lifting to  $V[G_{j^{n-1}(\kappa)}]$  of)  $j$ . The latter image has a lower bound in  $Q_{j^n(\kappa)}$ , so the only difficulty is the construction of a  $Q_{j^n(\kappa)}$ -generic. We cannot use the argument of Theorem 7, as  $j^n(\kappa)$  (when minimised) has cofinality  $(j^{n-2}(\kappa))^+$  whereas its cardinal successor in  $M$  has cofinality  $(j^{n-1}(\kappa))^+$ . But using the  $j^n(\kappa)$ -closure of the forcing, we can meet all predense sets with names of the form  $j(f)(a)$  for  $j^{n-1}(\kappa)$ -many  $f : V_{j^{n-1}(\kappa)} \rightarrow V_{j^{n-1}(\kappa)+1}$ 's and fewer than  $j^n(\kappa)$ -many  $a$ 's from  $V_{j^n(\kappa)}$ . Fixing the set of  $a$ 's, we can in  $(j^{n-1}(\kappa))^+$  steps meet all predense sets with names of the form  $j(f)(a)$ ,  $f$  arbitrary. Then we can repeat this in  $(j^{n-2}(\kappa))^+$  steps for larger and larger sets of  $a$ 's, thereby meeting all predense sets in  $M[G_{j^n(\kappa)}]$ .

Finally,  $\omega$ -superstrength is handled just as in Theorem 2.  $\square$

**Theorem 10.** (*Large cardinals and Gap 1 morasses*) *If  $\kappa$  is superstrong then there is an outer model in which  $\kappa$  is still superstrong and gap 1 morasses exist at each regular cardinal. The same holds for hyperstrong,  $n$ -superstrong for finite  $n$  and  $\omega$ -superstrong.*

*Proof.* For the definition of a gap 1 morass we refer the reader to [3]. Assume GCH and let  $\kappa$  be superstrong. We apply the reverse Easton iteration  $P$  where at each regular stage  $\alpha$ ,  $Q_\alpha$  adds a gap 1 morass at  $\alpha$ . A condition in  $Q_\alpha$  is a size  $< \alpha$  initial segment of a morass up to some top level, together with a map of an initial segment of this top level into  $\alpha^+$  which obeys the requirements of a morass map. To extend a condition, we end-extend the morass up to its top level and require that the map from the given initial segment of its top level into  $\alpha^+$  factor as the composition of a map into the top level of the stronger condition followed by the map given by the stronger condition into  $\alpha^+$ . The forcing  $Q_\alpha$  is  $\alpha$ -closed and, using a  $\Delta$ -system argument, is  $\alpha^+$ -cc.

To obtain the desired  $G^*$ , we must build a  $Q_{j(\kappa)}$ -generic which extends the image under  $j$  of the  $Q_\kappa$ -generic  $g_\kappa$ . As in the case of  $\square$  we use minimisation of  $j(\kappa)$  to ensure that it has cofinality  $\kappa^+$  and then build a  $Q_{j(\kappa)}$ -generic in  $\kappa^+$  steps. Note that any condition in  $j[g_\kappa]$  is extended by one which has top level  $\kappa$  and maps an initial segment of the top level into  $j(\kappa)^+$  using  $j$ . Now given fewer than  $j(\kappa)$  maximal antichains in  $M[G_{j(\kappa)}]$ , we can choose  $\alpha < j(\kappa)^+$  of cofinality  $j(\kappa)$  in  $M$  so that these maximal antichains are maximal when restricted to conditions which are “below  $\alpha$ ” in the sense that they map an initial segment of their top level into  $\alpha$ . Moreover, there is a condition which serves as a lower bound to all conditions in  $j[g_\kappa]$  which are below  $\alpha$  in this sense. Therefore we can choose a condition below  $\alpha$  meeting all of the given maximal antichains compatibly with the conditions in  $j[g_\kappa]$  which are below  $\alpha$ , and therefore compatibly with all conditions in  $j[g_\kappa]$ . Repeating this in  $\kappa^+$  steps for increasingly large  $\alpha < j(\kappa)^+$  of  $M$ -cofinality  $j(\kappa)$  yields the desired  $Q_{j(\kappa)}$ -generic. The remainder of the generic  $G^*$  can be formed using Lemma 3.

Now suppose that  $\kappa$  is hyperstrong. We may assume that  $j$  is given by a hyperextender and therefore  $j$  is cofinal from  $\kappa^{++}$  into  $j(\kappa)^{++}$  of  $M$ . Let  $S$  consist of those morass points at the top level (i.e., level  $\kappa^+$ ) of  $g_{\kappa^+}$  which have cofinality  $\kappa^+$ . For each  $\sigma$  in  $S$  let  $g_{\kappa^+} \upharpoonright \sigma$  denote the set of conditions in  $g_{\kappa^+}$  which are below  $\sigma$ . Then  $j[g_{\kappa^+} \upharpoonright \sigma]$  has a greatest lower bound  $p_\sigma$  in  $Q_{j(\kappa)}$ .

The collection of maximal antichains of  $Q_{j(\kappa)}$  which belong to  $M$  can be written as a union  $\bigcup_{i < j(\kappa)^+} X_i$  where for each  $i$  and each  $\sigma$ ,  $X_i \upharpoonright j(\sigma)$  (the subset of  $X_i$  consisting of those maximal antichains all of whose elements are below  $j(\sigma)$ ) is a set of size at most  $j(\kappa)$  in  $M$ . By induction on  $\sigma \in S$  choose a condition  $q_\sigma$  extending  $p_\sigma$  and all  $q_\tau$ ,  $\tau \in S \cap \sigma$ , which meets all antichains in  $X_0 \upharpoonright j(\sigma)$ . By hyperstrength, the sequence of  $q_\tau \upharpoonright j(\sigma)$  has a greatest lower bound  $p_\sigma^1$  for each  $\sigma \in S$ . Now repeat this construction for  $X_1, X_2, \dots$  for  $j(\kappa)^+$  steps, resulting in a set of conditions which generates a generic  $g_{j(\kappa)^+}$  for  $Q_{j(\kappa)^+}$ . As before, the remainder of the generic  $G^*$  can be generated using Lemma 3.

The cases of  $n$ -superstrength,  $2 \leq n$  finite, are handled using the above argument together with the proof of Theorem 9.  $\omega$ -superstrength is handled as in the proof of Theorem 2.  $\square$

*Questions.* 1. The proofs of Theorems 2 and 7 show that one can force the GCH and  $\square$  preserving the superstrength of *all* superstrong cardinals and GCH preserving the hyperstrength of *all* hyperstrong cardinals. Is it

possible to force GCH preserving the 2-superstrength of all 2-superstrong cardinals?

2. It is possible to force a definable wellordering of the universe over a model of GCH preserving the superstrength of all superstrong cardinals, at the cost of some cardinal collapsing. Is it possible to do this without cardinal collapsing? Is it possible to preserve the superstrength of all superstrong cardinals while forcing not only the universe but also each  $H(\kappa)$ ,  $\kappa > \omega_1$ , to have a definable wellordering?

3. Is it consistent with a superstrong cardinal to have a gap 2 morass at every regular cardinal?

4. To what extent are the condensation and hyperfine structural properties of  $L$  (see [6]) consistent with large cardinals?

#### REFERENCES

- [1] Beller, A., Jensen, R. and Welch, P. *Coding the Universe*, London Math Society Lecture Note Series, **47** (1982) Cambridge University Press.
- [2] Burke, D. Generic embeddings and the failure of Box, Proceedings of the American Mathematical Society, Vol. 123, No. 9, pp. 2867–2871, 1995.
- [3] Devlin, K. *Constructibility*, Springer-Verlag, 1984.
- [4] Friedman, S. Coding over a measurable cardinal, Journal of Symbolic Logic Vol. 54, No. 4, pp. 1145–1159, 1989.
- [5] Friedman, S. *Fine structure and class forcing*, de Gruyter, 2000.
- [6] Friedman, S. and Koepke, P. An elementary approach to the fine structure of  $L$ , Bulletin of Symbolic Logic, Vol. 3, No. 4, pp. 453–468, 1997.
- [7] Kunen, K. Elementary embeddings and infinitary combinatorics, Journal of Symbolic Logic 36, pp. 407–413, 1971.
- [8] Mitchell, W. An introduction to inner models and large cardinals, a chapter for the Handbook of Set Theory.
- [9] Neeman, I. Inner models in the region of a Woodin limit of Woodin cardinals, Annals of Pure and Applied Logic, Vol. 116, pp. 67–115, 2002.
- [10] Zeman, M. *Inner models and large cardinals*, de Gruyter Series in Logic and Its Applications Vol.5, 2002.