



CENTRE DE RECERCA MATEMÀTICA

Preprint núm. 1206

May 2015

Cylinders for non-symmetric
DG-operads via homological
perturbation theory

F. Muro

CYLINDERS FOR NON-SYMMETRIC DG-OPERADS VIA HOMOLOGICAL PERTURBATION THEORY

FERNANDO MURO

ABSTRACT. We construct small cylinders for cellular non-symmetric DG-operads over an arbitrary commutative ring by using the basic perturbation lemma from homological algebra. We show that our construction, applied to the A -infinity operad, yields the operad parametrizing A -infinity maps whose linear part is the identity. We also compute some other examples with non-trivial operations in arities 1 and 0.

INTRODUCTION

Cylinders are basic tools to do homotopy theory in any context. The existence of cylinders is guaranteed by the axioms of model categories. In cofibrantly generated model categories, cylinders can be constructed out of generating cofibrations, but they are huge, not useful for explicit computations. In many specific model categories, there are nice and small cylinders for cofibrant objects, e.g. topological spaces (the product with the interval), chain complexes (well known), differential graded (DG) algebras, and connected commutative DG-algebras in characteristic 0, see [Bau89, §1].

Any DG-algebra has a cofibrant resolution of the form $(T(V), d + \partial)$, where $(T(V), d)$ is the free DG-algebra on a chain complex V and ∂ is a perturbation of the differential d . The cylinder of such a DG-algebra is $(T(IV), d + \partial_I)$, where $(T(IV), d)$ is the free DG-algebra on the cylinder IV of the chain complex V and ∂_I is a perturbation defined from ∂ in a straightforward way, using a chain homotopy compatible with the associative algebra structure. The commutative case starts similarly, with a perturbed free commutative DG-algebra, but quickly diverges. The differential is defined in terms of locally nilpotent derivations and formal exponentials, where factorial denominators appear.

DG-operads are closely related to DG-algebras, the arity 1 part $\mathcal{O}(1)$ of a DG-operad \mathcal{O} is a DG-algebra. However, one of the DG-operad laws contains a switch,

$$(x \circ_i y) \circ_j z = (-1)^{|y||z|} (x \circ_j z) \circ_{i+n-1} y, \quad j < i, \quad z \in \mathcal{O}(n).$$

2010 *Mathematics Subject Classification.* 18D50, 18G55.

Key words and phrases. Operad, cylinder, homotopy.

Therefore, in any possible construction of cylinders for DG-operads we can expect some of the complications of the commutative DG-algebra case. We work with non-symmetric DG-operads, which avoids further switches associated to symmetric group actions. Unlike in the symmetric case, the category of non-symmetric DG-operads is endowed with a model structure over any commutative ground ring [Mur11, Lyu11].

Any DG-operad has a cofibrant resolution of the form $(\mathcal{F}(V), d + \partial)$, where $(\mathcal{F}(V), d)$ is the free DG-operad on a sequence of chain complexes $V = \{V(n)\}_{n \geq 0}$ and ∂ is a perturbation of the differential d . We construct a cylinder for such a DG-operad of the form $(\mathcal{F}(IV), d + \partial_I)$, where $(\mathcal{F}(IV), d)$ is the free DG-operad on the sequence of chain cylinders $IV = \{IV(n)\}_{n \geq 0}$ and ∂_I is a perturbation defined from ∂ by using tools from homological perturbation theory. The definition of ∂_I is recursive. We compute some examples in Sections 2 and 3. The most remarkable one is the A -infinity operad \mathcal{A}_∞ , which has the previous form. Maps from \mathcal{A}_∞ to an endomorphism operad correspond to A -infinity algebra structures, and homotopies with respect to our cylinder correspond to A -infinity morphisms whose linear part is the identity.

Fresse [Fre09] defined cylinders for symmetric DG-operads arising as the cobar construction of a coaugmented connected DG-cooperad. The A -infinity operad \mathcal{A}_∞ arises in this way, it is the cobar construction of the Koszul dual cooperad of the associative operad. Fresse's cylinder of \mathcal{A}_∞ coincides with ours, modulo symmetrization. His formulas are closed, not recursive, and work for symmetric operads. Our construction does not rule out operads with non-trivial operations in arities 0 and 1, such as the unital A -infinity operad [HM12, Lyu11, MT14] or the DG-operad for homotopy associative algebras with derivation [Lod10], considered in Example 2.20, and works straightaway in the relative case. We actually describe the cylinder of a DG-operad concentrated in arities 0 and 1, observing that our construction generalizes the classical cylinder of DG-algebras. In the final section, we consider a family of DG-operads, called linear, where our formulas greatly simplify. This family contains interesting examples, mainly in the relative case.

Acknowledgments. The author was partially supported by the Andalusian Ministry of Economy, Innovation and Science under the grant FQM-5713 and by the Spanish Ministry of Economy under the MEC-FEDER grant MTM2013-42178-P. This article was finished at the Centre de Recerca Matemàtica while the author was taking part in the Research program IRTATCA: Interactions between Representation Theory, Algebraic Topology and Commutative Algebra. He is very grateful to this institution and to the organizers of the program for excellent working conditions and research atmosphere.

1. CYLINDERS

We work over an arbitrary commutative ground ring \mathbb{k} . The symmetry constraint in the monoidal categories of $(\mathbb{Z}$ -)graded $(\mathbb{k}$ -)modules and chain complexes uses the Koszul sign rule. Differentials have degree -1 . As a graded module, the suspension ΣX of a chain complex X is $(\Sigma X)_n = X_{n-1}$ with differential $d_{\Sigma X} = -d_X$. We denote by $\sigma: X \rightarrow \Sigma X$ the degree $+1$ isomorphism that is the identity degreewise, which satisfies $d_{\Sigma X}\sigma = -\sigma d_X$. Therefore, given $x \in X_{n-1}$ we often write $\sigma(x)$ for x itself regarded as an element in $(\Sigma X)_n$.

Definition 1.1. A *strong deformation retraction*, or simply an *SDR*, consists of two chain complexes X and Y and a diagram

$$X \begin{array}{c} \xleftarrow{i} \\ \xrightarrow{p} \end{array} Y \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h$$

where i and p are chain maps, h is a chain homotopy from ip to the identity in Y , i.e. a degree $+1$ map satisfying

$$ip - 1_Y = dh + hd,$$

and the following equations are satisfied,

$$pi = 1_X, \quad ph = 0, \quad hi = 0, \quad h^2 = 0.$$

A *strong pseudo-cylinder* of a chain complex X is a diagram

$$X \begin{array}{c} \xrightarrow{i_0} \\ \xrightarrow{i_1} \end{array} C \begin{array}{c} \xrightarrow{p} \\ \xrightarrow{p} \end{array} X \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h$$

where C is a chain complex, i_0, i_1 and p are chain maps, h is a chain homotopy such that

$$X \begin{array}{c} \xleftarrow{i_0} \\ \xrightarrow{p} \end{array} C \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h$$

is an SDR, and $pi_1 = 1_X$. We call it a *strong cylinder* if $(i_0, i_1): X \oplus X \rightarrow C$ is a cofibration in the projective model structure on chain complexes [Hov99, Theorem 2.3.11], and hence the underlying diagram obtained by forgetting h is a cylinder object factorization in the model theoretic sense.

The *canonical strong pseudo-cylinder* of a chain complex X is given by

$$X \begin{array}{c} \xrightarrow{i_0} \\ \xrightarrow{i_1} \end{array} IX \begin{array}{c} \xrightarrow{p} \\ \xrightarrow{p} \end{array} X \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h_I$$

where IX , as a graded module, is

$$IX = X \oplus \Sigma X \oplus X.$$

The differential of IX is

$$d_{IX} = \begin{pmatrix} d_X & 1 & 0 \\ 0 & -d_X & 0 \\ 0 & -1 & d_X \end{pmatrix}$$

and the structure maps are

$$i_0 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad i_1 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad p = (1 \ 0 \ 1), \quad h_I = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The canonical strong pseudo-cylinder is a strong cylinder if and only if X is a cofibrant chain complex.

Definition 1.2. In this paper, all operads are non-symmetric. Hence, a graded operad or DG-operad \mathcal{O} is a sequence of objects $\{\mathcal{O}(n)\}_{n \geq 0}$ equipped with structure maps

$$\begin{aligned} \mathcal{O}(n) \otimes \mathcal{O}(p_1) \otimes \cdots \otimes \mathcal{O}(p_n) &\longrightarrow \mathcal{O}(p_1 + \cdots + p_n), & n \geq 1, \quad p_1, \dots, p_n \geq 0, \\ x_0 \otimes x_1 \otimes \cdots \otimes x_n &\mapsto x_0(x_1, \dots, x_n), \end{aligned}$$

satisfying

$$(1.3) \quad x_0(x_1, \dots, x_n)(y_1, \dots, y_{p_1+\dots+p_n}) = (-1)^\epsilon x_0(x_1(y_1, \dots, y_{p_1}), \dots, x_n(y_{p_1+\dots+p_{n-1}+1}, \dots, y_{p_1+\dots+p_n})),$$

where the sign is simply determined by the Koszul rule

$$\epsilon = \sum_{i=2}^n \sum_{j=1}^{\sum_{k=1}^{i-1} p_k} |x_i| |y_j|,$$

and an identity element $\text{id} = \text{id}_{\mathcal{O}} \in \mathcal{O}(1)$ satisfying

$$\text{id}(x) = x = x(\text{id}, \dots, \text{id}).$$

We call $\mathcal{O}(n)$ the *arity* n part of \mathcal{O} . In the case of DG-operads, the structure maps being chain maps translates in the *operadic Leibniz rule*,

$$d(x_0(x_1, \dots, x_n)) = d(x_0)(x_1, \dots, x_n) + \sum_{i=1}^n (-1)^{\sum_{j=0}^{i-1} |x_j|} x_0(\dots, x_{i-1}, d(x_i), x_{i+1}, \dots),$$

which implies $d(\text{id}) = 0$. In $x_0(x_1, \dots, x_n)$ we usually omit those x_i , $1 \leq i \leq n$, which are $x_i = \text{id}$. Operads can also be defined in terms of *operadic compositions*,

$$\circ_i: \mathcal{O}(p) \otimes \mathcal{O}(q) \longrightarrow \mathcal{O}(p+q-1), \quad 1 \leq i \leq p, \quad q \geq 0, \quad x \circ_i y = x^{(i-1, \cdot)} y^{(\cdot, i)}.$$

In this case, the laws are

$$(1.4) \quad \begin{aligned} x \circ_i (y \circ_j z) &= (x \circ_i y) \circ_{i+j-1} z; \\ (x \circ_i y) \circ_j z &= (-1)^{|y||z|} (x \circ_j z) \circ_{i+n-1} y, \quad j < i, \quad z \in \mathcal{O}(n); \\ \text{id} \circ_1 x &= x = x \circ_i \text{id}; \end{aligned}$$

and, for DG-operads, the operadic Leibniz rule is equivalent to

$$(1.5) \quad d(x \circ_i y) = d(x) \circ_i y + (-1)^{|x|} x \circ_i d(y).$$

Definition 1.6. A *strong pseudo-cylinder* of a DG-operad \mathcal{O} is a sequence of strong pseudo-cylinders of chain complexes

$$\mathcal{O} \begin{array}{c} \xrightarrow{i_0} \\ \xleftarrow{i_1} \end{array} \mathcal{P} \xrightarrow{p} \mathcal{O} \quad \begin{array}{c} \curvearrowright \\ h \end{array}$$

such that \mathcal{P} is a DG-operad and i_0, i_1 and p are DG-operad maps. We call it a *strong cylinder* if $(i_0, i_1): \mathcal{O} \amalg \mathcal{O} \rightarrow \mathcal{P}$ is a cofibration in the model structure on DG-operads transferred from the projective model structure on chain complexes, see [Mur11, Theorem 1.1] or [Lyu11, Proposition 1.8]. Abusing terminology, we sometimes say that \mathcal{P} is a strong (pseudo-)cylinder of \mathcal{O} . We sometimes write $h = h_{\mathcal{O}}$ in order to avoid ambiguity.

No compatibility condition with the DG-operad structure is required for the chain homotopy h . Many explicit constructions below will however satisfy $h(\text{id}) = 0$. There seems to be no canonical strong pseudo-cylinders for arbitrary DG-operads. The aim of this paper is to construct nice strong pseudo-cylinders for a big class of quasi-free DG-operads, i.e. DG-operads with free underlying graded operad. Nice means that they are strong cylinders under some extra assumptions, like canonical strong pseudo-cylinders of chain complexes. As a toy example, we will start with honestly free DG-operads.

We consider planted planar trees with leaves, that we simply call *trees*, and endow the set of inner vertices with the path order. Two vertices satisfy $v \leq w$ if, when we draw the paths from the root to v and w , the path to v bifurcates to the left or is contained in the path to w , e.g. in the following tree the path order is indicated by the subscripts




see [Mur11, §3] for formal definitions. The *arity* of a vertex v is the number \tilde{v} of edges adjacent from above, e.g. in the previous tree $\tilde{v}_1 = 2, \tilde{v}_2 = 3, \tilde{v}_3 = 0, \tilde{v}_4 = 3,$ and $\tilde{v}_5 = 1$. We just depict *inner vertices*, i.e. we do not draw the top vertices of the leaves or the bottom vertex of the root. An *inner edge* is an edge which is neither a leaf nor the root, i.e. such that the two adjacent vertices are inner vertices.

Given a sequence of graded modules or chain complexes $V = \{V(n)\}_{n \geq 0}$ and a tree T , we denote

$$V(T) = \bigotimes_v V(\tilde{v}).$$

This tensor product is indexed by the inner vertices of T , and it is taken in the path order. A tensor in $V(T)$ is usually denoted by labeling each inner vertex v with an element in $V(\tilde{v})$, e.g.



Labeled trees are also used to denote iterated compositions in operads. There is only one bracketing compatible with the path order, e.g. in the previous tree $x_1(x_2(-, x_3, x_4), x_5)$. This bracketing, that we call *nested*, is the only one with no


$$\dots)(\dots$$

We can nest any bracketing by iterating (1.3). As we will next see, the previous labeled tree can be regarded as an iterated composition in the free operad.

The underlying sequence of the free operad $\mathcal{F}(V)$ spanned by a sequence of graded modules or chain complexes V is

$$\mathcal{F}(V)(n) = \bigoplus_T V(T),$$

see for instance [Mur11, §5]. This direct sum is indexed by the trees with n leaves. The natural sequence of maps $V \rightarrow \mathcal{F}(V)$ is the inclusion of the direct summands indexed by corollas, $n \geq 0$,



The composition law \circ_i is given by the symmetry isomorphisms $V(T) \otimes V(T') \cong V(T \circ_i T')$, where $T \circ_i T'$ is the tree obtained by grafting T' in the i^{th} leaf of T (notice that the inner vertices of $T \circ_i T'$ are the disjoint union of the inner vertices of T and T'). Using labeled trees, this is just grafting up to sign determined by the path order and the Koszul rule. The identity element is $\text{id} = 1 \in V(|) = \mathbb{k}$.

Lemma 1.8 ([Lam93, Lemma 2.3]). *Given two SDRs*

$$X \begin{array}{c} \xrightarrow{i} \\ \xleftarrow{p} \end{array} Y \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h, \quad X' \begin{array}{c} \xrightarrow{i'} \\ \xleftarrow{p'} \end{array} Y' \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h',$$

the following diagram is also an SDR, that we call tensor product SDR,

$$X \otimes X' \begin{array}{c} \xrightarrow{i \otimes i'} \\ \xleftarrow{p \otimes p'} \end{array} Y \otimes Y' \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h \otimes 1_{Y'} + i p \otimes h' .$$

Tensor product SDRs behave well with respect to associativity constraints, so we can define tensor products of several SDRs by iteration without specifying a bracketing. However, this construction is not symmetric, so the order of tensor factors does matter. This is why we insisted on the path order in the definition of free operads. We could have equally worked with the chain homotopy $h \otimes i'p' + 1_Y \otimes h'$, but we must fix some convention, and we have decided to fix that in Lemma 1.8.

Tensor products of strong pseudo-cylinders of chain complexes are defined by using tensor products of SDRs in the obvious way. The canonical strong pseudo-cylinder of a sequence of chain complexes is defined aritywise as in Definition 1.1. The empty tensor product is the trivial strong pseudo-cylinder of \mathbb{k} , regarded as a chain complex concentrated in degree 0,

$$\mathbb{k} \begin{array}{c} \xrightarrow{1} \\ \xrightarrow{1} \end{array} \mathbb{k} \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \mathbb{k} \xrightarrow{1} \mathbb{k} .$$

Definition 1.9. The *canonical strong pseudo-cylinder* of a free DG-operad $\mathcal{F}(V)$ on a sequence V of chain complexes

$$\mathcal{F}(V) \begin{array}{c} \xrightarrow{\mathcal{F}(i_0)} \\ \xrightarrow{\mathcal{F}(i_1)} \end{array} \mathcal{F}(IV) \begin{array}{c} \xrightarrow{\mathcal{F}(p)} \\ \xrightarrow{\mathcal{F}(p)} \end{array} \mathcal{F}(V)$$

is defined on each $V(T)$ as the tensor product (in the path order) of the canonical strong pseudo-cylinders of the sequence of chain complexes V .

The empty tensor product appears when $T = |$, hence $h_V(\text{id}) = 0$.

Remark 1.10. It is very easy to evaluate h_V on labeled trees, e.g.

$$h_V \left(\begin{array}{c} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \\ \begin{array}{c} x_3 \quad x_4 \\ x_2 \quad x_5 \\ x_1 \end{array} \end{array} \right) = \begin{array}{c} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \\ \begin{array}{c} x_3 \quad x_4 \\ x_2 \quad x_5 \\ h_I(x_1) \end{array} \end{array} + (-1)^{|x_1|} \begin{array}{c} \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \\ \begin{array}{c} x_3 \quad x_4 \\ x_2 \quad x_5 \\ i_0 p(x_1) \end{array} \end{array}$$

$$\begin{aligned}
& + (-1)^{|x_1|+|x_2|} \begin{array}{c} \begin{array}{c} h_I(x_3) \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ i_0p(x_2) \quad x_4 \\ \diagdown \quad \diagup \\ i_0p(x_1) \quad x_5 \end{array} \end{array} + (-1)^{|x_1|+|x_2|+|x_3|} \begin{array}{c} \begin{array}{c} i_0p(x_3) \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ i_0p(x_2) \quad h_I(x_4) \\ \diagdown \quad \diagup \\ i_0p(x_1) \quad x_5 \end{array} \end{array} \\
& + (-1)^{|x_1|+|x_2|+|x_3|+|x_4|} \begin{array}{c} \begin{array}{c} i_0p(x_3) \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ i_0p(x_2) \quad i_0p(x_4) \\ \diagdown \quad \diagup \\ i_0p(x_1) \quad h_I(x_5) \end{array} \end{array} .
\end{aligned}$$

In terms of formulas, given $x_0 \in IV(n)$ and $x_1, \dots, x_n \in \mathcal{F}(IV)$,

(1.11)

$$\begin{aligned}
h_V(x_0(x_1, \dots, x_n)) &= h_I(x_0)(x_1, \dots, x_n) \\
&+ \sum_{i=1}^n (-1)^{\sum_{j=0}^{i-1} |x_j|} i_0p(x_0)(i_0p(x_1), \dots, i_0p(x_{i-1}), h_V(x_i), x_{i+1}, \dots, x_n).
\end{aligned}$$

This yields a straightforward way of computing h_V on nested bracketings of elements in V . This equation need not hold if $x_0 \in \mathcal{F}(IV)$ is an arbitrary element.

The canonical strong pseudo-cylinder of a free operad is a strong cylinder when V is a sequence of cofibrant chain complexes.


We now consider strong pseudo-cylinders on the coproduct of an arbitrary DG-operad and a free one.

Given two sequences of graded modules or chain complexes, U and V , and a tree T , we define

$$(U, V)(T) \cong \bigotimes_{u \text{ odd}} U(\tilde{u}) \otimes \bigotimes_{v \text{ even}} V(\tilde{v}).$$

The first (resp. second) tensor product is indexed by the inner vertices of odd (resp. even) level, and the order of tensor factors should be the path order (in the right hand side of the formula we have separated odd and even inner vertices for lack of a better notation, the isomorphism is defined by the symmetry constraint). In (1.7), v_1 has level 1, v_2 and v_5 have level 2, and v_3 and v_4 have level 3. As above, a tensor in $(U, V)(T)$ is usually denoted by labeling each inner vertex u of odd level with an element in $U(\tilde{u})$ and each inner vertex v of even level with

an element in $V(\tilde{v})$, e.g.




Given an arbitrary operad \mathcal{O} and a sequence of graded modules or chain complexes V , the sequence underlying the coproduct $\mathcal{O} \amalg \mathcal{F}(V)$ in the category of graded or DG-operads, see [Mur11, §5] or [Mur14, Remark 3.9], is

$$(1.12) \quad (\mathcal{O} \amalg \mathcal{F}(V))(n) = \bigoplus_T (\mathcal{O}, V)(T).$$


This direct sum is indexed by the trees with n leaves, all of them in even level. In (1.7), the three topmost leaves have level 4, and the two other leaves have level 3, so this not an indexing tree here, but the following similar example is,



Using (labeled) trees, structure maps are easily defined as follows. The inclusion of the first factor $\mathcal{O} \rightarrow \mathcal{O} \amalg \mathcal{F}(V)$ is the inclusion of the direct summands indexed by corollas. The canonical map $V \rightarrow \mathcal{O} \amalg \mathcal{F}(V)$ sends an element $y \in V(n)$ to the following labeled tree,



Composition \circ_i is given by grafting into the i^{th} leaf (taking into account the path order and the Koszul sign rule) and, if u (resp. v) is the bottom (resp. top) vertex of the inner edge created by grafting and v is in the j^{th} edge adjacent to u from above, then contracting this inner edge and labelling the vertex obtained by merging u and v with the element in \mathcal{O} obtained by applying \circ_j to the labels of u and v (this also involves a sign, according to the standing conventions, which is -1 up to the product of the degree of the label of v by the sum of the degrees of the labels of the vertices strictly between u and v), e.g.



Definition 1.14. Given a DG-operad \mathcal{O} equipped with a strong pseudo-cylinder

$$\mathcal{O} \begin{array}{c} \xrightarrow{i_0} \\ \xleftarrow{i_1} \end{array} \mathcal{P} \xrightarrow{p} \mathcal{O} \quad \begin{array}{c} \xrightarrow{h_{\mathcal{O}}} \\ \xrightarrow{p} \end{array}$$

and a sequence of chain complexes V , the *canonical induced strong pseudo-cylinder* of the coproduct $\mathcal{O} \amalg \mathcal{F}(V)$

$$\mathcal{O} \amalg \mathcal{F}(V) \begin{array}{c} \xrightarrow{i_0=i_0 \amalg \mathcal{F}(i_0)} \\ \xleftarrow{i_1=i_1 \amalg \mathcal{F}(i_1)} \end{array} \mathcal{P} \amalg \mathcal{F}(IV) \begin{array}{c} \xrightarrow{p=p \amalg \mathcal{F}(p)} \\ \xrightarrow{h_{\mathcal{O},V}} \end{array} \mathcal{O} \amalg \mathcal{F}(V)$$

is defined on each $(\mathcal{O}, V)(T)$ as the tensor product of the strong pseudo-cylinder of \mathcal{O} and the canonical strong pseudo-cylinder of the sequence of chain complexes V .

If \mathcal{O} is the initial DG-operad and we take the trivial strong pseudo-cylinder on it (Definition 1.25), with $\mathcal{P} = \mathcal{O}$, $i_0 = i_1 = p = 1_{\mathcal{O}}$, and $h_{\mathcal{O}} = 0$, then we recover the canonical strong pseudo-cylinder of $\mathcal{F}(V)$.

Remark 1.15. Computing $h_{\mathcal{O},V}$ on labeled trees is as easy as with h_V above, e.g.

$$h_{\mathcal{O},V} \left(\begin{array}{c} \begin{array}{c} x_3 \\ x_4 \\ y_2 \\ x_1 \end{array} \end{array} \right) = \begin{array}{c} \begin{array}{c} x_3 \\ x_4 \\ y_2 \\ h_{\mathcal{O}}(x_1) \end{array} \end{array} + (-1)^{|x_1|} \begin{array}{c} \begin{array}{c} x_3 \\ x_4 \\ h_I(y_2) \\ i_0 p(x_1) \end{array} \end{array}$$

and a degree -1 map $\partial: Y \rightarrow Y$, called perturbation, such that $\partial^2 + d_Y \partial + \partial d_Y = 0$ and the infinite sum $\Sigma_\infty = \sum_{n \geq 0} (\partial h)^n \partial$ is well defined, i.e. almost all summands vanish when evaluated at a given $y \in Y$, then there is a new SDR

$$(X, d_X + p\Sigma_\infty i) \xrightleftharpoons[p+p\Sigma_\infty h]{i+h\Sigma_\infty i} (Y, d_Y + \partial) \curvearrowright h+h\Sigma_\infty h .$$

Remark 1.18. The vanishing condition is fulfilled if Y is equipped with an exhaustive increasing filtration

$$0 = F_{-1}Y \subset F_0Y \subset \dots \subset F_nY \subset F_{n+1}Y \subset \dots \subset Y, \quad Y = \bigcup_{n \geq 0} F_nY,$$

such that for $n \geq 0$

$$\partial(F_nY) \subset F_{n-1}Y, \quad h(F_nY) \subset F_nY.$$

This implies that, if $y \in F_nY$, then $\Sigma_\infty(y) = \sum_{j=0}^{n-1} (\partial h)^j \partial(y)$. The maps i and p often preserve the filtration, like h .

Remark 1.19. The new chain homotopy $h' = h + h\Sigma_\infty h$ satisfies the following equation,

$$h' = h + h' \partial h.$$

Remark 1.20. The differential of the twisted coproduct $\mathcal{O} \amalg_\partial \mathcal{F}(V)$ is obtained by perturbing the differential of the honest coproduct $\mathcal{O} \amalg \mathcal{F}(V)$. The perturbation $d_\partial - d_{\mathcal{O} \amalg \mathcal{F}(V)}$, that we also call ∂ , is the only degree -1 self-map of $\mathcal{O} \amalg \mathcal{F}(V)$ satisfying $\partial(\mathcal{O}) = 0$, the operadic Leibniz rule, and restricting to the original $\partial: V \rightarrow \mathcal{O}$ on V . Applying ∂ to a labeled tree with m inner vertices of even level, we obtain a sum of labeled trees with the same shape (with signs coming from the path order and the Koszul sign rule), one for each $1 \leq i \leq m$, where all labels are the same except for label of the i^{th} inner vertex of even level, which changes from y to $\partial(y)$. These labeled trees are to be regarded as iterated compositions in $\mathcal{O} \amalg \mathcal{F}(V)$, since there may be adjacent labels in \mathcal{O} , e.g.

$$\begin{aligned} \partial \left(\begin{array}{c} x_3 \\ | \\ y_2 \bullet \\ | \\ \bullet \\ / \quad | \quad \backslash \\ x_1 \end{array} \right) &= (-1)^{|x_1|} \begin{array}{c} x_3 \\ | \\ \partial(y_2) \bullet \\ | \\ \bullet \\ / \quad | \quad \backslash \\ x_1 \end{array} + (-1)^{|x_1|+|y_2|+|x_3|} \begin{array}{c} x_3 \\ | \\ y_2 \bullet \\ | \\ \bullet \\ / \quad | \quad \backslash \\ x_1 \end{array} \\ &= (-1)^{|x_1|} \begin{array}{c} y_4 \\ | \\ \bullet \\ / \quad | \quad \backslash \\ x_1 \circ_1 (\partial(y_2) \circ_1 x_3) \end{array} + (-1)^{|x_1|+|y_2|+|x_3|+(y_4-1)(|y_2|+|x_3|)} \begin{array}{c} x_3 \\ | \\ y_2 \bullet \\ | \\ \bullet \\ / \quad | \quad \backslash \\ x_1 \circ_4 \partial(y_4) \end{array} \end{aligned}$$

Theorem 1.21. *Let $\mathcal{O} \amalg_{\partial} \mathcal{F}(V)$ be a twisted coproduct as in Definition 1.16. Assume we have chosen a strong pseudo-cylinder for \mathcal{O} as in Definition 1.6. Then there is a strong pseudo-cylinder for $\mathcal{O} \amalg_{\partial} \mathcal{F}(V)$, that we call canonical induced strong pseudo-cylinder,*

$$\mathcal{O} \amalg_{\partial} \mathcal{F}(V) \begin{array}{c} \xrightarrow{i_0=i_0 \amalg \mathcal{F}(i_0)} \\ \xrightarrow{i_1=i_1 \amalg \mathcal{F}(i_1)} \end{array} \mathcal{P} \amalg_{\partial_I} \mathcal{F}(IV) \xrightarrow{p=p \amalg \mathcal{F}(p)} \mathcal{O} \amalg_{\partial} \mathcal{F}(V)$$

$\begin{array}{c} h_{\partial} \\ \curvearrowright \end{array}$

such that the degree -1 map $\partial_I: V \oplus \Sigma V \oplus V \rightarrow \mathcal{P}$ is

$$\partial_I = (i_0 \partial \quad -h_{\mathcal{O}} i_1 \partial \quad i_1 \partial)$$

and

$$h_{\partial} = \sum_{n \geq 0} (h_{\mathcal{O}, V} \partial_I)^n h_{\mathcal{O}, V},$$

where ∂_I denotes here the extension to $\mathcal{P} \amalg \mathcal{F}(IV)$ in Remark 1.20.

Proof. We first check the necessary equation to define the twisted coproduct $\mathcal{P} \amalg_{\partial_I} \mathcal{F}(IV)$,

$$\begin{aligned} \partial_I d_{IV} + d_{\mathcal{P}} \partial_I &= (i_0 \partial \quad -h_{\mathcal{O}} i_1 \partial \quad i_1 \partial) \begin{pmatrix} d_V & 1 & 0 \\ 0 & -d_V & 0 \\ 0 & -1 & d_V \end{pmatrix} + d_{\mathcal{P}} (i_0 \partial \quad -h_{\mathcal{O}} i_1 \partial \quad i_1 \partial) \\ &= (i_0 \partial d_V \quad i_0 \partial + h_{\mathcal{O}} i_1 \partial d_V - i_1 \partial \quad i_1 \partial d_V) + (d_{\mathcal{P}} i_0 \partial \quad -d_{\mathcal{P}} h_{\mathcal{O}} i_1 \partial \quad d_{\mathcal{P}} i_1 \partial) \\ &= (-i_0 d_{\mathcal{O}} \partial \quad i_0 \partial - h_{\mathcal{O}} i_1 d_{\mathcal{O}} \partial - i_1 \partial \quad -i_1 d_{\mathcal{O}} \partial) + (i_0 d_{\mathcal{O}} \partial \quad -d_{\mathcal{P}} h_{\mathcal{O}} i_1 \partial \quad i_1 d_{\mathcal{O}} \partial) \\ &= (0 \quad i_0 p i_1 \partial - i_1 \partial - h_{\mathcal{O}} d_{\mathcal{P}} i_1 \partial - d_{\mathcal{P}} h_{\mathcal{O}} i_1 \partial \quad 0) \\ &= (0 \quad (i_0 p - 1 - h_{\mathcal{O}} d_{\mathcal{P}} - d_{\mathcal{P}} h_{\mathcal{O}}) i_1 \partial \quad 0) \\ &= (0 \quad 0 \quad 0). \end{aligned}$$

Here we use the equations $\partial d_V + d_{\mathcal{O}} \partial = 0$, $i_0 p - 1_{\mathcal{P}} = d_{\mathcal{P}} h_{\mathcal{O}} + h_{\mathcal{O}} d_{\mathcal{P}}$, and $p i_1 = 1_{\mathcal{O}}$, and the fact that $i_0, i_1: \mathcal{O} \rightarrow \mathcal{P}$ are DG-maps. The perturbation equation in Lemma 1.17 is a consequence of this. We want to apply this lemma to the sequence of SDRs

$$\mathcal{O} \amalg \mathcal{F}(V) \begin{array}{c} \xleftarrow{i_0} \\ \xrightarrow{p} \end{array} \mathcal{P} \amalg \mathcal{F}(IV) \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} h_{\mathcal{O}, V}$$

The vanishing condition follows from the criterion in Remark 1.18. The sequence of subcomplexes $F_n(\mathcal{P} \amalg \mathcal{F}(IV))$ is aritywise the direct subsum indexed by the trees T with $\leq n$ vertices of even level. The chain homotopy $h_{\mathcal{O}, V}$ preserves filtration levels since it restricts to each direct summand. It is easy to see that the extension of ∂_I to $\mathcal{P} \amalg \mathcal{F}(IV)$ strictly decreases filtration levels by using the Leibniz rule and the facts that $\partial_I(\mathcal{P}) = 0$ and $\partial_I(IV) \subset \mathcal{P}$.

The perturbation lemma applies, but we still have to check that the maps i_0 and p do not change and that the induced perturbation on $\mathcal{O} \amalg \mathcal{F}(V)$ is ∂ , i.e.

$$h_{\mathcal{O}, V} \Sigma_{\infty} i_0 = 0, \quad p \Sigma_{\infty} h_{\mathcal{O}, V} = 0, \quad p \Sigma_{\infty} i_0 = \partial.$$

Since $\partial_I i_0 = i_0 \partial: \mathcal{O} \amalg \mathcal{F}(V) \rightarrow \mathcal{P} \amalg \mathcal{F}(IV)$, $h_{\mathcal{O},V} \partial_I i_0 = h_{\mathcal{O},V} i_0 \partial = 0 \partial = 0$. This clearly proves the first equation. Also the third one, since it implies that $p \Sigma_\infty i_0 = p \partial_I i_0 = p i_0 \partial = \partial$.

The middle equation follows from $p \partial_I h_{\mathcal{O},V} = 0$, that we now check. Since $ph_{\mathcal{O}} = 0: \mathcal{P} \rightarrow \mathcal{O}$ and $ph_I = 0: IV \rightarrow V$, p applied to a labeled tree containing a label of the form $h_{\mathcal{O}}(x)$, $x \in \mathcal{P}$, or $h_I(y)$, $y \in IV$, yields 0. If we apply $\partial_I h_{\mathcal{O},V}$ to a labeled tree in $\mathcal{P} \amalg \mathcal{F}(IV)$, we obtain a linear combination of labeled trees, many of them containing labels of the previous form, except for those where the label of an inner vertex of even degree has changed from $y \in IV$ to $\partial_I h_I(y)$, but

$$p \partial_I h_I = p \begin{pmatrix} i_0 \partial & -h_{\mathcal{O}} i_1 \partial & i_1 \partial \\ 0 & 0 & \sigma \\ 0 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -ph_{\mathcal{O}} i_1 \partial \sigma \\ 0 & 0 & -0 i_1 \partial \sigma \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Therefore, $p \partial_I h_{\mathcal{O},V} = 0$. \square

For $\partial = 0$, we recover the canonical induced strong pseudo-cylinder of the coproduct. The differential in the canonical strong pseudo-cylinder of an element of the form $\sigma(x)$, $x \in V$, is

$$(1.22) \quad d\sigma(x) = i_0(x) - i_1(x) - h_{\mathcal{O}} i_1 \partial(x).$$

In general, the computation of h_∂ is somewhat involved. Let us just check that it extends the chain homotopies of the canonical strong pseudo-cylinder of V and the chosen strong pseudo-cylinder of \mathcal{O} .

Lemma 1.23. *If $\mathcal{P} \amalg_{\partial_I} \mathcal{F}(IV)$ is the canonical induced strong pseudo-cylinder of a twisted coproduct, $x \in \mathcal{P}$, and $y \in IV$, then*

$$h_\partial(x) = h_{\mathcal{O}}(x), \quad h_\partial(y) = h_I(y).$$

Proof. We use Remark 1.18 and the filtration in the proof of Theorem 1.21. As an element of $\mathcal{P} \amalg_{\partial_I} \mathcal{F}(IV)$, $x \in \mathcal{P}$ has filtration degree 0, hence

$$h_\partial(x) = h_{\mathcal{O},V}(x) = h_{\mathcal{O}}(x).$$

Moreover, $y \in IV$ has filtration degree 1, so

$$\begin{aligned} h_\partial(y) &= h_{\mathcal{O},V}(y) + h_{\mathcal{O}} \partial_I h_{\mathcal{O},V}(y) \\ &= h_I(y) + h_{\mathcal{O}} \partial_I h_I(y) \\ &= h_I(y) + h_{\mathcal{O}} \begin{pmatrix} i_0 \partial & -h_{\mathcal{O}} i_1 \partial & i_1 \partial \\ 0 & 0 & \sigma \\ 0 & 0 & 0 \end{pmatrix} (y) \\ &= h_I(y) + \begin{pmatrix} 0 & 0 & -h_{\mathcal{O}}^2 i_1 \partial \sigma \end{pmatrix} (y) \\ &= h_I(y). \end{aligned}$$

Here we use that $h_{\mathcal{O}}^2 = 0$. \square

We finally consider strong pseudo-cylinders on DG-operads constructed as iterated twisted coproducts.

Definition 1.24. A DG-operad \mathcal{O} is *relatively pseudo-cellular* if it is equipped with an increasing filtration $\{\mathcal{O}_\beta\}_{\beta \leq \alpha}$ indexed by an ordinal α , that we call *length*, which is *exhaustive*, i.e. $\mathcal{O} = \mathcal{O}_\alpha$, *continuous*, i.e. if $\beta \leq \alpha$ is a limit ordinal then $\mathcal{O}_\beta = \operatorname{colim}_{\gamma < \beta} \mathcal{O}_\gamma$ (here colim can be replaced with \cup), and such that if $\beta + 1 \leq \alpha$ then $\mathcal{O}_{\beta+1}$ is a twisted coproduct of the following form,

$$\mathcal{O}_{\beta+1} = \mathcal{O}_\beta \amalg_{\partial_\beta} \mathcal{F}(V_\beta).$$

We say that \mathcal{O} is (*absolutely*) *pseudo-cellular* if in addition \mathcal{O}_0 is the initial DG-operad.

If the complexes of the sequences V_β are cofibrant for all $\beta < \alpha$, then the inclusion $\mathcal{O}_0 \rightarrow \mathcal{O}$ is a cofibration. In particular, \mathcal{O} is cofibrant in the absolute case. If the V_β are sequences of free graded modules with trivial differential, then $\mathcal{O}_0 \rightarrow \mathcal{O}$ is a relative cell complex with respect to the standard set of generating cofibrations in the model category of DG-operads, see [Hov99, Theorem 2.3.11] and [Mur11, proof of Theorem 1.1].

Strong pseudo-cylinders are closed under filtered colimits, for both chain complexes and DG-operads (filtered colimits of DG-operads are computed in the underlying sequences of chain complexes).

Definition 1.25. The *canonical strong pseudo-cylinder* of a relatively pseudo-cellular DG-operad \mathcal{O}

$$\mathcal{O} \begin{array}{c} \xrightarrow{i_0} \\ \xrightarrow{i_1} \end{array} I\mathcal{O} \begin{array}{c} \xrightarrow{h_{\mathcal{O}}} \\ \xrightarrow{p} \end{array} \mathcal{O}$$

is defined by induction on the length α in the following way. If $\alpha = 0$, we take the *trivial strong pseudo-cylinder*,

$$\mathcal{O} \begin{array}{c} \xrightarrow{1} \\ \xrightarrow{1} \end{array} \mathcal{O} \begin{array}{c} \xrightarrow{0} \\ \xrightarrow{1} \end{array} \mathcal{O}.$$

If α is a limit ordinal, we define the canonical strong pseudo-cylinder of \mathcal{O} as the colimit of the canonical strong pseudo-cylinders of \mathcal{O}_β , $\beta < \alpha$. If $\alpha = \beta + 1$, the canonical strong pseudo-cylinder of $\mathcal{O} = \mathcal{O}_\beta \amalg_{\partial_\beta} \mathcal{F}(V_\beta)$ is defined by applying Theorem 1.21 to the canonical strong pseudo-cylinder of \mathcal{O}_β ,

$$\mathcal{O}_\beta \amalg_{\partial_\beta} \mathcal{F}(V_\beta) \begin{array}{c} \xrightarrow{i_0=i_0 \amalg \mathcal{F}(i_0)} \\ \xrightarrow{i_1=i_1 \amalg \mathcal{F}(i_1)} \end{array} I\mathcal{O}_\beta \amalg_{\partial_{\beta,I}} \mathcal{F}(IV_\beta) \begin{array}{c} \xrightarrow{h_{\mathcal{O}}=h_{\partial_\beta}} \\ \xrightarrow{p=p \amalg \mathcal{F}(p)} \end{array} \mathcal{O}_\beta \amalg_{\partial_\beta} \mathcal{F}(V_\beta),$$

in particular, $I\mathcal{O} = I\mathcal{O}_\beta \amalg_{\partial_{\beta,I}} \mathcal{F}(IV_\beta)$.

The DG-operad $I\mathcal{O}$ is relatively pseudo-cellular of the same length as \mathcal{O} and $(I\mathcal{O})_0 = \mathcal{O}_0$. In particular, if \mathcal{O} is absolutely pseudo-cellular, then so is $I\mathcal{O}$.

If the complexes of the sequences V_β , $\beta < \alpha$, are cofibrant, the canonical strong pseudo-cylinder of \mathcal{O} is a relative cylinder in the model theoretic sense, i.e. $(i_0, i_1): \mathcal{O} \cup_{\mathcal{O}_0} \mathcal{O} \rightarrow I\mathcal{O}$ is a cofibration. In particular, if \mathcal{O} is absolutely pseudo-cellular, its canonical strong pseudo-cylinder is a strong cylinder.

As a graded operad, any relatively pseudo-cellular DG-operad of length α is $\mathcal{O} = \mathcal{O}_0 \amalg \mathcal{F}(V)$, $V = \bigoplus_{\beta < \alpha} V_\beta$, and the differential on the free part is determined by the equations $d(x) = d_{V_\beta}(x) + \partial_\beta(x)$, $x \in V_\beta$. The canonical strong pseudo-cylinder, as a graded operad, is $I\mathcal{O} = \mathcal{O}_0 \amalg \mathcal{F}(IV)$. On elements of the form $i_0(x)$ and $i_1(x)$, $x \in V_\beta$, the differential is simply given by

$$di_j(x) = i_j d(x) = i_j d_{V_\beta}(x) + i_j \partial_\beta(x), \quad j = 0, 1,$$

since $i_0, i_1: \mathcal{O} \rightarrow I\mathcal{O}$ are DG-operad maps. On elements of the form σx , $x \in V_\beta$, the differential depends on the involved inductive definition of the chain homotopy

$$(1.26) \quad d\sigma(x) = i_0(x) - i_1(x) - h_{\mathcal{O}_\beta} i_1 \partial_\beta(x),$$

see (1.22). The chain homotopy is easy on \mathcal{O}_0 and on generators of the free part, as a corollary of Lemma 1.23.

Corollary 1.27. *If \mathcal{O} is a relatively pseudo-cellular DG-operad of length α , $x \in \mathcal{O}_0$, and $y \in IV_\beta$ for some $\beta < \alpha$, then*

$$h_{\mathcal{O}}(x) = 0, \quad h_{\mathcal{O}}(y) = h_I(y).$$


In particular, $h_{\mathcal{O}}(\text{id}) = 0$.

The underlying graded operad of an absolutely pseudo-cellular DG-operad \mathcal{O} is free, $\mathcal{O} = \mathcal{F}(V)$, $V = \bigoplus_{\beta < \alpha} V_\beta$, since \mathcal{O}_0 is initial and hence it disappears in coproducts. In order to prove a useful vanishing condition for the chain homotopy, we introduce the following set of linear generators of $I\mathcal{O} = \mathcal{F}(V)$ (as a sequence of graded modules).


Definition 1.28. Let \mathcal{O} be an absolutely pseudo-cellular DG-operad of length α as above. A *standard labeled tree* in $I\mathcal{O}$ is a labeled tree such that each label is of the form $i_0(x)$, $i_1(x)$ or $\sigma(x)$, $x \in V_\beta$, $\beta < \alpha$.

Lemma 1.29. *Let \mathcal{O} be an absolutely pseudo-cellular DG-operad and $t \in I\mathcal{O}$ a standard labeled tree satisfying one of the two following conditions:*


- (1) *The bottommost label is $\sigma(x)$.*



(2) The bottommost label is $i_0(x)$



and t does not contain any forbidden edge, i.e. an inner edge with bottom label $i_0(y)$ and top label $i_1(z)$.



Then, $h_{\mathcal{O}}(t) = 0$.

Proof. By induction on the length α . If $\alpha = 0$, then the chain homotopy is $h_{\mathcal{O}} = 0$, so there is nothing to check. If α is a limit ordinal, then $h_{\mathcal{O}}(t) = h_{\mathcal{O}_\beta}(t)$ for some $\beta < \alpha$, and $h_{\mathcal{O}_\beta}(t) = 0$ by induction hypothesis.

Suppose $\alpha = \beta + 1$. We prove, by induction on the number of inner vertices, that $h_{\mathcal{O}_\beta, V_\beta}(t) = 0$. This clearly suffices.

Suppose t satisfies (1). If $x \in V_\beta$, we can write $t = \sigma(x)(x_1, \dots, x_n)$. The formula in the second paragraph of Remark 1.15 yields

$$h_{\mathcal{O}_\beta, V_\beta}(\sigma(x)(x_1, \dots, x_n)) = h_I(\sigma(x))(x_1, \dots, x_n) \pm \text{terms of the form } i_0 p(\sigma(x))(\dots, h_{\mathcal{O}_\beta, V_\beta}(x_i), \dots).$$

All summands vanish since $h_I \sigma = 0$ and $p\sigma = 0$.

Otherwise, we can write $t = x_0(x_1, \dots, x_n)$. Here, $x_0 \in IO_\beta$ is a standard labeled tree with the same bottommost label as t , $\sigma(x)$. Hence, $h_{\mathcal{O}_\beta}(x_0) = 0$ by the first induction hypothesis, and $p(x_0) = 0$ since $p\sigma = 0$. The formula in the third paragraph of Remark 1.15 yields

$$(1.30) \quad h_{\mathcal{O}_\beta, V_\beta}(x_0(x_1, \dots, x_n)) = h_{\mathcal{O}_\beta}(x_0)(x_1, \dots, x_n) \pm \text{terms of the form } i_0 p(x_0)(\dots, h_{\mathcal{O}_\beta, V_\beta}(x_i), \dots),$$

which therefore vanishes.

Suppose now that t satisfies (2). If $x \in V_\beta$, we can write the standard labeled tree as $i_0(x)(x_1, \dots, x_n)$, where the elements x_1, \dots, x_n are standard labeled trees with less inner vertices than t . We will check that $h_{\mathcal{O}_\beta, V_\beta}(x_i) = 0$, $1 \leq i \leq n$.

Hence, the formula in the second paragraph of Remark 1.15 yields

$$h_{\mathcal{O}_\beta, V_\beta}(i_0(x)(x_1, \dots, x_n)) = h_I i_0(x)(x_1, \dots, x_n) \\ \pm \text{terms of the form } i_0 p i_0(x)(\dots, h_{\mathcal{O}_\beta, V_\beta}(x_i), \dots),$$

which vanishes since $h_I i_0 = 0$.

If the standard labeled tree x_i has no inner vertices, then $x_i = \text{id}$ and $h_{\mathcal{O}_\beta, V_\beta}(\text{id}) = h_0(\text{id}) = 0$. If it has inner vertices, let us look at the bottommost label. It cannot be $i_1(y)$, since it is adjacent to $i_0(x)$ in t , and we are assuming that t does not contain forbidden edges. Hence it is $\sigma(y)$ or $i_0(y)$. Not containing a forbidden edge is a property inherited by subtrees. Hence, x_i does not contain forbidden edges. Therefore, x_i satisfies (1) or (2) and has less inner vertices than t , so $h_{\mathcal{O}_\beta, V_\beta}(x_i) = 0$ by the second induction hypothesis.

If $x \in V_\gamma$ for some $\gamma < \beta$, then we can write the standard labeled tree as $x_0(x_1, \dots, x_n)$, where $x_0 \in I\mathcal{O}_\beta$ is a standard labeled tree and, for $1 \leq i \leq n$, either $x_i = \text{id}$ or $x_i = y_{i,0}(y_{i,1}, \dots, y_{i,p_i})$ is a standard labeled tree with less inner vertices than t where $y_{i,0}$ is $i_0(y'_i)$, $\sigma(y'_i)$ or $i_1(y'_i)$ for some $y'_i \in V_\beta$. Formula (1.30) also applies in this case. Hence, it suffices to prove that $h_{\mathcal{O}_\beta}(x_0) = 0$ and either $p(x_0) = 0$ or $h_{\mathcal{O}_\beta, V_\beta}(x_i) = 0$ for all $1 \leq i \leq n$.

The bottommost label of x_0 is the same as in t , $i_0(x)$, and x_0 cannot contain a forbidden edge, since it is a subtree of t . Hence $h_{\mathcal{O}_\beta}(x_0) = 0$ by the first induction hypothesis. For $1 \leq i \leq n$, if $x_i = \text{id}$ then we know that $h_{\mathcal{O}_\beta, V_\beta}(x_i) = 0$. Otherwise, let us argue with the possible values of $y_{i,0}$. Being a subtree of t , x_i does not contain any forbidden edge. Hence, if $y_{i,0}$ is $i_0(y'_i)$ or $\sigma(y'_i)$, $h_{\mathcal{O}_\beta, V_\beta}(x_i) = 0$ by the second induction hypothesis, since x_i has less inner vertices than t . The bottommost vertex of x_i is adjacent to a vertex in x_0 . Therefore, if $y_{i,0} = i_1(y'_i)$ then the adjacent vertex in x_0 is $\sigma(z)$, otherwise t would contain a forbidden edge. In this case $p(x_0) = 0$ since $p\sigma = 0$. \square

2. EXAMPLES

We start this section on examples by illustrating how our canonical strong cylinder construction works on the most widely used cellular DG-operad.

Definition 2.1. The *A-infinity DG-operad* \mathcal{A}_∞ is freely generated as a graded operad by

$$\mu_n \in \mathcal{A}_\infty(n)_{n-2}, \quad n \geq 2,$$

with differential defined by

$$d(\mu_n) = \sum_{\substack{p+q=n+1 \\ 1 \leq i \leq p}} (-1)^{p-i+q(i-1)} \mu_p \circ_i \mu_q.$$

Here we use the sign conventions in [LH03], but we should point out that Lefèvre-Hasegawa uses cohomological grading and, modulo this, an *A-infinity algebra* $(X, m_1, m_2, \dots, m_n, \dots)$ in his sense is an \mathcal{A}_∞ -algebra structure on the

chain complex $(X, -m_1)$. This DG-operad is cellular. The following result computes its canonical strong cylinder.

Theorem 2.2. *The canonical strong cylinder of the A -infinity DG-operad is the DG-operad $I\mathcal{A}_\infty$, freely generated as a graded operad by*

$$i_0\mu_n, i_1\mu_n \in I\mathcal{A}_\infty(n)_{n-2}, \quad \sigma\mu_n \in I\mathcal{A}_\infty(n)_{n-1}, \quad n \geq 2,$$

with differential determined by the fact that $i_0, i_1: \mathcal{A}_\infty \rightarrow I\mathcal{A}_\infty$ are DG-operad maps and by the following formula, $n \geq 2$,

$$\begin{aligned} d(\sigma\mu_n) = & i_0\mu_n - i_1\mu_n - \sum_{\substack{p+q=n+1 \\ 1 \leq j \leq p}} (-1)^{p-j+q(j-1)} \sigma\mu_p \circ_j i_1\mu_q \\ & + \sum_{\substack{1 \leq s \leq r \\ j_0+t_1+j_1+\dots+t_s+j_s=n}} (-1)^{\sum_{k=1}^s (t_k-1)(j_0+\sum_{l=1}^{k-1} (t_l+j_l))} i_0\mu_r(\cdot^{j_0}, \sigma\mu_{t_1}, \cdot^{j_1}, \sigma\mu_{t_2}, \dots, \sigma\mu_{t_s}, \cdot^{j_s}). \end{aligned}$$

This theorem follows from (1.26) and the last formula in Lemma 2.10.

Corollary 2.3. *Given a chain complex X and two maps to its endomorphism operad $\varphi, \varphi': \mathcal{A}_\infty \rightarrow \mathcal{E}(X)$, see Remark 2.5 below, which correspond with two A -infinity structures on X , $(X, \{m_n\}_{n \geq 2})$ and $(X, \{m'_n\}_{n \geq 2})$, respectively, a homotopy $H: I\mathcal{A}_\infty \rightarrow \mathcal{E}(X)$ between them, $Hi_0 = \varphi'$, $Hi_1 = \varphi$, is the same as an A -infinity morphism [LH03, Définition 1.2.1.2]*

$$\{f_n\}_{n \geq 1}: (X, \{m_n\}_{n \geq 2}) \longrightarrow (X, \{m'_n\}_{n \geq 2})$$

whose linear part is the identity in X , $f_1 = 1_X$.

Proof. The correspondence is simply given by $m'_n = \varphi'(\mu_n) = H(i_0\mu_n)$, $m_n = \varphi(\mu_n) = H(i_1\mu_n)$, and $f_n = H(\sigma\mu_n)$, $n \geq 2$. \square

In order to simplify computations, we use the formalism of operadic suspensions and brace algebras.

Definition 2.4. Given a DG-operad \mathcal{O} , its *operadic suspension* is the DG-operad $\Lambda\mathcal{O}$ such that $\Lambda\mathcal{O}(n) = \mathcal{O}(n)$ as plain modules, $n \geq 0$, with the following new grading

$$\|x\| = |x| + 1 - \text{arity of } x.$$

The differential is the same as in \mathcal{O} . Compositions in $\Lambda\mathcal{O}$, that we here denote by \bullet_i in order to avoid confusion, are defined as follows, $x \in \mathcal{O}(p)$, $y \in \mathcal{O}(q)$,

$$x \bullet_i y = (-1)^{\|y\|(p-i)+|y|(i-1)} x \circ_i y,$$

and the identity is the same $\text{id}_{\Lambda\mathcal{O}} = \text{id}_{\mathcal{O}}$.

Remark 2.5. The functor Λ is an automorphism of the category of DG-operads. It preserves free operads, $\Lambda\mathcal{F}(V) = \mathcal{F}(\Lambda V)$, where ΛV is defined as above, (twisted) coproducts, (relatively) pseudo-cellular DG-operads, (canonical) strong

(pseudo-)cylinders, etc. It also preserves the homotopical structure (fibrations, cofibrations, and weak equivalences).

Recall that, given chain complexes X and Y , the inner $\text{Hom}(X, Y)$ is the chain complex consisting of the modules $\text{Hom}(X, Y)_n$ of degree n maps $f: X \rightarrow Y$ with differential $d(f) = d_Y f - (-1)^{|f|} f d_X$, in particular chain maps $X \rightarrow Y$ are 0-cycles in $\text{Hom}(X, Y)$. The endomorphism operad of a chain complex X is

$$\mathcal{E}(X) = \{\text{Hom}(X^{\otimes n}, X)\}_{n \geq 0},$$

the operation \circ_i is composition at the i^{th} slot, the operadic identity is the identity map $\text{id}_{\mathcal{E}(X)} = 1_X$, and an \mathcal{O} -algebra structure on X is an operad map $\mathcal{O} \rightarrow \mathcal{E}(X)$. There is an isomorphism of DG-operads

$$\Lambda \mathcal{E}(X) \cong \mathcal{E}(\Sigma X),$$

defined by mapping $f: X^{\otimes n} \rightarrow X$ to $(-1)^{|f|} \sigma f (\sigma^{-1})^{\otimes n}: (\Sigma X)^{\otimes n} \rightarrow \Sigma X$, $n \geq 0$. Therefore, an \mathcal{O} -algebra structure on X is the same as a $\Lambda \mathcal{O}$ -algebra structure on ΣX .

The *Hadamard product* of two DG-operads $\mathcal{O} \otimes_{\mathbb{H}} \mathcal{P}$ is the DG-operad with $(\mathcal{O} \otimes_{\mathbb{H}} \mathcal{P})(n) = \mathcal{O}(n) \otimes \mathcal{P}(n)$, $n \geq 0$, compositions

$$(x_1 \otimes x_2) \circ_i (y_1 \otimes y_2) = (-1)^{|x_2||y_1|} (x_1 \circ_i y_1) \otimes (x_2 \circ_i y_2),$$

and identity $\text{id}_{\mathcal{O} \otimes_{\mathbb{H}} \mathcal{P}} = \text{id}_{\mathcal{O}} \otimes \text{id}_{\mathcal{P}}$. The operadic suspension $\Lambda \mathcal{O}$ can be naturally identified with $\mathcal{O} \otimes_{\mathbb{H}} \mathcal{E}(\Sigma \mathbb{k})$. The natural isomorphism

$$\Lambda \mathcal{O} \cong \mathcal{O} \otimes_{\mathbb{H}} \mathcal{E}(\Sigma \mathbb{k})$$

maps $x \in \mathcal{O}(n)$ to $x \otimes (\sigma \varphi_n (\sigma^{-1})^{\otimes n})$, where $\varphi_n: \mathbb{k}^{\otimes n} \rightarrow \mathbb{k}$ is defined by $\varphi_n(1 \otimes \cdots \otimes 1) = 1$.

Definition 2.6. A *graded* or *DG-brace algebra* is a graded module or chain complex B equipped with maps, called *braces*, $n \geq 1$,

$$\begin{aligned} B^{\otimes(n+1)} &\longrightarrow B, \\ x_0 \otimes x_1 \otimes \cdots \otimes x_n &\mapsto x_0 \{x_1, \dots, x_n\}, \end{aligned}$$

satisfying

$$\begin{aligned} x \{y_1, \dots, y_p\} \{z_1, \dots, z_q\} &= \sum_{0 \leq i_1 \leq j_1 \leq \cdots \leq i_p \leq j_p \leq q} (-1)^\epsilon x \{z_1, \dots, z_{i_1}, y_1 \{z_{i_1+1}, \dots, z_{j_1}\}, \dots \\ &\quad \dots, y_p \{z_{i_p+1}, \dots, z_{j_p}\}, z_{j_p+1}, \dots, z_q\}. \end{aligned}$$

The sign $(-1)^\epsilon$ is simply determined by the Koszul sign rule,

$$\epsilon = \sum_{k=1}^p \sum_{l=1}^{i_k} |y_k| |z_l|.$$

In the DG-case, the fact that braces are chain maps is equivalent to the following *brace Leibniz rule*:

$$d(x_0\{x_1, \dots, x_n\}) = d(x_0)\{x_1, \dots, x_n\} + \sum_{i=1}^n (-1)^{\sum_{j=0}^{i-1} |x_j|} x_0\{x_1, \dots, d(x_i), \dots, x_n\}.$$

Remark 2.7. If \mathcal{O} is a graded or DG-operad, then $\bigoplus_{n \geq 0} \mathcal{O}(n)$ has a brace algebra structure defined as follows. Given $x_0, \dots, x_n \in \mathcal{O}$, with $x_0 \in \mathcal{O}(m)$,

$$x_0\{x_1, \dots, x_n\} = \sum_{i_0 + \dots + i_n = m-n} x_0(\cdot^{i_0}, x_1, \cdot^{i_1}, x_2, \dots, x_n, \cdot^{i_n}).$$

Note that $x_0\{x_1, \dots, x_n\} = 0$ if $n > m$, since the summation is empty in this case.

Remark 2.8. The DG-operad $\Lambda\mathcal{A}_\infty$ is freely generated as a graded operad by

$$\mu_n \in (\Lambda\mathcal{A}_\infty)(n)_{-1}, \quad n \geq 2,$$

with differential defined by

$$d(\mu_n) = \sum_{p+q=n+1} \mu_p\{\mu_q\}.$$

This is a cellular DG-operad with $\Lambda\mathcal{A}_0 = \Lambda\mathcal{A}_1$ the initial DG-operad and, for $n \geq 2$,

$$\Lambda\mathcal{A}_n = \Lambda\mathcal{A}_{n-1} \amalg_{\partial_{n-1}} \mathcal{F}(\mathbb{k} \cdot \mu_n), \quad \partial_{n-1}(\mu_n) = d(\mu_n).$$

Here $\mathbb{k} \cdot \mu_n$ denotes the sequence of graded modules freely generated by μ_n in arity n and degree -1 endowed with the trivial differential.

Remark 2.9. The chain homotopy h_V of the canonical strong pseudo-cylinder of a free DG-operad $\mathcal{F}(V)$ is well behaved with respect to braces. Given $x_0 \in IV$ and $x_1, \dots, x_n \in \mathcal{F}(IV)$,

$$\begin{aligned} h_V(x_0\{x_1, \dots, x_n\}) &= h_I(x_0)\{x_1, \dots, x_n\} \\ &+ \sum_{i=1}^n (-1)^{\sum_{j=0}^{i-1} |x_j|} i_0p(x_0)\{i_0p(x_1), \dots, i_0p(x_{i-1}), h_V(x_i), x_{i+1}, \dots, x_n\}. \end{aligned}$$

This follows from Remark 1.10.

The chain homotopy $h_{\mathcal{O},V}$ of the canonical induced strong pseudo-cylinder of a coproduct $\mathcal{O} \amalg \mathcal{F}(V)$, see Definition 1.14, satisfies the previous formula if $x_0 \in IV$ and $x_1, \dots, x_n \in \mathcal{P} \amalg \mathcal{F}(IV)$, replacing h_V with $h_{\mathcal{O},V}$. Also if $x_0 \in \mathcal{P}$ and each $x_i \in \mathcal{P} \amalg \mathcal{F}(IV)$, $1 \leq i \leq n$, is either $x_i = \text{id}$ or $x_i = y_{i,0}\{y_{i,1}, \dots, y_{i,p_i}\}$ with $y_{i,0} \in IV$, replacing h_V with $h_{\mathcal{O},V}$ and h_I with $h_{\mathcal{O}}$. This follows from Remark 1.15.

The following lemma contains the formulas which prove Theorem 2.2. The first technical series of formulas is auxiliary. What really matters is the last one.

Lemma 2.10. *The following equations hold in $I\Lambda\mathcal{A}_n$,*

$$\begin{aligned}
(2.11) \quad & h_{\Lambda\mathcal{A}_n}(i_0\mu_r(\dots, \sigma\mu_{t_1}, \dots, \sigma\mu_{t_{j-1}}, \dots, i_1\mu_q, \dots, \sigma\mu_{t_j}, \dots, \sigma\mu_{t_s}, \dots)) \\
& = h_{\Lambda\mathcal{A}_{n-1, \mathbb{k}, \mu_n}}(\text{the same element}) \quad \text{for } n = \max\{r, q, t_1, \dots, t_s\}, \\
& = -i_0\mu_r(\dots, \sigma\mu_{t_1}, \dots, \sigma\mu_{t_{j-1}}, \dots, \sigma\mu_q, \dots, \sigma\mu_{t_j}, \dots, \sigma\mu_{t_s}, \dots) \\
& \quad \text{if } t_1, \dots, t_{j-1} > r \geq q, \\
& \quad \text{or if } t_1, \dots, t_{j-1} > q > r \text{ and } q \leq t_j, \dots, t_s; \\
& = 0 \quad \text{otherwise;}
\end{aligned}$$

$$h_{\Lambda\mathcal{A}_n}i_1d(\mu_{n+1}) = \sum_{p+q=n+2} \sigma\mu_p\{i_1\mu_q\} - \sum_{\substack{1 \leq s \leq r \\ t_1 + \dots + t_s = n+1+s-r}} i_0\mu_r\{\sigma\mu_{t_1}, \dots, \sigma\mu_{t_s}\}.$$

Proof. We simultaneously prove all equations by induction on n . Notice that there is nothing to check for $n = 0, 1$. Take a bigger n and assume that the formulas are true for smaller values. We start with the first series of equations. For the sake of simplicity, let us denote

$$x = i_0\mu_r(\dots, \sigma\mu_{t_1}, \dots, \sigma\mu_{t_{j-1}}, \dots, i_1\mu_q, \dots, \sigma\mu_{t_j}, \dots, \sigma\mu_{t_s}, \dots).$$

If $n > r, q$, then n is reached at some of the t_i s. Hence,

$$x = i_0\mu_r(\dots, i_1\mu_q, \dots)(\dots, \sigma\mu_n, \dots, \sigma\mu_n, \dots),$$

where $i_0\mu_r(\dots, i_1\mu_q, \dots)$ has been obtained by removing all $\sigma\mu_n$ s from x . Therefore, by Remark 1.15,

$$\begin{aligned}
h_{\Lambda\mathcal{A}_{n-1, \mathbb{k}, \mu_n}}(x) & = h_{\Lambda\mathcal{A}_{n-1}}(i_0\mu_r(\dots, i_1\mu_q, \dots))(\dots, \sigma\mu_n, \dots, \sigma\mu_n, \dots) \\
& \quad + \sum i_0p(i_0\mu_r(\dots, i_1\mu_q, \dots))(\dots, h_I\sigma\mu_n, \dots).
\end{aligned}$$

We do not index the summation since all terms vanish anyway (recall that $h_I\sigma = 0$). The maximum subscript in $i_0\mu_r(\dots, i_1\mu_q, \dots)$ is smaller than n . Moreover, $i_0\mu_r(\dots, i_1\mu_q, \dots)$ satisfies assumption (2.11) if and only if x does. Therefore, using the induction hypothesis, we obtain that

$$\begin{aligned}
h_{\Lambda\mathcal{A}_{n-1, \mathbb{k}, \mu_n}}(x) & = -i_0\mu_r(\dots, \sigma\mu_q, \dots)(\dots, \sigma\mu_n, \dots, \sigma\mu_n, \dots) \\
& = -i_0\mu_r(\dots, \sigma\mu_{t_1}, \dots, \sigma\mu_{t_{j-1}}, \dots, \sigma\mu_q, \dots, \sigma\mu_{t_j}, \dots, \sigma\mu_{t_s}, \dots) \\
& \quad \text{if (2.11) holds,} \\
& = 0 \quad \text{otherwise.}
\end{aligned}$$

If $q = n > r$ and there is some $t_i < n$, then

$$x = i_0\mu_r(\dots, \sigma\mu_{t_i}, \dots)(\dots, \sigma\mu_n, \dots, i_1\mu_n, \dots, \sigma\mu_n, \dots),$$

where $i_0\mu_r(\dots, \sigma\mu_{t_i}, \dots)$ has been obtained by removing $i_1\mu_n$ and all $\sigma\mu_n$ s from x . By Remark 1.15,

$$h_{\Lambda\mathcal{A}_{n-1, \mathbb{k}, \mu_n}}(x) = h_{\Lambda\mathcal{A}_{n-1}}(i_0\mu_r(\dots, \sigma\mu_{t_i}, \dots))(\dots, \sigma\mu_n, \dots, i_1\mu_n, \dots, \sigma\mu_n, \dots)$$

$$\begin{aligned}
& - \sum i_0 p(i_0 \mu_r(\dots, \sigma \mu_{t_i}, \dots))(\dots) \\
& = 0.
\end{aligned}$$

The first term is 0 by Lemma 1.29 (2). The summation vanishes since $p\sigma = 0$. In this case (2.11) cannot hold.

If $q = t_1 = \dots = t_s = n > r$, then by Remark 1.15,

$$\begin{aligned}
h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x) &= h_{\Lambda \mathcal{A}_{n-1}}(i_0 \mu_r)(\dots, \sigma \mu_n, \dots, i_1 \mu_n, \dots, \sigma \mu_n, \dots) \\
& - \sum i_0 p(i_0 \mu_r)(\dots, h_I \sigma \mu_n, \dots, i_1 \mu_n, \dots, \sigma \mu_n, \dots) \\
& - i_0 p(i_0 \mu_r)(\dots, i_0 p \sigma \mu_n, \dots, h_I i_1 \mu_n, \dots, \sigma \mu_n, \dots) \\
& + \sum i_0 p(i_0 \mu_r)(\dots, i_0 p \sigma \mu_n, \dots, i_0 p i_1 \mu_n, \dots, h_I \sigma \mu_n, \dots).
\end{aligned}$$

Using that $h_{\Lambda \mathcal{A}_{n-1}} i_0 = 0$, $h_I \sigma = 0$, and $p\sigma = 0$, we see that all factors vanish unless $i_1 \mu_n$ is the first element in the brackets in x , i.e. $j = 1$. In that case, there is a single non-vanishing term:

$$\begin{aligned}
h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x) &= -i_0 p i_0(\mu_r)(\dots, h_I i_1 \mu_n, \dots, \sigma \mu_n, \dots) \\
& = -i_0(\mu_r)(\dots, \sigma \mu_n, \dots, \sigma \mu_n, \dots).
\end{aligned}$$

If $r = n$, Remark 1.15 also applies directly, and using Corollary 1.27,

$$\begin{aligned}
h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x) &= h_I i_0 \mu_n(\dots, \sigma \mu_{t_1}, \dots, \sigma \mu_{t_{j-1}}, \dots, i_1 \mu_q, \dots, \sigma \mu_{t_j}, \dots, \sigma \mu_{t_s}, \dots) \\
& - \sum i_0 p i_0(\mu_n)(\dots, h_I \sigma \mu_{t_i}, \dots, i_1 \mu_q, \dots) \\
& - i_0 p i_0(\mu_n)(\dots, i_0 p \sigma \mu_{t_i}, \dots, h_I i_1 \mu_q, \dots) \\
& + \sum i_0 p i_0(\mu_n)(\dots, i_0 p i_1 \mu_q, \dots, h_I \sigma \mu_{t_i}, \dots).
\end{aligned}$$

Using again that $h_I i_0$, $h_I \sigma = 0$, and $p\sigma = 0$, we deduce that all terms vanish unless $i_1 \mu_q$ is the first element in the brackets in x , i.e. $j = 1$. In that case, there is again a single non-vanishing term,

$$\begin{aligned}
h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x) &= -i_0 p i_0(\mu_n)(\dots, h_I i_1 \mu_q, \dots, \sigma \mu_{t_1}, \dots, \sigma \mu_{t_s}, \dots) \\
& = -i_0(\mu_n)(\dots, \sigma \mu_q, \dots, \sigma \mu_{t_1}, \dots, \sigma \mu_{t_s}, \dots).
\end{aligned}$$

We have finally established the formula for $h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x)$. Let us see that it coincides with $h_{\Lambda \mathcal{A}_n}(x)$. Using the induction hypothesis for the formula of $h_{\Lambda \mathcal{A}_{n-1}} i_1 d(\mu_n)$, we see that $\partial_{n-1, I} h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x)$ is a linear combination of standard labeled trees satisfying the hypotheses of Lemma 1.29 (2). Therefore,

$$h_{\Lambda \mathcal{A}_n} \partial_{n-1, I} h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x) = 0.$$

This implies $h_{\Lambda \mathcal{A}_n}(x) = h_{\Lambda \mathcal{A}_{n-1}, \mathbb{k}, \mu_n}(x)$ by Remark 1.19.

We now attack the final formula. In order to apply $h_{\Lambda\mathcal{A}_n}$, we divide $i_1d(\mu_{n+1})$ in two three blocks, according to the parity of n ,

$$\begin{aligned} i_1d(\mu_{n+1}) &= \sum_{\substack{p+q=n+2 \\ p < q}} i_1\mu_p\{i_1\mu_q\} + \sum_{\substack{p+q=n+2 \\ p > q}} i_1\mu_p\{i_1\mu_q\} \\ &\quad + i_1\mu_{\frac{n}{2}+1}\{i_1\mu_{\frac{n}{2}+1}\} \quad \text{if } n \text{ is even,} \end{aligned}$$

which will be considered in this order.

If $p < q$, $i_1\mu_p\{i_1\mu_q\} \in I\Lambda\mathcal{A}_q$ is in filtration degree 1, hence

$$\begin{aligned} h_{\Lambda\mathcal{A}_n}(i_1\mu_p\{i_1\mu_q\}) &= h_{\Lambda\mathcal{A}_q}(i_1\mu_p\{i_1\mu_q\}) \\ &= h_{\Lambda\mathcal{A}_{q-1}, \mathbb{k}\cdot\mu_q}(i_1\mu_p\{i_1\mu_q\}) \\ (2.12) \quad &\quad + h_{\Lambda\mathcal{A}_{q-1}}\partial_{q-1, I}h_{\Lambda\mathcal{A}_{q-1}, \mathbb{k}\cdot\mu_q}(i_1\mu_p\{i_1\mu_q\}). \end{aligned}$$

By Remark 2.9 and Corollary 1.27, the first summand is

$$\begin{aligned} h_{\Lambda\mathcal{A}_{q-1}, \mathbb{k}\cdot\mu_q}(i_1\mu_p\{i_1\mu_q\}) &= h_{\Lambda\mathcal{A}_{q-1}}i_1\mu_p\{i_1\mu_q\} - i_0p i_1\mu_p\{h_I i_1\mu_q\} \\ (2.13) \quad &= \sigma\mu_p\{i_1\mu_q\} - i_0\mu_p\{\sigma\mu_q\}. \end{aligned}$$

Let us see that (2.12) = 0. If we apply $\partial_{q-1, I}$ to the summands in (2.13), we obtain

$$\partial_{q-1, I}(\sigma\mu_p\{i_1\mu_q\}) = \sigma\mu_p\{i_1d(\mu_q)\}, \quad \partial_{q-1, I}(i_0\mu_p\{\sigma\mu_q\}) = i_0\mu_p\{h_{\Lambda\mathcal{A}_{q-1}}i_1d(\mu_q)\}.$$

The first element is a linear combination of standard labeled trees as in Lemma 1.29 (1). Moreover, using the induction hypothesis for $h_{\Lambda\mathcal{A}_{q-1}}i_1d(\mu_q)$ we see that the second term is a linear combination of standard labeled trees as in Lemma 1.29 (2). Therefore, this lemma proves that

$$h_{\Lambda\mathcal{A}_{q-1}}(\sigma\mu_p\{i_1d(\mu_q)\}) = 0, \quad h_{\Lambda\mathcal{A}_{q-1}}(i_0\mu_p\{h_{\Lambda\mathcal{A}_{q-1}}i_1d(\mu_q)\}) = 0.$$

If $p > q$, $i_1\mu_p\{i_1\mu_q\} \in I\Lambda\mathcal{A}_p$ is in filtration degree 1 and

$$\begin{aligned} h_{\Lambda\mathcal{A}_n}(i_1\mu_p\{i_1\mu_q\}) &= h_{\Lambda\mathcal{A}_p}(i_1\mu_p\{i_1\mu_q\}) \\ &= h_{\Lambda\mathcal{A}_{p-1}, \mathbb{k}\cdot\mu_p}(i_1\mu_p\{i_1\mu_q\}) \\ (2.14) \quad &\quad + h_{\Lambda\mathcal{A}_{p-1}}\partial_{p-1, I}h_{\Lambda\mathcal{A}_{p-1}, \mathbb{k}\cdot\mu_p}(i_1\mu_p\{i_1\mu_q\}). \end{aligned}$$

As above,

$$\begin{aligned} h_{\Lambda\mathcal{A}_{p-1}, \mathbb{k}\cdot\mu_p}(i_1\mu_p\{i_1\mu_q\}) &= h_I i_1\mu_p\{i_1\mu_q\} - i_0p i_1\mu_p\{h_{\Lambda\mathcal{A}_{p-1}}i_1\mu_q\} \\ (2.15) \quad &= \sigma\mu_p\{i_1\mu_q\} - i_0\mu_p\{\sigma\mu_q\}. \end{aligned}$$

In this case, (2.14) need not be 0. Using the induction hypothesis for $h_{\Lambda\mathcal{A}_{p-1}}i_1d(\mu_p)$, we see that $\partial_{p-1, I}$ of the two summands in (2.15) is

$$\begin{aligned} \partial_{p-1, I}(i_0\mu_p\{\sigma\mu_q\}) &= i_0d(\mu_p)\{\sigma\mu_q\}, \\ \partial_{p-1, I}(\sigma\mu_p\{i_1\mu_q\}) &= -h_{\Lambda\mathcal{A}_{p-1}}i_1d(\mu_p)\{i_1\mu_q\} \\ &= -\sum_{k+l=p+1} \sigma\mu_k\{i_1\mu_l\}\{i_1\mu_q\} \end{aligned}$$

$$+ \sum_{\substack{1 \leq s \leq r \\ t_1 + \dots + t_s = p + s - r}} i_0 \mu_r \{ \sigma \mu_{t_1}, \dots, \sigma \mu_{t_s} \} \{ i_1 \mu_q \}.$$

The brace relation yields

$$\begin{aligned} \sigma \mu_k \{ i_1 \mu_l \} \{ i_1 \mu_q \} &= \sigma \mu_k \{ i_1 \mu_l, i_1 \mu_q \} + \sigma \mu_k \{ i_1 \mu_l \{ i_1 \mu_q \} \} \\ &\quad - \sigma \mu_k \{ i_1 \mu_q, i_1 \mu_l \}, \\ i_0 \mu_r \{ \sigma \mu_{t_1}, \dots, \sigma \mu_{t_s} \} \{ i_1 \mu_q \} &= \sum_{j=1}^s i_0 \mu_r \{ \dots, \sigma \mu_{t_j} \{ i_1 \mu_q \}, \dots \} \\ &\quad + \sum_{j=1}^{s+1} i_0 \mu_r \{ \dots, \sigma \mu_{t_{j-1}}, i_1 \mu_q, \sigma \mu_{t_j}, \dots \} \text{ if } s < r, \end{aligned}$$

see Remark 2.7.

By Lemma 1.29,

$$h_{\Lambda \mathcal{A}_{p-1}}(\sigma \mu_k \{ i_1 \mu_l \} \{ i_1 \mu_q \}) = 0, \quad h_{\Lambda \mathcal{A}_{p-1}}(i_0 \mu_r \{ \dots, \sigma \mu_{t_j} \{ i_1 \mu_q \}, \dots \}) = 0.$$

Moreover, by the first series of equations in the statement, already checked up to n ,

$$h_{\Lambda \mathcal{A}_{p-1}}(i_0 \mu_r \{ \dots, \sigma \mu_{t_{j-1}}, i_1 \mu_q, \sigma \mu_{t_j}, \dots \}) = \begin{cases} -i_0 \mu_r \{ \dots, \sigma \mu_{t_{j-1}}, \sigma \mu_q, \sigma \mu_{t_j}, \dots \} \\ \quad \text{if } t_1, \dots, t_{j-1} > r \geq q, \\ \quad \text{or if } t_1, \dots, t_{j-1} > q > r \\ \quad \quad \text{and } q \leq t_j, \dots, t_s; \\ 0, \quad \text{otherwise.} \end{cases}$$

If n is even, by Remark 1.19,

$$\begin{aligned} h_{\Lambda \mathcal{A}_n}(i_1 \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \}) &= h_{\Lambda \mathcal{A}_{\frac{n}{2}, \mathbb{k} \cdot \mu_{\frac{n}{2}+1}}}(i_1 \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \}) \\ &\quad + h_{\Lambda \mathcal{A}_{\frac{n}{2}+1}} \partial_{\frac{n}{2}, I} h_{\Lambda \mathcal{A}_{\frac{n}{2}, \mathbb{k} \cdot \mu_{\frac{n}{2}+1}}}(i_1 \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \}). \end{aligned}$$

The first summand is computed as in the previous two cases,

$$\begin{aligned} h_{\Lambda \mathcal{A}_{\frac{n}{2}, \mathbb{k} \cdot \mu_{\frac{n}{2}+1}}}(i_1 \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \}) &= h_I i_1 \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \} - i_0 p i_1 \mu_{\frac{n}{2}+1} \{ h_I i_1 \mu_{\frac{n}{2}+1} \} \\ &= \sigma \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \} - i_0 \mu_{\frac{n}{2}+1} \{ \sigma \mu_{\frac{n}{2}+1} \}, \end{aligned}$$

Let us check that the second one vanishes. We have

$$\begin{aligned} \partial_{\frac{n}{2}, I}(i_0 \mu_{\frac{n}{2}+1} \{ \sigma \mu_{\frac{n}{2}+1} \}) &= i_0 d(\mu_{\frac{n}{2}+1}) \{ \sigma \mu_{\frac{n}{2}+1} \} + i_0 \mu_{\frac{n}{2}+1} \{ h_{\Lambda \mathcal{A}_{\frac{n}{2}}} i_1 d(\mu_{\frac{n}{2}+1}) \}, \\ \partial_{\frac{n}{2}, I}(\sigma \mu_{\frac{n}{2}+1} \{ i_1 \mu_{\frac{n}{2}+1} \}) &= -h_{\Lambda \mathcal{A}_{\frac{n}{2}}} i_1 d(\mu_{\frac{n}{2}+1}) \{ i_1 \mu_{\frac{n}{2}+1} \} + \sigma \mu_{\frac{n}{2}+1} \{ i_1 d(\mu_{\frac{n}{2}+1}) \}. \end{aligned}$$

Clearly $i_0 d(\mu_{\frac{n}{2}+1}) \{ \sigma \mu_{\frac{n}{2}+1} \}$ and $\sigma \mu_{\frac{n}{2}+1} \{ i_1 d(\mu_{\frac{n}{2}+1}) \}$ are linear combinations of standard labeled trees as in Lemma 1.29. Using the formula for $h_{\Lambda \mathcal{A}_{\frac{n}{2}}} i_1 d(\mu_{\frac{n}{2}+1})$,

that we know by induction hypothesis, we see that $i_0\mu_{\frac{n}{2}+1}\{h_{\Lambda\mathcal{A}_{\frac{n}{2}}}\}i_1d(\mu_{\frac{n}{2}+1})$ too. Therefore,

$$\begin{aligned} h_{\Lambda\mathcal{A}_{\frac{n}{2}+1}}(i_0d(\mu_{\frac{n}{2}+1})\{\sigma\mu_{\frac{n}{2}+1}\}) &= 0, & h_{\Lambda\mathcal{A}_{\frac{n}{2}+1}}(i_0\mu_{\frac{n}{2}+1}\{hi_1d(\mu_m)\}) &= 0, \\ h_{\Lambda\mathcal{A}_{\frac{n}{2}+1}}(\sigma\mu_{\frac{n}{2}+1}\{i_1d(\mu_{\frac{n}{2}+1})\}) &= 0. \end{aligned}$$

Moreover,

$$(2.16) \quad \begin{aligned} h_{\Lambda\mathcal{A}_{\frac{n}{2}}}\{i_1d(\mu_{\frac{n}{2}+1})\{i_1\mu_{\frac{n}{2}+1}\}\} &= \sum_{p+q=\frac{n}{2}+2} \sigma\mu_p\{i_1\mu_q\}\{i_1\mu_{\frac{n}{2}+1}\} \\ &- \sum_{\substack{1 \leq s \leq r \\ t_1 + \dots + t_s = \frac{n}{2} + 1 + s - r}} i_0\mu_r\{\sigma\mu_{t_1}, \dots, \sigma\mu_{t_s}\}\{i_1\mu_{\frac{n}{2}+1}\}. \end{aligned}$$

Using the brace equation, we can check as above that $\sigma\mu_p\{i_1\mu_q\}\{i_1\mu_{\frac{n}{2}+1}\}$ is a linear combination of standard labeled trees as in Lemma 1.29 (1), so

$$h_{\Lambda\mathcal{A}_{\frac{n}{2}+1}}(\sigma\mu_p\{i_1\mu_q\}\{i_1\mu_{\frac{n}{2}+1}\}) = 0.$$

Furthermore, the summands in (2.16) are

$$(2.17) \quad \begin{aligned} i_0\mu_r\{\sigma\mu_{t_1}, \dots, \sigma\mu_{t_s}\}\{i_1\mu_{\frac{n}{2}+1}\} &= \sum_{j=1}^s i_0\mu_r\{\dots, \sigma\mu_{t_j}\{i_1\mu_{\frac{n}{2}+1}\}, \dots\} \\ &+ \sum_{j=1}^{s+1} i_0\mu_r\{\dots, \sigma\mu_{t_{j-1}}, i_1\mu_{\frac{n}{2}+1}, \sigma\mu_{t_j}, \dots\} \text{ if } s < r, \end{aligned}$$

see Remark 2.7. Here, $i_0\mu_r\{\dots, \sigma\mu_{t_j}\{i_1\mu_{\frac{n}{2}+1}\}, \dots\}$ is a linear combination of standard labeled trees as in Lemma 1.29 (2), so

$$h_{\Lambda\mathcal{A}_{\frac{n}{2}+1}}(i_0\mu_r\{\dots, \sigma\mu_{t_j}\{i_1\mu_{\frac{n}{2}+1}\}, \dots\}) = 0.$$

Looking at the index of the summation in (2.16), we see that, in (2.17), $r, t_1, \dots, t_s \leq \frac{n}{2}$. Therefore, by the first series of equations in the statement, already checked up to n , we see that

$$h_{\Lambda\mathcal{A}_{\frac{n}{2}+1}}(i_0\mu_r\{\dots, \sigma\mu_{t_{j-1}}, i_1\mu_{\frac{n}{2}+1}, \sigma\mu_{t_j}, \dots\}) = 0.$$

Collecting previous equations, we obtain

$$\begin{aligned} hi_1d(\mu_{n+1}) &= \sum_{p+q=n+2} (\sigma\mu_p\{i_1\mu_q\} - i_0\mu_p\{\sigma\mu_q\}) \\ &- \overbrace{\sum_{\substack{p+q=n+2 \\ p>q}} \sum_{s=1}^{r-1} \sum_{j=1}^{s+1} \sum_{\substack{t_1+\dots+t_s=p+s-r \\ \text{and either} \\ t_1, \dots, t_{j-1} > r \geq q \\ \text{or} \\ t_1, \dots, t_{j-1} > q > r \\ \text{and } t_j, \dots, t_s \geq q}} i_0\mu_r\{\dots, \sigma\mu_{t_{j-1}}, \sigma\mu_q, \sigma\mu_{t_j}, \dots\}}^{(*)}. \end{aligned}$$

Now, it is enough to check that

$$(2.18) \quad (\star) = \sum_{\substack{2 \leq \bar{s} \leq \bar{r} \\ \bar{t}_1 + \dots + \bar{t}_{\bar{s}} = n+1 + \bar{s} - \bar{r}}} i_0 \mu_{\bar{r}} \{ \sigma \mu_{\bar{t}_1}, \dots, \sigma \mu_{\bar{t}_{\bar{s}}} \}.$$

The summand in (\star) corresponding to certain $p, q, j, r, s, t_1, \dots, t_s$ is the same as the summand on the right hand side of (2.18) corresponding to

$$\bar{r} = r, \quad \bar{s} = s + 1, \quad \bar{t}_i = t_i \text{ for } 1 \leq i < j, \quad \bar{t}_j = q, \quad \bar{t}_i = t_{i-1} \text{ for } j < i \leq \bar{s}.$$

Obviously $\bar{s} = s + 1 \geq 1 + 1 = 2$; $\bar{s} = s + 1 \leq r - 1 + 1 = \bar{r}$; and, using that $p + q = n + 2$,

$$\bar{t}_1 + \dots + \bar{t}_{\bar{s}} = t_1 + \dots + t_s + q = p + s - r + q = n + 1 + s + 1 - r = n + 1 + \bar{s} - \bar{r}.$$

Consider now a summand on the right hand side of (2.18), corresponding to certain $\bar{r}, \bar{s}, \bar{t}_1, \dots, \bar{t}_{\bar{s}}$. Suppose that some \bar{t}_i is smaller or equal than \bar{r} . Let j be the smallest value $1 \leq j \leq \bar{s}$ such that $\bar{t}_j \leq \bar{r}$, in particular $\bar{t}_1, \dots, \bar{t}_{j-1} > \bar{r} \geq \bar{t}_j$. Then the summand on the right hand side of (2.18) is the same as the summand in (\star) corresponding to

$$(2.19) \quad \begin{aligned} r &= \bar{r}, & t_i &= \bar{t}_i \text{ for } 1 \leq i < j, \\ s &= \bar{s} - 1, & t_{i-1} &= \bar{t}_i \text{ for } j < i \leq \bar{s}, \\ q &= \bar{t}_j, & p &= \bar{t}_1 + \dots + \bar{t}_{j-1} + \bar{t}_{j+1} + \dots + \bar{t}_{\bar{s}} + \bar{r} - \bar{s} + 1. \end{aligned}$$

Note that $s = \bar{s} - 1 \geq 2 - 1 = 1$; $1 \leq j \leq \bar{s} = s + 1$; $r = \bar{r} \geq \bar{s} > \bar{s} - 1 = s$; $p + q = \bar{t}_1 + \dots + \bar{t}_{\bar{s}} + \bar{r} - \bar{s} + 1 = n + 2$; for $1 \leq i < j$, $t_i = \bar{t}_i > \bar{r} = r \geq \bar{t}_j = q$; and, since $\bar{t}_i > 1$ for all i , $p > \bar{s} - 1 + \bar{r} - \bar{s} + 1 = \bar{r} \geq \bar{t}_j = q$.

Othwerwise, all \bar{t}_i s are bigger than \bar{r} . Let j be the smallest $1 \leq j \leq \bar{s}$ such that \bar{t}_j attains the minimum value among all \bar{t}_i s, in particular $\bar{t}_1, \dots, \bar{t}_{j-1} > \bar{t}_j > \bar{r}$ and $\bar{t}_{j+1}, \dots, \bar{t}_{\bar{s}} \geq \bar{t}_j$. Once again, one can straightforwardly check that the summand on the right hand side of (2.18) is the same as the summand in (\star) corresponding to the formulas in (2.19). \square

We now compute examples with non-trivial operations in arities 1 and 0.

Example 2.20. We can consider the following extension \mathcal{A}_∞^D of the A -infinity operad which has non-trivial elements in arity 1. As a graded operad, it is freely generated by

$$\mu_n \in \mathcal{A}_\infty^D(n)_{n-2}, \quad n \geq 2; \quad D_n \in \mathcal{A}_\infty^D(n)_{n-1}, \quad n \geq 1.$$

The differential on the new generators is given by

$$d(D_n) = \sum_{\substack{p+q=n+1 \\ 1 \leq i \leq p}} ((-1)^{(q-1)(i-1)} \mu_p \circ_i D_q - (-1)^{p-i+q(i-1)} D_p \circ_i \mu_q).$$

This is the Koszul resolution of the operad whose algebras are associative algebras equipped with a degree 0 derivation, considered by Loday in [Lod10]. We have simplified notation and adapted sign conventions to our setting. We should warn

the reader that [Lod10] contains an obvious mistake in the grading of generators coming from the A -infinity operad (in fact, Loday's differential would not be homogeneous with his grading). We have fixed this mistake and its consequences on signs. Let us sketch how the canonical strong cylinder $I\mathcal{A}_\infty^D$ can be computed as above, by using operadic suspensions.

The operadic suspension $\Lambda\mathcal{A}_\infty^D$ is freely generated as a graded operad by

$$\mu_n \in \Lambda\mathcal{A}_\infty^D(n)_{-1}, \quad n \geq 2; \quad D_n \in \Lambda\mathcal{A}_\infty^D(n)_0, \quad n \geq 1.$$

The differential of the generators not coming from $\Lambda\mathcal{A}_\infty$ is then

$$d(D_n) = \sum_{p+q=n+1} (\mu_p\{D_q\} - D_p\{\mu_q\}).$$

This operad is cellular with $\Lambda\mathcal{A}_0^D = \Lambda\mathcal{A}_1^D$ the initial DG-operad and

$$\Lambda\mathcal{A}_n^D = \Lambda\mathcal{A}_{n-1}^D \amalg_{\partial_{n-1}} \mathcal{F}(\mathbb{k} \cdot \{D_{n-1}, \mu_n\}), \quad n \geq 2.$$

Here ∂_{n-1} is defined as the differential. Lemma 2.10 can be extended to show

$$\begin{aligned} h_{\Lambda\mathcal{A}_n^D} i_1 d(D_n) &= \sum_{p+q=n+1} \sigma\mu_p\{i_1 D_q\} \\ &\quad - \sum_{\substack{0 \leq s < r \\ t_1 + \dots + t_s + q = n+1+s-r \\ 1 \leq j \leq s+1}} i_0 \mu_r \{ \sigma\mu_{t_1}, \dots, \sigma\mu_{t_{j-1}}, \sigma D_q, \sigma\mu_{t_j}, \dots, \sigma\mu_{t_s} \} \\ &\quad - \sum_{p+q=n+1} \sigma D_p\{i_1 \mu_q\} - \sum_{\substack{1 \leq s \leq r \\ t_1 + \dots + t_s = n+s-r}} i_0 D_r \{ \sigma\mu_{t_1}, \dots, \sigma\mu_{t_s} \}. \end{aligned}$$

This formula, together with (1.26) and the fact that the inclusions $I\Lambda\mathcal{A}_\infty \subset I\Lambda\mathcal{A}_\infty^D$ and $i_0, i_1: \Lambda\mathcal{A}_\infty^D \rightarrow I\Lambda\mathcal{A}_\infty^D$ are DG-operad morphisms, completely determines $I\Lambda\mathcal{A}_\infty^D$ as a DG-operad, and hence $I\mathcal{A}_\infty^D$.

Our canonical strong pseudo-cylinder generalizes the classical cylinder of DG-algebras, regarded as DG-operads concentrated in arity 1. This is a consequence the following result, see (1.26) and [Bau89, §I.7].

Lemma 2.21. *Let \mathcal{O} be an absolutely pseudo-cellular DG-operad of length α as in Definition 1.24, such that V_β is concentrated in arities 0 and 1, $\beta < \alpha$. Then, the following equation holds for any $x \in \mathcal{O}(1)$ and any $y \in \mathcal{O}$,*

$$h_{\mathcal{O}} i_1(x \circ_1 y) = h_{\mathcal{O}} i_1(x) \circ_1 i_1(y) + (-1)^{|x|} i_0(x) \circ_1 h_{\mathcal{O}} i_1(y).$$

Proof. A simple computation shows that the statement follows if we prove that, given $x_i \in V_{\beta_i}(1)$, $1 \leq i < n$, and $x_n \in V_{\beta_n}$,

$$\begin{aligned} h_{\mathcal{O}} i_1(x_1 \circ_1 \dots \circ_1 x_n) &= h_{\mathcal{O}_{\gamma+1}} i_1(x_1 \circ_1 \dots \circ_1 x_n) \\ &= h_{\mathcal{O}_{\gamma}, V_{\gamma}} i_1(x_1 \circ_1 \dots \circ_1 x_n) \end{aligned}$$

$$= \sum_{j=1}^n (-1)^{\sum_{k=1}^{j-1} |x_k|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n),$$

where $\gamma = \max\{\beta_1, \dots, \beta_n\}$. The first equation holds by definition. We check the other two ones by induction, first on γ and then on the filtration degree of $x_1 \circ_1 \cdots \circ_1 x_n \in \mathcal{O}_\gamma \amalg_{\partial_\gamma} \mathcal{F}(V_\gamma)$ with respect to the filtration in the proof of Theorem 1.21, i.e. on the (positive) amount of numbers $1 \leq i \leq n$ with $\beta_i = \gamma$. Let i be the smallest $1 \leq i \leq n$ such that $\beta_i = \gamma$. Then

$$\begin{aligned} & h_{\mathcal{O}_\gamma, V_\gamma}(x_1 \circ_1 \cdots \circ_1 x_n) \\ &= h_{\mathcal{O}_\gamma} i_1(x_1 \circ_1 \cdots \circ_1 x_{i-1}) \circ_1 i_1(x_i \circ_1 \cdots \circ_1 x_n) \\ &\quad + (-1)^{\sum_{j=0}^{i-1} |x_j|} i_0 p i_1(x_1 \circ_1 \cdots \circ_1 x_{i-1}) \circ_1 h_{\mathcal{O}_\gamma, V_\gamma} i_1(x_i \circ_1 \cdots \circ_1 x_n) \\ &= h_{\mathcal{O}_\gamma} i_1(x_1 \circ_1 \cdots \circ_1 x_{i-1}) \circ_1 i_1(x_i \circ_1 \cdots \circ_1 x_n) \\ &\quad + (-1)^{\sum_{j=1}^{i-1} |x_j|} i_0 p i_1(x_1 \circ_1 \cdots \circ_1 x_{i-1}) \circ_1 h_I i_1(x_i) \circ_1 i_1(x_{i+1} \circ_1 \cdots \circ_1 x_n) \\ &\quad + (-1)^{\sum_{j=1}^i |x_j|} i_0 p i_1(x_1 \circ_1 \cdots \circ_1 x_{i-1}) \circ_1 i_0 p i_1(x_i) \circ_1 h_{\mathcal{O}_\gamma, V_\gamma} i_1(x_{i+1} \circ_1 \cdots \circ_1 x_n) \\ &= \sum_{j=1}^{i-1} (-1)^{\sum_{k=1}^{j-1} |x_k|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n) \\ &\quad + (-1)^{\sum_{j=1}^{i-1} |x_j|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{i-1}) \circ_1 \sigma(x_i) \circ_1 i_1(x_{i+1}) \circ_1 \cdots \circ_1 i_1(x_n) \\ &\quad + \sum_{j=i+1}^n (-1)^{\sum_{k=1}^{j-1} |x_k|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n) \\ &= \sum_{j=1}^n (-1)^{\sum_{k=1}^{j-1} |x_k|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n). \end{aligned}$$

In the first two equations, we use Remark 1.15. In the third one, we apply the two induction hypotheses. Indeed, $x_1 \circ_1 \cdots \circ_1 x_{i-1} \in \mathcal{O}_\gamma$ and either $x_{i+1} \circ_1 \cdots \circ_1 x_n \in \mathcal{O}_\gamma \amalg \mathcal{F}(V_\gamma)$ has positive but smaller filtration degree or $x_{i+1} \circ_1 \cdots \circ_1 x_n \in \mathcal{O}_\gamma$.

In order to check that $h_{\mathcal{O}_{\gamma+1}}(x_1 \circ_1 \cdots \circ_1 x_n) = h_{\mathcal{O}_\gamma, V_\gamma}(x_1 \circ_1 \cdots \circ_1 x_n)$, observe that

$$\begin{aligned} & \partial_{\gamma, I}(i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n)) \\ &= -(-1)^{\sum_{k=1}^{j-1} |x_k|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 h_{\mathcal{O}_\gamma} i_1 \partial_\gamma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n) \\ &\quad - (-1)^{\sum_{k=1}^j |x_k|} i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1 \partial_\gamma(x_{j+1} \circ_1 \cdots \circ_1 x_n). \end{aligned}$$

The second summand is clearly a linear combination of standard labeled trees as in Lemma 1.29, hence

$$h_{\mathcal{O}_{\gamma+1}}(i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 \sigma(x_j) \circ_1 i_1 \partial_\gamma(x_{j+1} \circ_1 \cdots \circ_1 x_n)) = 0.$$

Moreover, using the induction hypothesis on $h_{\mathcal{O}_\gamma} i_1$, we see that the first summand is also such a linear combination, therefore

$$h_{\mathcal{O}_\gamma, V_\gamma}(i_0(x_1) \circ_1 \cdots \circ_1 i_0(x_{j-1}) \circ_1 h_{\mathcal{O}_\gamma} i_1 \partial_\gamma(x_j) \circ_1 i_1(x_{j+1}) \circ_1 \cdots \circ_1 i_1(x_n)) = 0.$$

This proves that $h_{\mathcal{O}_{\gamma+1}}(x_1 \circ_1 \cdots \circ_1 x_n) = h_{\mathcal{O}_\gamma, V_\gamma}(x_1 \circ_1 \cdots \circ_1 x_n)$ by Remark 1.19. \square

In the conditions of the previous lemma, the operad \mathcal{O} consists of just a DG-algebra $\mathcal{O}(1)$ and a left $\mathcal{O}(1)$ -module $\mathcal{O}(0)$. It is trivial in higher arities. The full computation of $h_{\mathcal{O}} i_1$ has been possible in this case since all labeled trees in \mathcal{O} are linear and the path order coincides with the linear order.

3. LINEAR DG-OPERADS

In this section we analyze the canonical strong pseudo-cylinder construction in a class of relatively pseudo-cellular DG-operads, that we call *linear*, where formulas are easy. Classical examples, such as the A -infinity DG-operad, are not linear, but relative examples do show up, as we will see below.

Given a graded operad \mathcal{O} , recall that an \mathcal{O} -module [Mar96, Definition 1.4] is a sequence of graded modules $M = \{M(n)\}_{n \geq 0}$ equipped with compositions, $1 \leq i \leq p$, $q \geq 0$

$$\circ_i: M(p) \otimes \mathcal{O}(q) \longrightarrow M(p+q-1), \quad \circ_i: \mathcal{O}(p) \otimes M(q) \longrightarrow M(p+q-1),$$

satisfying the same laws as graded operads (1.4) when one of the variables is in M and the rest in \mathcal{O} . These are the same as the linear modules introduced in [BJT97, Definition 2.13] and the infinitesimal bimodules from [MV09a, §3.1]. Any graded operad is a module over itself, and restriction of scalars is defined in the obvious way.

The (aritywise) suspension ΣM of an \mathcal{O} -module M is again an \mathcal{O} -module with structure

$$(\sigma x) \circ_i y = \sigma(x \circ_i y), \quad y \circ_i (\sigma x) = (-1)^{|y|} \sigma(y \circ_i x), \quad x \in M, \quad y \in \mathcal{O}.$$

Suppose now that \mathcal{O} is a relatively pseudo-cellular DG-operad of length α , as in Definition 1.24. Recall that its underlying graded operad is $\mathcal{O}_0 \amalg \mathcal{F}(V)$, $V = \bigoplus_{\beta < \alpha} V_\beta$. The sub- \mathcal{O}_0 -module of \mathcal{O} spanned by the identity element $\text{id}_{\mathcal{O}}$ is \mathcal{O}_0 . The sub- \mathcal{O}_0 -module $\langle V \rangle_{\mathcal{O}_0} \subset \mathcal{O}$ spanned by V corresponds, in the direct sum decomposition for graded operad coproducts (1.12), to the direct subsum

The maps $i_0, i_1: \mathcal{O}_\beta \rightarrow I\mathcal{O}_\beta$ (co)restrict to the left and right vertical \mathcal{O}_0 -module maps in the following diagram, respectively,

$$\begin{array}{c} \mathcal{O}_0 \oplus \langle \bigoplus_{\gamma < \beta} V_\gamma \rangle_{\mathcal{O}_0} \\ \left(\begin{array}{c} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{array} \right) \downarrow \downarrow \left(\begin{array}{c} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{array} \right) \\ \mathcal{O}_0 \oplus \langle \bigoplus_{\gamma < \beta} V_\gamma \rangle_{\mathcal{O}_0} \oplus \langle \bigoplus_{\gamma < \beta} \Sigma V_\gamma \rangle_{\mathcal{O}_0} \oplus \langle \bigoplus_{\gamma < \beta} V_\gamma \rangle_{\mathcal{O}_0} \end{array}$$

Since $\partial_{\beta, I} = (i_0 \partial_\beta \quad -h_{\mathcal{O}_\beta} i_1 \partial_\beta \quad i_1 \partial_\beta)$, using the induction hypothesis on $h_{\mathcal{O}_\beta}$, we deduce that

$$h_{\mathcal{O}_\beta} \left(\begin{array}{c} \dots \quad \dots \\ \cdot \quad \cdot \\ x_1 \quad \dots \quad x_n \\ \cdot \quad \cdot \quad \cdot \\ \partial_{\beta, I} h_I(y) \quad \dots \\ \cdot \\ x_0 \end{array} \right) = (-1)^{|x_0|} \begin{array}{c} \dots \quad \dots \\ \cdot \quad \cdot \\ x_1 \quad \dots \quad x_n \\ \cdot \quad \cdot \quad \cdot \\ \phi(y) \quad \dots \\ \cdot \\ x_0 \end{array},$$

where $\phi = h_{\mathcal{O}_\beta} \partial_{\beta, I} h_I$ is the following product of (block) matrices,

$$\phi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma \\ 0 & 0 & 0 & 0 \end{pmatrix} \left(\begin{array}{c} \left(\begin{array}{c} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{array} \right) \left(\begin{array}{c} \partial_\beta^0 \\ \partial_\beta^1 \end{array} \right) \\ \hline 0 \\ \hline \left(\begin{array}{c} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{array} \right) \left(\begin{array}{c} \partial_\beta^0 \\ \partial_\beta^1 \end{array} \right) \end{array} \right) \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \partial_\beta^0 \\ \partial_\beta^1 \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \sigma \\ 0 & 0 & 0 \end{pmatrix}.$$

A straightforward computation shows that $\phi = 0$, hence the previous claim follows.

As part of the proof of the previous statement, we have shown that, for any linear DG-operad \mathcal{O} of any length α , and for any $\beta < \alpha$, $h_{\mathcal{O}_\beta} i_1 \partial_\beta: V_\beta \rightarrow I\mathcal{O}_\beta$ corestricts to

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sigma \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \partial_\beta^0 \\ \partial_\beta^1 \end{pmatrix} = \begin{pmatrix} 0 \\ \sigma \partial_\beta^1 \\ 0 \end{pmatrix}: V_\beta \longrightarrow \langle \bigoplus_{\gamma < \beta} V_\gamma \rangle_{\mathcal{O}_0} \oplus \langle \bigoplus_{\gamma < \beta} \Sigma V_\gamma \rangle_{\mathcal{O}_0} \oplus \langle \bigoplus_{\gamma < \beta} V_\gamma \rangle_{\mathcal{O}_0}.$$

The statement of the proposition is equivalent to this, see (1.26). \square

Example 3.2. Let \mathcal{O} be the operad obtained by quotienting out μ_n , $n \geq 3$, from \mathcal{A}_∞^D in Example 2.20. Its algebras are non-unital DG-algebras with an up-to-homotopy derivation. As a graded operad, \mathcal{O} is generated by

$$\mu_2 \in \mathcal{O}_\infty(2)_0, \quad D_n \in \mathcal{O}_\infty(n)_{n-1}, \quad n \geq 1,$$

with a single relation

$$\mu_2 \circ_1 \mu_2 = \mu_2 \circ_2 \mu_2,$$

and differential

$$d(\mu_2) = 0, \quad d(D_n) = \mu_2 \circ_1 D_{n-1} + (-1)^n \mu_2 \circ_2 D_{n-1} + \sum_{i=1}^{n-1} (-1)^{n+i} D_{n-1} \circ_i \mu_2.$$

The operadic suspension $\Lambda\mathcal{O}$ is therefore generated by

$$\mu_2 \in \Lambda\mathcal{O}_\infty(2)_{-1}, \quad D_n \in \Lambda\mathcal{O}_\infty(n)_0, \quad n \geq 1,$$

with a single relation, the same as above, and differential

$$d(\mu_2) = 0, \quad d(D_n) = \mu_2\{D_{n-1}\} - D_{n-1}\{\mu_2\}.$$

We regard $\Lambda\mathcal{O}$ as a relatively cellular DG-operad with \mathcal{O}_0 the associative DG-operad, i.e. $\Lambda\mathcal{O}_0$ is the sub-DG-operad generated by μ_2 , and

$$\Lambda\mathcal{O}_n = \Lambda\mathcal{O}_{n-1} \amalg_{\partial_{n-1}} \mathcal{F}(\mathbb{k} \cdot D_n), \quad \partial_{n-1}(D_n) = d(D_n).$$

Then it is strictly linear with

$$d^1(D_n) = d(D_n), \quad n \geq 1.$$

Therefore, Proposition 3.1 yields

$$\begin{aligned} d(\sigma D_n) &= i_0(D_n) - i_1(D_n) - \sigma d(D_n) \\ &= i_0(D_n) - i_1(D_n) - \sigma(\mu_2\{D_{n-1}\} - D_{n-1}\{\mu_2\}) \\ &= i_0(D_n) - i_1(D_n) + \mu_2\{\sigma D_{n-1}\} + \sigma D_{n-1}\{\mu_2\}. \end{aligned}$$

It is easy to check that this is indeed the formula obtained from (1.26) and the formula for $h_{\Lambda\mathcal{A}_\infty^D} i_1 d(D_n)$ in Example 2.20, killing $\sigma\mu_n$ for $n \geq 2$, $i_0\mu_n, i_1\mu_n$ for $n \geq 3$, and identifying $i_0\mu_2 = i_1\mu_2 = \mu_2$.

In the following example, a non-strict linear DG-operad is also considered.

Example 3.3. The following linear relatively pseudo-cellular DG-operad \mathcal{O} appears in [Mur14]. Fix some $m > 0$. As a graded operad, \mathcal{O} is generated by

$$u \in \mathcal{O}(0)_0, \quad \mu \in \mathcal{O}(2)_0, \quad \nu_n^S \in \mathcal{O}(n-m)_{n-2+m},$$

where $n \geq m$ and $S \subset \{1, \dots, n\}$ runs over all subsets of cardinality m , with relations

$$\mu \circ_1 \mu = \mu \circ_2 \mu, \quad \mu \circ_1 u = \text{id} = \mu \circ_2 u.$$

The differential is defined by

$$d(u) = 0, \quad d(\mu) = 0;$$

if $(n, m) \neq (2, 1), (1, 1)$,

$$\begin{aligned} d(\nu_n^S) &= (-1)^n \mu \circ_1 \nu_{n-1}^S \quad \text{unless } l_m = n \\ &\quad + \mu \circ_2 \nu_{n-1}^{S-1} \quad \text{unless } l_1 = 1 \\ &\quad + \sum_{\substack{1 \leq v \leq m+1 \\ l_{v-1} < i+v-1 < l_v-1}} (-1)^{i+v-1} \nu_{n-1}^{S_v \cup (S'_v-1)} \circ_i \mu; \end{aligned}$$

and if $m = 1$ also

$$d(\nu_1^{\{1\}}) = 0, \quad d(\nu_2^{\{1\}}) = \mu \circ_1 \nu_1^{\{1\}} - \text{id}, \quad d(\nu_2^{\{2\}}) = \mu \circ_2 \nu_1^{\{1\}} - \text{id}.$$

Here we denote $S = \{l_1, \dots, l_{|S|}\}$, $l_0 = 0$, $l_{|S|+1} = n+1$, $S+t = \{l_1+t, \dots, l_{|S|}+t\}$, and

$$S_v = \{l_1, \dots, l_{v-1}\}, \quad S'_v = S \setminus S_v = \{l_v, \dots, l_{|S|}\}, \quad 1 \leq v \leq |S| + 1.$$

Unlike in previous cases, operadic suspension and braces do not simplify the definition of \mathcal{O} .

This DG-operad is relatively cellular with $\mathcal{O}_0 = \dots = \mathcal{O}_{m-1}$ the unital associative operad, i.e. the suboperad generated by u and μ , and

$$\mathcal{O}_n = \mathcal{O}_{n-1} \amalg_{\partial_{n-1}} \mathcal{F}(\mathbb{k} \cdot \{\nu_n^S\}_{\substack{S \subset \{1, \dots, n\} \\ |S|=m}}), \quad n \geq m.$$

Here ∂_{n-1} is defined as the differential above. Note that \mathcal{O} is clearly linear, even strictly for $m > 1$, but not for $m = 1$, since the constant part of the differential satisfies

$$d^0(\nu_2^{\{1\}}) = d^0(\nu_2^{\{2\}}) = -\text{id}.$$

Consider the retraction $r: \mathcal{O} \rightarrow \mathcal{O}_0$ defined as follows,

$$r(\nu_n^S) = 0 \text{ if } (n, m) \neq (1, 1), \quad r(\nu_1^{\{1\}}) \mapsto u \text{ if } m = 1.$$

Compatibility with differentials is checked in [Mur14, Lemma 5.9]. Denote by $j: \mathcal{O}_0 \rightarrow \mathcal{O}$ the inclusion. The equations in the proof of [Mur14, Lemma 5.9] show that there is a homotopy $H: I\mathcal{O} \rightarrow \mathcal{O}$, relative to \mathcal{O}_0 , from the identity in \mathcal{O} to jr , defined by

$$H(i_0\nu_n^S) = \nu_n^S, \quad H(\sigma\nu_n^S) = (-1)^{l_1+1}\nu_{n+1}^{S+1} \circ_{l_1} u, \quad H(i_1\nu_n^S) = jr(\nu_n^S).$$

Therefore j is the inclusion of a strong deformation retract, in particular a homotopy equivalence.

We finally compute some structure maps on canonical strong pseudo-cylinders of linear DG-operads.

Definition 3.4. Let \mathcal{O} be a relatively pseudo-cellular DG-operad. Consider the pasting of two canonical strong pseudo-cylinders $I\mathcal{O}_{i_1 \cup i_0} I\mathcal{O}$ obtained by identifying the top copy of \mathcal{O} ($i_1\mathcal{O}$) in the first $I\mathcal{O}$ with the bottom copy of \mathcal{O} ($i_0\mathcal{O}$) in the second $I\mathcal{O}$. A *doubling map* is a map

$$\nu: I\mathcal{O} \longrightarrow I\mathcal{O}_{i_1 \cup i_0} I\mathcal{O},$$

compatible with the projections onto \mathcal{O} , sending the bottom (resp. top) copy of \mathcal{O} in the source to the bottom (resp. top) copy of \mathcal{O} in the first (resp. second) $I\mathcal{O}$ in the target. A *reversing map* is a map

$$\iota: I\mathcal{O} \longrightarrow I\mathcal{O}$$

compatible with the projections onto \mathcal{O} , sending the bottom (resp. top) copy of \mathcal{O} in the source to the top (resp. bottom) copy of \mathcal{O} in the target.

Doubling and reversing maps allow to vertically compose and invert homotopies. They are important since, if \mathcal{O} is *based*, i.e. equipped with a retraction $\mathcal{O} \rightarrow \mathcal{O}_0$, they give rise to the up-to-homotopy cogroup structure of the model theoretic (relative) suspension $\Sigma\mathcal{O}$ of \mathcal{O} and to the coaction of $\Sigma\mathcal{O}$ on the relative cone $C\mathcal{O}$. Using the chain homotopy in $I\mathcal{O}$, we could give formulas for doubling and reversing maps in all cases. Formulas are not easy in general, so we will content ourselves with the linear case.

Proposition 3.5. *Let us place ourselves in the context of Proposition 3.1. Denote by $j_0, j_1: I\mathcal{O} \rightarrow I\mathcal{O}_{i_1} \cup_{i_0} I\mathcal{O}$ the inclusion of the first and second factor, respectively, which satisfy $j_0 i_1 = j_1 i_0$. The following formulas define a doubling map and a reversing map in the sense of Definition 3.4, $x \in V$,*

$$\begin{aligned} \nu i_0(x) &= j_0 i_0(x), & \nu i_1(x) &= j_1 i_1(x), & \nu(\sigma x) &= j_0 \sigma x + j_1 \sigma x, \\ i_0(x) &= i_1(x), & i_1(x) &= i_0(x), & \iota(\sigma x) &= -\sigma x. \end{aligned}$$

Proof. The conditions for ν to be a doubling map are $(p, p)\nu = p$, $\nu i_0 = j_0 i_0$, and $\nu i_1 = j_1 i_1$. The only part about ν which is not completely trivial is compatibility with differentials in the third case. In order to check this, note that the third formula is actually true for any $x \in \langle V \rangle_{\mathcal{O}_0}$, since i_0, i_1, j_0, j_1 are maps relative to \mathcal{O}_0 . Therefore,

$$\begin{aligned} d\nu(\sigma x) &= dj_0 \sigma x + dj_1 \sigma x \\ &= j_0 d(\sigma x) + j_1 d(\sigma x) \\ &= j_0 i_0(x) - j_0 i_1(x) - j_0 \sigma d^1(x) \\ &\quad + j_1 i_0(x) - j_1 i_1(x) - j_1 \sigma d^1(x) \\ &= j_0 i_0(x) - j_1 i_1(x) - j_0 \sigma d^1(x) - j_1 \sigma d^1(x) \\ &= \nu i_0(x) - \nu i_1(x) - \nu \sigma d^1(x) \\ &= \nu d(\sigma x). \end{aligned}$$

One can similarly check that ι is a reversing map. □

REFERENCES

- [Bau89] H.-J. Baues, *Algebraic Homotopy*, Cambridge University Press, 1989.
- [BJT97] H.-J. Baues, M. Jibladze, and A. Tonks, *Cohomology of monoids in monoidal categories*, Proceedings of the Renaissance Conferences (Providence, RI) (J.-L. Loday, J. D. Stasheff, and A. A. Voronov, eds.), Contemp. Math., vol. 202, Amer. Math. Soc., 1997, pp. 137–165.
- [Bro65] R. Brown, *The twisted Eilenberg-Zilber theorem*, Simposio di Topologia (Messina, 1964), Edizioni Oderisi, Gubbio, 1965, pp. 33–37.
- [Fre09] Benoit Fresse, *Operadic cobar constructions, cylinder objects and homotopy morphisms of algebras over operads*, Alpine perspectives on algebraic topology, Contemp. Math., vol. 504, Amer. Math. Soc., Providence, RI, 2009, pp. 125–188.
- [HM12] J. Hirsh and J. Millès, *Curved Koszul duality theory*, Math. Ann. **354** (2012), no. 4, 1465–1520.

- [Hov99] M. Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999.
- [Lam93] L. A. Lambe, *Resolutions which split off of the bar construction*, J. Pure Appl. Algebra **84** (1993), no. 3, 311–329.
- [LH03] K. Lefèvre-Hasegawa, *Sur les A_∞ -catégories*, Ph.D. thesis, Université Paris 7, 2003, [arXiv:math/0310337v1](https://arxiv.org/abs/math/0310337v1) [math.CT].
- [Lod10] J.-L. Loday, *On the operad of associative algebras with derivation*, Georgian Math. J. **17** (2010), no. 2, 347–372.
- [Lyu11] V. Lyubashenko, *Homotopy unital A_∞ -algebras*, J. Algebra **329** (2011), 190–212.
- [Mar96] M. Markl, *Models for operads*, Comm. Algebra **24** (1996), no. 4, 1471–1500.
- [MT14] F. Muro and A. Tonks, *Unital associahedra*, Forum Math. **26** (2014), no. 2, 593–620.
- [Mur11] F. Muro, *Homotopy theory of nonsymmetric operads*, Algebr. Geom. Topol. **11** (2011), 1541–1599.
- [Mur14] ———, *Homotopy units in A -infinity algebras*, [arXiv:1111.2723](https://arxiv.org/abs/1111.2723) [math.AT], to appear in Trans. Amer. Math. Soc., 2014.
- [MV09a] S. Merkulov and B. Vallette, *Deformation theory of representations of prop(erad)s. I*, J. Reine Angew. Math. **634** (2009), 51–106.
- [MV09b] ———, *Deformation theory of representations of prop(erad)s. II*, J. Reine Angew. Math. **636** (2009), 123–174.

UNIVERSIDAD DE SEVILLA
 FACULTAD DE MATEMÁTICAS
 DEPARTAMENTO DE ÁLGEBRA
 AVDA. REINA MERCEDES S/N
 41012 SEVILLA, SPAIN

E-mail address: fmuro@us.es

URL: <http://personal.us.es/fmuro>

