

Perturbation theory and canonical coordinates in celestial mechanics

18th School on Interactions Between Dynamical Systems and Partial Differential
Equations (JISD2022)*

Gabriella Pinzari[†]

May 27, 2022

Abstract

KAM theory owes most of its success to its initial motivation: the application to problems of celestial mechanics. The masterly application was offered by V.I. Arnold in the 60s who worked out a theorem, that he named the “Fundamental Theorem”, especially designed for the planetary problem. This is the problem of $1 + n$ point masses, one “sun” and n “planets”, undergoing gravitational attraction. However, Arnold’s Fundamental Theorem could be really used at that purpose only when, about 50 years later, the “right” canonical set was discovered. In these lectures I shall talk about the complex interplay between perturbation theories and canonical coordinates in problems of celestial mechanics.

Contents

1	Some sets of canonical coordinates for many–body problems	2
1.1	$(1 + n)$ –body problem, Delaunay–Poincaré coordinates and Arnold’s theorem	2
1.2	The rotational degeneracy	6
1.3	Jacobi reduction of the nodes	7
1.4	Deprit coordinates	8
1.5	The map \mathcal{K}	13
1.6	The reduction of perihelia \mathcal{P}	19
1.7	The behavior of \mathcal{K} and \mathcal{P} under reflections	22
2	Applications	23
2.1	Arnold’s Theorem	23
2.2	Global Kolmogorov tori	25
2.3	On the co–existence of stable and whiskered tori	32
A	The m_0–centric reduction	44
	References	45

*I warmly thank M. Guardia and T. M–Seara for their interest and the Centre de Recerca Matemàtica of Bellaterra (Barcelona) for its kind hospitality during the 18th School of Interaction Between Dynamical Systems and Partial Differential Equations, 2022.

Sections 1.5, 1.6, 1.7, 2.2 and 2.3 are based on work done while the author was funded by the ERC grant 677793 StableChaoticPlanetM (2016–2022).

[†]Department of Mathematics, University of Padua, e–mail address: pinzari@math.unipd.it

1 Some sets of canonical coordinates for many–body problems

1.1 $(1+n)$ –body problem, Delaunay–Poincaré coordinates and Arnold’s theorem

In the masterpiece [1], a young a brilliant mathematician, named Vladimir Igorevich Arnold, stated, and partly proved, the following result.

Theorem 1.1 “Theorem of stability of planetary motions”, [1, Chapter III, p. 125]

For the majority of initial conditions under which the instantaneous orbits of the planets are close to circles lying in a single plane, perturbation of the planets on one another produces, in the course of an infinite interval of time, little change on these orbits provided the masses of the planets are sufficiently small. [...] In particular [...] in the n -body problem there exists a set of initial conditions having a positive Lebesgue measure and such that, if the initial positions and velocities of the bodies belong to this set, the distances of the bodies from each other will remain perpetually bounded.

Let us summarize the main ideas behind the statement above.

After the symplectic reduction of the linear momentum, the $(1+n)$ –body problem with masses m_0, m_1, \dots, m_n is governed by the $3n$ –degrees of freedom Hamiltonian (see Appendix A)

$$\mathcal{H} = \sum_{1 \leq i \leq n} \left(\frac{|\mathbf{y}_i|^2}{2\mu_i} - \frac{\mu_i M_i}{|\mathbf{x}_i|} \right) + \sum_{1 \leq i < j \leq n} \left(\frac{\mathbf{y}_i \cdot \mathbf{y}_j}{m_0} - \frac{m_i m_j}{|\mathbf{x}_i - \mathbf{x}_j|} \right) \quad (1)$$

where \mathbf{x}_i represent the difference between the position of the i^{th} planet and the mass m_0 , \mathbf{y}_i are the associated symplectic momenta, $\mathbf{x} \cdot \mathbf{y} = \sum_{1 \leq i \leq 3} x_i y_i$ and $|\mathbf{x}| := (\mathbf{x} \cdot \mathbf{x})^{1/2}$ denote, respectively, the standard inner product in \mathbb{R}^3 and the Euclidean norm;

$$\mu_i := \frac{m_0 m_i}{m_0 + m_i}, \quad M_i := m_0 + m_i \quad (2)$$

The phase space is the “collisionless” domain of $\mathbb{R}^{3n} \times \mathbb{R}^{3n}$

$$\left\{ (\mathbf{y}, \mathbf{x}) = ((\mathbf{y}_1, \dots, \mathbf{y}_n), (\mathbf{x}_1, \dots, \mathbf{x}_n)) \text{ s.t. } 0 \neq \mathbf{x}_i \neq \mathbf{x}_j, \forall i \neq j \right\}, \quad (3)$$

endowed with the standard symplectic form

$$\omega = \sum_{i=1}^n d\mathbf{y}_i \wedge d\mathbf{x}_i = \sum_{i=1}^n \sum_{j=1}^3 dy_{ij} \wedge dx_{ij}$$

where $\mathbf{y}_{ij}, \mathbf{x}_{ij}$ denote the j^{th} component of $\mathbf{y}_i, \mathbf{x}_i$.

The *planetary case* is when m_1, \dots, m_n are of the same order, and much smaller that m_0 . In such a case, letting $m_i \rightarrow \mu m_i, \mathbf{y}_i \rightarrow \mu \mathbf{y}_i$, with $0 < \mu \ll 1$, one obtains

$$\mathcal{H} = \sum_{1 \leq i \leq n} \left(\frac{|\mathbf{y}_i|^2}{2\mu_i} - \frac{\mu_i M_i}{|\mathbf{x}_i|} \right) + \mu \sum_{1 \leq i < j \leq n} \left(\frac{\mathbf{y}_i \cdot \mathbf{y}_j}{m_0} - \frac{m_i m_j}{|\mathbf{x}_i - \mathbf{x}_j|} \right) \quad (4)$$

with

$$\mu_i := \frac{m_0 m_i}{m_0 + \mu m_i}, \quad M_i := m_0 + \mu m_i \quad (5)$$

Consider the *two–body* Hamiltonians

$$h_i(\mathbf{y}_i, \mathbf{x}_i) := \frac{|\mathbf{y}_i|^2}{2\mu_i} - \frac{\mu_i M_i}{|\mathbf{x}_i|}. \quad (6)$$

Assume that $h_i(\mathbf{y}_i, \mathbf{x}_i) < 0$ so that the Hamiltonian flow $\phi_{h_i}^t$ evolves on a Keplerian ellipse \mathcal{E}_i and assume that the eccentricity $e_i \in (0, 1)$. Let a_i, \mathbf{P}_i denote, respectively, the *semimajor axis* and the *perihelion* of \mathcal{E}_i . Let \mathbf{C}_i denote the i^{th} angular momentum

$$\mathbf{C}_i(\mathbf{y}_j, \mathbf{x}_j) := \mathbf{x}_i \times \mathbf{y}_i. \quad (7)$$

Define the *Delaunay nodes*

$$\bar{\mathbf{n}}_i := \mathbf{k} \times \mathbf{C}_i \quad (8)$$

and, for $u, v \in \mathbb{R}^3$ lying in the plane orthogonal to a vector w , let $\alpha_w(u, v)$ denote the positively oriented angle (mod 2π) between u and v (orientation follows the “right hand rule”).

The *Delaunay action-angle variables*

$$\mathcal{D}_{el,aa} := (\mathbf{Z}, \mathbf{G}, \mathbf{\Lambda}, \zeta, \mathbf{g}, \boldsymbol{\ell}) \quad (9)$$

with

$$\begin{aligned} \mathbf{Z} &= (Z_1, \dots, Z_n), & \zeta &= (\zeta_1, \dots, \zeta_n) \\ \mathbf{G} &= (G_1, \dots, G_n), & \mathbf{g} &= (g_1, \dots, g_n) \\ \mathbf{\Lambda} &= (\Lambda_1, \dots, \Lambda_n), & \boldsymbol{\ell} &= (\ell_1, \dots, \ell_n) \end{aligned}$$

are defined as

$$\begin{cases} \Lambda_i := \mu_i \sqrt{M_i a_i} \\ \ell_i := \text{mean anomaly of } \mathbf{x}_i \text{ on } \mathcal{E}_i \end{cases} \quad \begin{cases} G_i := |\mathbf{C}_i| = \Lambda_i \sqrt{1 - e_i^2} \\ g_i := \alpha_{\mathbf{C}_i}(\bar{\mathbf{n}}_i, \mathbf{P}_i) \end{cases} \quad \begin{cases} Z_i := \mathbf{C}_i \cdot \mathbf{k} \\ \zeta_i := \alpha_{\mathbf{k}}(\mathbf{i}, \bar{\mathbf{n}}_i) \end{cases} \quad (10)$$

The *Poincaré variables*

$$\mathcal{P}_{oinc} := ((\boldsymbol{\eta}, \mathbf{p}, \mathbf{\Lambda}), (\boldsymbol{\xi}, \mathbf{q}, \boldsymbol{\lambda}))$$

with

$$\begin{aligned} \boldsymbol{\eta} &= (\eta_1, \dots, \eta_n), & \boldsymbol{\xi} &= (\xi_1, \dots, \xi_n) \\ \mathbf{p} &= (p_1, \dots, p_n), & \mathbf{q} &= (q_1, \dots, q_n) \\ \mathbf{\Lambda} &= (\Lambda_1, \dots, \Lambda_n), & \boldsymbol{\lambda} &= (\lambda_1, \dots, \lambda_n) \end{aligned}$$

with the Λ_i 's as in (10) and

$$\begin{aligned} \lambda_i = \ell_i + G_i + \theta_i & \quad \begin{cases} \eta_i = \sqrt{2(\Lambda_i - G_i)} \cos(\zeta_i + g_i) \\ \xi_i = -\sqrt{2(\Lambda_i - G_i)} \sin(\zeta_i + g_i) \end{cases} \\ & \quad \begin{cases} p_i = \sqrt{2(G_i - Z_i)} \cos \zeta_i \\ q_i = -\sqrt{2(G_i - Z_i)} \sin \zeta_i \end{cases} \end{aligned} \quad (11)$$

In Poincaré coordinates the Hamiltonian (4) takes the form

$$\mathcal{H}_P(\Lambda, \lambda, z) = h_K(\Lambda) + \mu f_P(\Lambda, \lambda, z), \quad z := (\boldsymbol{\eta}, \mathbf{p}, \boldsymbol{\xi}, \mathbf{q}) \in \mathbb{R}^{4n} \quad (12)$$




Figure 1: Delaunay coordinates Z_i, ζ_i, G_i .




Figure 2: Delaunay coordinates G_i, g_i, l_i .

where $(\Lambda, \lambda) \in \mathbb{R}^n \times \mathbb{T}^n$; the ‘‘Kepler’’ unperturbed term h_K , coming from h_{plt} in (1), becomes

$$h_K := \sum_{i=1}^n h_K^{(i)}(\Lambda) = - \sum_{i=1}^n \frac{\mu_i^3 M_i^2}{2\Lambda_i^2}. \quad (13)$$

Because of rotation (with respect the \mathbf{k} -axis) and reflection (with respect to the coordinate planes) invariance of the Hamiltonian (1), the perturbation f_P in (12) satisfies well known symmetry relations called *d’Alembert rules*, see [4]. By such symmetries, in particular, the averaged perturbation

$$f_P^{\text{av}}(\Lambda, z) := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} f_P(\Lambda, \lambda, z) d\lambda \quad (14)$$

is even around the origin $z = 0$ and its expansion in powers of z has the form¹

$$f_P^{\text{av}} = C_0(\Lambda) + \mathcal{Q}_h(\Lambda) \cdot \frac{\eta^2 + \xi^2}{2} + \mathcal{Q}_v(\Lambda) \cdot \frac{p^2 + q^2}{2} + O(|z|^4), \quad (15)$$

where $\mathcal{Q}_h, \mathcal{Q}_v$ are suitable quadratic forms. The explicit expression of such quadratic forms can be found, e.g., in [8, (36), (37)].

By such expansion, the (secular) origin $z = 0$ is an *elliptic equilibrium* for f_P^{av} and corresponds to co-planar and co-circular motions. It is therefore natural to put (15) into Birkhoff Normal Form (BNF, from now on) in a small neighborhood of the secular origin; see, e.g., [10] for general information on BNFs for Birkhoff theory for rotational invariant Hamiltonian systems.

As a preliminary step, one can diagonalize (15), i.e., find a symplectic transformation defined by $\Lambda \rightarrow \Lambda$ and

$$\lambda = \tilde{\lambda} + \varphi(\Lambda, \tilde{z}), \quad \eta = \rho_h(\Lambda)\tilde{\eta}, \quad \xi = \rho_h(\Lambda)\tilde{\xi}, \quad p = \rho_v(\Lambda)\tilde{p}, \quad q = \rho_v(\Lambda)\tilde{q}, \quad (16)$$

with $\rho_h, \rho_v \in \text{SO}(n)$ diagonalizing $\mathcal{Q}_h, \mathcal{Q}_v$. In this way, (12) takes the form

$$\tilde{\mathcal{H}}_P(\Lambda, \tilde{\lambda}, \tilde{z}) = h_K(\Lambda) + \mu \tilde{f}(\Lambda, \tilde{\lambda}, \tilde{z}), \quad (17)$$

with the average over $\tilde{\lambda}$ of \tilde{f}^{av} given by

$$\tilde{f}^{\text{av}}(\Lambda, \tilde{z}) = C_0(\Lambda) + \sum_{i=1}^m \Omega_i(\Lambda) \frac{\tilde{u}_i^2 + \tilde{v}_i^2}{2} + O(|\tilde{z}|^4), \quad \tilde{z} = (\tilde{u}, \tilde{v}) = ((\tilde{\eta}, \tilde{p}), (\tilde{\xi}, \tilde{q})). \quad (18)$$

with $m = 2n$, and the vector $\Omega(\Lambda) := (\sigma_1(\Lambda), \dots, \sigma_n(\Lambda), \varsigma_1(\Lambda), \dots, \varsigma_n(\Lambda))$ being formed by the eigenvalues of the matrices \mathcal{Q}_h and \mathcal{Q}_v .

Theorem 1.2 (Birkhoff) *Let \mathcal{H} be a Hamiltonian having the form in (17)–(18). Assume that there exists $\tilde{\varepsilon} > 0$, $\mathcal{A} \subset \mathbb{R}^n$ and $s \in \mathbb{N}$ such that \mathcal{H} is smooth on an open set $\tilde{\mathcal{M}}_{\tilde{\varepsilon}}^{2m+2n} = \mathcal{A} \times \mathbb{T}^n \times B_{\tilde{\varepsilon}}^{2m}$ and that*

$$\sum_{i=0}^m \Omega_i(\Lambda) k_i \neq 0 \quad \forall k = (k_1, \dots, k_m) \in \mathbb{Z}^m : 0 < |k|_1 \leq 2s, \quad \forall \Lambda \in \mathcal{A}. \quad (19)$$

Then there exists $0 < \varepsilon \leq \tilde{\varepsilon}$ and a symplectic map (‘‘Birkhoff transformation’’)

$$\Phi_B : (\Lambda, \mathbf{l}, \bar{\mathbf{w}}) \in \mathcal{M}_{\varepsilon}^{2m+2n} \rightarrow (\Lambda, \tilde{\lambda}, \tilde{z}) \in \Phi_B(\mathcal{M}_{\varepsilon}^{2m+2n}) \subseteq \mathcal{M}_{\tilde{\varepsilon}}^{2m+2n} \quad (20)$$

which puts the Hamiltonian (17) into the form

$$\mathcal{H}_B(\Lambda, \mathbf{l}, \bar{\mathbf{w}}) := \tilde{\mathcal{H}}_P \circ \Phi_B = h_K(\Lambda) + \mu f_B(\Lambda, l, w) \quad (21)$$

¹ $\mathcal{Q} \cdot u^2$ denotes the 2-indices contraction $\sum_{i,j} \mathcal{Q}_{ij} u_i u_j$ (\mathcal{Q}_{ij}, u_i denoting the entries of \mathcal{Q}, u).

where the average $f_B^{\text{av}}(\Lambda, w) := \int_{\mathbb{T}^n} f_B dl$ is in BNF of order s :

$$f_B^{\text{av}}(\Lambda, w) = C_0 + \Omega \cdot r + P_s(r) + O(|w|^{2s+1}) \quad w := (u, v) \quad r_i := \frac{u_i^2 + v_i^2}{2}, \quad (22)$$

P_s being homogeneous polynomial in r of order s , with coefficients depending on Λ . In particular, if (19) holds with $s = 4$,

$$f_B^{\text{av}}(\Lambda, w) = C_0(\Lambda) + \Omega(\Lambda) \cdot r + r \cdot \tau(\Lambda)r + O(|w|^5) \quad w := (u, v) \quad r_i := \frac{u_i^2 + v_i^2}{2}, \quad (23)$$

with some square matrix $\tau(\Lambda)$ of order m (“torsion”, or “second-order Birkhoff invariants”).

Theorem 1.3 (“The Fundamental Theorem”, V. I. Arnold, [1]) *If the Hessian matrix of h and the matrix $\tau(\Lambda)$ do not vanish identically, and if μ is suitably small with respect to ε , the system affords a positive measure set $\mathcal{K}_{\mu, \varepsilon}$ of quasi-periodic motions in phase space such that its density goes to one as $\varepsilon \rightarrow 0$.*

Remark 1.1 (Arnold, Herman) It turns out that such invariants satisfy identically the following two secular resonances

$$\varsigma_n(\Lambda) \equiv 0, \quad \sum_{i=1}^n (\sigma_i(\Lambda) + \varsigma_i(\Lambda)) \equiv 0 \quad (24)$$

Such resonances strongly violate the assumption (19) of Theorem 1.2.

We remark that the former equality in (24) is mentioned in [1], while the latter been pointed out by M. Herman in the 1990s. Note that (24) do not appear in the planar problem, because the matrix \mathcal{Q}_v , hence the ς_i ’s, do not exist in that case. Being aware of such difficulty, Arnold completely proved Theorem 1.1 via Theorem 2.2 in the case of the planar three-body problem, checking explicitly the non vanishing of the 2×2 torsion matrix for that case. However, in the case of the spatial problem, the question remained open until 2004, when M. Herman and J. Féjóz [8] proved Theorem 1.1 via a completely different strategy, which does need Birkhoff normal form. We refer to [6] for more details.

1.2 The rotational degeneracy

In [1], Arnold wrote – without giving the details – that the former resonance in (24) was to be ascribed to the conservation of the total angular momentum of the system:

$$\mathbf{C} = \sum_{j=1}^n \mathbf{C}_j, \quad \mathbf{C}_j = \mathbf{x}_j \times \mathbf{y}_j. \quad (25)$$

An argument which clearly shows this goes as follows. Using Poincaré coordinates, the planets’ angular momenta have the expressions

$$\begin{aligned} \mathbf{C}_j &= \begin{pmatrix} -q_j \sqrt{\Lambda_j - \frac{\eta_j^2 + \xi_j^2}{2} - \frac{p_j^2 + q_j^2}{4}} \\ -p_j \sqrt{\Lambda_j - \frac{\eta_j^2 + \xi_j^2}{2} - \frac{p_j^2 + q_j^2}{4}} \\ \Lambda_j - \frac{\eta_j^2 + \xi_j^2}{2} - \frac{p_j^2 + q_j^2}{2} \end{pmatrix} \\ &= \begin{pmatrix} -\sqrt{\Lambda_j} q_j + O(|z|^3) \\ -\sqrt{\Lambda_j} p_j + O(|z|^3) \\ \Lambda_j + O(|z|^2) \end{pmatrix} \end{aligned}$$

In particular, the two former components of the total angular momentum (25) are given by

$$C_1 = -\sum_{j=1}^n \sqrt{\Lambda_j} q_j + O(|z|^3), \quad C_2 = -\sum_{j=1}^n \sqrt{\Lambda_j} p_j + O(|z|^3) \quad (26)$$

On the other hand, it is possible to find a canonical transformation

$$(\Lambda, \check{\lambda}, \check{\eta}, \check{p}, \check{\xi}, \check{q}) \rightarrow (\Lambda, \lambda, \eta, p, \xi, q) \quad (27)$$

having the form (16) with $\rho_h = \text{id}$ and $\rho_v \in SO(n)$ chosen such in a way that the last row of ρ_v^{-1} is

$$N(\Lambda)(\sqrt{\Lambda_1}, \dots, \sqrt{\Lambda_n}) \quad (28)$$

where $N(\Lambda) = \frac{1}{\sqrt{\sum_{i=1}^n \Lambda_i}}$ fixes the Euclidean norm of (28) to 1. With such choice, we have

$$\check{p}_n = \rho_v^{-1} \begin{pmatrix} p_1 \\ \vdots \\ p_n \end{pmatrix}_n = N(\Lambda) \sum_{j=1}^n \sqrt{\Lambda_j} p_j$$

and, similarly,

$$\check{q}_n = N(\Lambda) \sum_{j=1}^n \sqrt{\Lambda_j} q_j$$

Therefore, (122) become

$$C_1 = -N(\Lambda)^{-1} \check{q}_n + O(|\check{z}|^3), \quad C_2 = -N(\Lambda)^{-1} \check{p}_n + O(|\check{z}|^3) \quad (29)$$

Now, as the projection of the transformation (27) on $\check{\lambda}$'s is a $\check{\lambda}$ -independent translation, the averaged perturbing function using the new coordinates can be obtained applying such transformation to the function in (15). We denote it as

$$\check{f}^{\text{av}} = C_0(\Lambda) + \check{Q}_h(\Lambda) \cdot \frac{\check{\eta}^2 + \check{\xi}^2}{2} + \check{Q}_v(\Lambda) \cdot \frac{\check{p}^2 + \check{q}^2}{2} + O(|\check{z}|^4),$$

with $\check{Q}_h(\Lambda) = Q_h(\Lambda)$ and $\check{Q}_v(\Lambda) = \rho_v(\Lambda)^{-1} Q_v(\Lambda) \rho_v(\Lambda)$. Note that $\check{Q}_v(\Lambda)$ has the same eigenvalues as $Q_v(\Lambda)$, as $\rho_v \in SO(n)$. Let us now use

$$\{\check{f}^{\text{av}}, C_1\} = 0 = \{\check{f}^{\text{av}}, C_2\} \quad (30)$$

which hold because they are true for f , and \mathbf{C} is $\check{\lambda}$ -independent. Using (29), it is immediate to see that (30) imply that the quadratic form

$$\check{Q}_v(\Lambda) \cdot \frac{\check{p}^2 + \check{q}^2}{2}$$

is independent of \check{p}_n, \check{q}_n . Hence, the n^{th} row and column of $\check{Q}_v(\Lambda)$ vanish identically. This implies that $\check{Q}_v(\Lambda)$, hence $Q_v(\Lambda)$, has an identically vanishing eigenvalue, which is $\varsigma_n(\Lambda)$ in (24).

1.3 Jacobi reduction of the nodes

In the case $n = 2$, Arnold in [1] suggested to get rid of the rotation invariance (described in the previous section) by means of the classical so-called *Jacobi reduction of the nodes*. This is a classical procedure with a remarkable geometric meaning, which goes as follows. Let us consider

a reference frame $(\mathbf{i}, \mathbf{j}, \mathbf{k})$ whose third axis \mathbf{k} is along the direction of the total angular momentum $\mathbf{C} = \mathbf{C}_1 + \mathbf{C}_2$, while \mathbf{i} coincides with the intersection of the planes orthogonal to $\mathbf{C}_1, \mathbf{C}_2$. Such intersection is well defined provided that $\mathbf{C}_1 \not\parallel \mathbf{C}_2$, namely, when the problem is not planar. With such a choice of the reference frame, one cannot fix Delaunay coordinates completely freely. Indeed, by the choice of \mathbf{i} , we have that the ζ_j satisfy

$$\zeta_2 - \zeta_1 = \pi. \quad (31)$$

Moreover, a geometrical analysis of the triangle formed by $\mathbf{C}_1, \mathbf{C}_2$ and \mathbf{C} shows that the coordinates Z_j satisfy




Figure 3: The construction underlying Jacobi reduction of the nodes.

$$Z_1 = \frac{G}{2} + \frac{G_1^2 - G_2^2}{2G}, \quad Z_2 = \frac{G}{2} - \frac{G_1^2 - G_2^2}{2G} \quad (32)$$

where $G := |\mathbf{C}| = \sqrt{C_1^2 + C_2^2 + C_3^2}$ is the Euclidean norm of \mathbf{C} . As \mathbf{i} moves, the following fact is not obvious at all – in fact proved by R. Radau.

Theorem 1.4 (R. Radau, 1868, [15]) *Replacing relations (31)–(32) inside the Hamiltonian (1) with $n = 2$ written in Delaunay coordinates, one obtains a function, depending on $(\Lambda_j, \ell_j, G_j, g_j)$ ($j = 1, 2$) and G , whose Hamilton equations relatively to $(\Lambda_j, \ell_j, G_j, g_j)$ generate the motions of the coordinates $(\Lambda_j, \ell_j, G_j, g_j)$ referred to the rotating frame under the action of the Hamiltonian (1) with $n = 2$. The motion of Z_j and ζ_j can be recovered via (31)–(32).*

1.4 Deprit coordinates

Arnold commented on the general problem of rotational degeneracy as follows:

[1, Chap.III, §5, n. 5] *In the case of more than three bodies [$n > 2$] there is no such [analogue to Jacobi reduction of the nodes] elegant method of reducing the number of degrees of freedom [...].*

However, exactly 20 years later, in 1983, A. Deprit [7] discovered a set of canonical coordinates which, after a simple transformation, do the desired job and reduce to Jacobi’s when $n = 2$. Let us describe them.

Consider the “partial angular momenta”

$$\mathbf{S}_j(\mathbf{y}, \mathbf{x}) := \sum_{i=1}^j \mathbf{C}_i; \quad (33)$$

with \mathbf{C}_i as in (7). Notice that $\mathbf{S}_n = \mathbf{C}$ is the total angular momentum of the system. Define the ‘‘Deprit nodes’’

$$\begin{cases} \boldsymbol{\nu}_{i+1} := \mathbf{S}_{i+1} \times \mathbf{C}_{i+1}, & 1 \leq i \leq n-1 \\ \boldsymbol{\nu}_1 := \mathbf{S}_2 \times \mathbf{C}_1 = -\boldsymbol{\nu}_2 \\ \boldsymbol{\nu}_{n+1} := \mathbf{k} \times \mathbf{C} =: \bar{\boldsymbol{\nu}}. \end{cases} \quad (34)$$

If $n \geq 2$, Deprit’s coordinates

$$\mathcal{D}_{ep} = (\mathbf{R}, \mathbf{G}, \boldsymbol{\Psi}, \mathbf{r}, \boldsymbol{\varphi}, \psi) \quad (35)$$

with

$$\begin{aligned} \mathbf{R} &= (R_1, \dots, R_n), \quad \boldsymbol{\Psi} = (\Psi_1, \dots, \Psi_n), \quad \mathbf{G} = (G_1, \dots, G_n), \\ \mathbf{r} &= (r_1, \dots, r_n), \quad \boldsymbol{\psi} = (\psi_1, \dots, \psi_n), \quad \boldsymbol{\varphi} = (\varphi_1, \dots, \varphi_n). \end{aligned} \quad (36)$$

are defined as follows (compare also Figures 4, 5 and 6):

$$\begin{cases} R_i := \mathbf{y}_i \cdot \frac{\mathbf{x}_i}{|\mathbf{x}_i|} \\ r_i := |\mathbf{x}_i| \end{cases} \quad \begin{cases} G_i := |\mathbf{C}_i| \\ \varphi_i := \alpha_{\mathbf{C}_i}(\boldsymbol{\nu}_i, \mathbf{x}_i) \end{cases} \quad (37)$$

$$\Psi_i := \begin{cases} |\mathbf{S}_{i+1}| & 1 \leq i \leq n-2 \quad (n \geq 3) \\ C := |\mathbf{C}| & i = n-1 \\ Z := \mathbf{C} \cdot \mathbf{k} & i = n \end{cases}$$

$$\psi_i := \begin{cases} \alpha_{\mathbf{S}_{i+1}}(\boldsymbol{\nu}_{i+2}, \boldsymbol{\nu}_{i+1}) & 1 \leq i \leq n-2 \quad (n \geq 3) \\ \gamma := \alpha_{\mathbf{C}}(\bar{\boldsymbol{\nu}}, \boldsymbol{\nu}_n) & i = n-1 \\ \zeta := \alpha_{\mathbf{k}}(\mathbf{i}, \bar{\boldsymbol{\nu}}) & i = n \end{cases}$$

We have

Theorem 1.5 (A. Deprit, 1983, [7]) $\sum_{i=1}^n \mathbf{y}_i \cdot d\mathbf{x}_i = \mathbf{R} \cdot d\mathbf{r} + \boldsymbol{\Psi} \cdot d\boldsymbol{\psi} + \mathbf{G} \cdot d\boldsymbol{\varphi}$ for all $n \in \mathbb{N}$.

For later need, we formulate an equivalent statement of Theorem 1.5. We consider the coordinates

$$\mathcal{D}_{el} := (\mathbf{Z}, \mathbf{G}, \mathbf{R}, \boldsymbol{\zeta}, \boldsymbol{\phi}, \mathbf{r}) \quad (38)$$

with

$$\begin{aligned} \mathbf{Z} &= (Z_1, \dots, Z_n), \quad \mathbf{G} = (G_1, \dots, G_n), \quad \mathbf{R} = (R_1, \dots, R_n) \\ \boldsymbol{\zeta} &= (\zeta_1, \dots, \zeta_n), \quad \boldsymbol{\phi} = (\phi_1, \dots, \phi_n), \quad \mathbf{r} = (r_1, \dots, r_n) \end{aligned} \quad (39)$$

where Z_i, G_i, ζ_i , are as in (10), R_i, r_i are as in (37), and, finally,

$$\phi_i := \alpha_{\mathbf{C}_i}(\mathbf{n}_i, \mathbf{x}_i).$$

Let

$$\mathcal{R}_1(i) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos i & -\sin i \\ 0 & \sin i & \cos i \end{pmatrix}, \quad \mathcal{R}_3(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (40)$$




Figure 4: Deprit coordinates Z , C and ζ fix the angular momentum in the initial reference frame $(\mathbf{i}, \mathbf{j}, \mathbf{k})$.




Figure 5: Deprit coordinates Ψ_i , Ψ_{i-1} , G_{i+1} and ψ_i .




Figure 6: Deprit coordinates g_i , G_i , l_i .

and

$$\mathbf{x} = \mathcal{R}_3(\theta)\mathcal{R}_1(i)\bar{\mathbf{x}}, \quad \mathbf{y} = \mathcal{R}_3(\theta)\mathcal{R}_1(i)\bar{\mathbf{y}}, \quad \mathbf{C} := \mathbf{x} \times \mathbf{y}, \quad \bar{\mathbf{C}} := \bar{\mathbf{x}} \times \bar{\mathbf{y}}, \quad \mathbf{i} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{k} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

with $\mathbf{x}, \bar{\mathbf{x}}, \mathbf{y}, \bar{\mathbf{y}} \in \mathbb{R}^3$. The proof of the following fact is left to the reader.

Lemma 1.1 $\mathbf{y} \cdot d\mathbf{x} = \mathbf{C} \cdot \mathbf{k}d\theta + \bar{\mathbf{C}} \cdot \mathbf{i}di + \bar{\mathbf{y}} \cdot d\bar{\mathbf{x}}$.

Lemma 1.1 immediately implies

Lemma 1.2 $\mathbf{y}_j \cdot d\mathbf{x}_j = Z_j d\zeta_j + G_j d\phi_j + R_j dr_j \quad \forall j = 1, \dots, n, \quad \forall n \in \mathbb{N}$.

Indeed, we have

$$\begin{cases} \mathbf{x}_j = \mathcal{R}_3(\zeta_j)\mathcal{R}_1(i_j^*)\mathbf{x}_{j\text{pl}}^* \\ \mathbf{y}_j = \mathcal{R}_3(\zeta_j)\mathcal{R}_1(i_j^*)\mathbf{y}_{j\text{pl}}^* \end{cases} \quad j = 1, \dots, n$$

where i_j^* is the convex angle formed by \mathbf{k} and \mathbf{C}_j and, finally,

$$\mathbf{x}_{j\text{pl}}^* = \begin{pmatrix} r_j \cos \phi_j \\ r_j \sin \phi_j \\ 0 \end{pmatrix}, \quad \mathbf{y}_{j\text{pl}}^* = \begin{pmatrix} R_j \cos \varphi_j - \frac{G_j}{r_j} \sin \phi_j \\ R_j \sin \varphi_j + \frac{G_j}{r_j} \cos \phi_j \\ 0 \end{pmatrix} \quad (41)$$

verify, as well known,

$$\mathbf{y}_{j\text{pl}}^* \cdot d\mathbf{x}_{j\text{pl}}^* = R_j dr_j + G_j d\phi_j. \quad (42)$$

Then, by Lemma 1.1, (42) and as $\mathbf{C}_{j\text{pl}}^* \cdot \mathbf{i} = 0$, we have

$$\begin{aligned} \mathbf{y}_j \cdot d\mathbf{x}_j &= \mathbf{C}_j \cdot \mathbf{k}d\zeta_j + \mathbf{C}_{j\text{pl}}^* \cdot \mathbf{i}di_j + \mathbf{y}_{j\text{pl}}^* \cdot d\mathbf{x}_{j\text{pl}}^* \\ &= Z_j d\zeta_j + G_j d\phi_j + R_j dr_j. \quad \square \end{aligned}$$

We denote as

$$\phi_{\mathcal{D}_{el}}^{\mathcal{D}_{ep}} : \mathcal{D}_{el} = (\mathbf{Z}, \mathbf{G}, \mathbf{R}, \zeta, \phi, \mathbf{r}) \rightarrow \mathcal{D}_{ep} = (\Psi, \mathbf{G}, \mathbf{R}, \psi, \varphi, \mathbf{r})$$

the map which relates \mathcal{D}_{el} and \mathcal{D}_{ep} and as

$$\widehat{\phi}_{\mathcal{D}_{el}}^{\mathcal{D}_{ep}} : \widehat{\mathcal{D}}_{el} = (\mathbf{Z}, \mathbf{G}, \zeta, \phi) \rightarrow \widehat{\mathcal{D}}_{ep} = (\Psi, \mathbf{G}, \psi, \varphi)$$

is the natural projections on the coordinates above. It is easy to check that $\widehat{\phi}_{\mathcal{D}_{el}}^{\mathcal{D}_{ep}}$ is independent of \mathbf{R} and \mathbf{r} . Indeed, $\widehat{\phi}_{\mathcal{D}_{el}}^{\mathcal{D}_{ep}}$ has the expression

$$\begin{aligned} G_j &= G_j, \\ \varphi_j &= \phi_j + \alpha_{\mathbf{C}_i}(\boldsymbol{\nu}_j, \bar{\boldsymbol{\nu}}_j) \text{ with } \bar{\boldsymbol{\nu}}_j = \mathbf{k} \times \mathbf{C}_j, \\ \Psi_j &= \begin{cases} |\mathbf{S}_{j+1}| & j \neq n \\ Z_1 + \dots + Z_n & j = n \end{cases} \\ \psi_j &= \begin{cases} \alpha_{\mathbf{S}_{j+1}}(\boldsymbol{\nu}_{j+2}, \boldsymbol{\nu}_{j+1}) & j \neq n \\ \alpha_{\mathbf{k}}(\mathbf{i}, \bar{\boldsymbol{\nu}}) & j = n \end{cases} \end{aligned} \quad (43)$$

where the right hand sides are to be written as functions of \mathcal{D}_{el} . As the right hand sides are defined only in terms of \mathbf{C}_j , so they are functions of \mathbf{Z}, ζ and \mathbf{G} , while are independent of \mathbf{R} and \mathbf{r} .

Theorem 1.6 *Theorem 1.5 is equivalent to stress that*

$$\widehat{\phi}_{\mathcal{D}_{el}}^{\mathcal{D}_{ep}} \text{ verifies : } \mathbf{Z} \cdot d\zeta + \mathbf{G} \cdot d\phi = \Psi \cdot d\psi + \mathbf{G} \cdot d\varphi \text{ for all } n \in \mathbb{N}. \quad (44)$$

Proof Use Lemma 1.2 and that the coordinates (\mathbf{R}, \mathbf{r}) are shared by \mathcal{D}_{ep} and \mathcal{D}_{el} . \square

We prove Theorem 1.5 (\iff (44)) by induction on n , with $n \geq 2$, as in [11].

Base step We prove the statement 1.5 with $n = 2$. We first observe that, in such case, $(\mathbf{y}_j, \mathbf{x}_j)$ are expressed, through $(\mathbf{R}, \Psi, \mathbf{G}, \mathbf{r}, \psi, \varphi)$ via the formulae

$$\begin{cases} \mathbf{x}_j = \mathcal{R}_3(\zeta)\mathcal{R}_1(i)\mathcal{R}_3(\gamma)\mathcal{R}_1(i_j)\mathbf{x}_{j\text{pl}} \\ \mathbf{y}_j = \mathcal{R}_3(\zeta)\mathcal{R}_1(i)\mathcal{R}_3(\gamma)\mathcal{R}_1(i_j)\mathbf{y}_{j\text{pl}} \end{cases} \quad j = 1, 2$$

where i is the convex² angle formed by \mathbf{k} and \mathbf{C} ; i_j is the convex angle formed by \mathbf{C} and \mathbf{C}_j and, finally, $\mathbf{x}_{j\text{pl}}, \mathbf{y}_{j\text{pl}}$ are as in (41), with ϕ_j replaced by φ_j .

Using Lemma 1.1 twice, one easily finds

$$\begin{aligned} \mathbf{y}_j \cdot d\mathbf{x}_j &= \mathbf{C}_j \cdot \mathbf{k} d\zeta + \bar{\mathbf{C}}_j \cdot \mathbf{i} di + \bar{\mathbf{C}}_j \cdot \mathbf{k} d\gamma + \mathbf{C}_{j\text{pl}} \cdot \mathbf{i} d(i_j) \\ &\quad + \mathbf{y}_{j\text{pl}} \cdot d\mathbf{x}_{j\text{pl}} \\ &= \mathbf{C}_j \cdot \mathbf{k} d\zeta + \mathbf{C}_j \cdot \mathbf{e}_1 di + \mathbf{C}_j \cdot \mathbf{e}_3 d\gamma + \mathbf{y}_{j\text{pl}} \cdot d\mathbf{x}_{j\text{pl}} \end{aligned} \quad (45)$$

We have used $\mathbf{C}_{j\text{pl}} \cdot \mathbf{i} = 0$, $\mathbf{C}_j = \mathcal{R}_3(\zeta)\mathcal{R}_1(i)\bar{\mathbf{C}}_j$ and we have let

$$\mathbf{e}_1 := \mathcal{R}_3(\zeta)\mathcal{R}_1(i)\mathbf{i}, \quad \mathbf{e}_3 := \mathcal{R}_3(\zeta)\mathcal{R}_1(i)\mathbf{k}. \quad (46)$$

Taking the sum of (45) with $j = 1, 2$ and using (42) and recognizing that

$$\begin{cases} (\mathbf{C}_1 + \mathbf{C}_2) \cdot \mathbf{k} = \mathbf{C} \cdot \mathbf{k} = Z \\ (\mathbf{C}_1 + \mathbf{C}_2) \cdot \mathbf{e}_1 = \mathbf{C} \cdot \mathbf{e}_1 = 0 \\ (\mathbf{C}_1 + \mathbf{C}_2) \cdot \mathbf{e}_3 = \mathbf{C} \cdot \mathbf{e}_3 = C \end{cases}$$

we have the proof. \blacksquare

Induction The inductive step is made on the statement (44). The map $\widehat{\phi}_{\mathcal{D}_{el}}^{\mathcal{D}_{ep}}$ in (44) will be named $\widehat{\phi}_n$. We assume that (44) holds for a given $n \geq 2$ and prove it for $n + 1$. Consider the map

$$\phi_{n+1}^* : \widehat{\mathcal{D}}_{el, n+1} = (\mathbf{Z}, \mathbf{G}, \zeta, \phi) \rightarrow \widetilde{\mathcal{D}}_{ep, n+1} = (\Psi^*, \mathbf{G}^*, \psi^*, \varphi^*)$$

defined as follows. If

$$\mathbf{Z} = (\widetilde{\mathbf{Z}}, Z_{n+1}), \quad \mathbf{G} = (\widetilde{\mathbf{G}}, G_{n+1}), \quad \zeta = (\widetilde{\zeta}, \zeta_{n+1}), \quad \phi = (\widetilde{\phi}, \phi_{n+1})$$

where the tilded arguments have dimension $n + 1$, we let

$$(\widetilde{\Psi}, \widetilde{\mathbf{G}}, \widetilde{\psi}, \widetilde{\varphi}) = \phi_n(\widetilde{\mathbf{Z}}, \widetilde{\mathbf{G}}, \widetilde{\zeta}, \widetilde{\phi})$$

and then

$$\phi_{n+1}^*(\mathbf{Z}, \mathbf{G}, \zeta, \phi) := ((\widetilde{\Psi}, Z_{n+1}), (\widetilde{\mathbf{G}}, G_{n+1}), (\widetilde{\psi}, \zeta_{n+1}), (\widetilde{\varphi}, \phi_{n+1})) =: (\Psi^*, \mathbf{G}^*, \psi^*, \varphi^*)$$

²The expressions of i_1, i_2 and i – not needed here – can easily be deduced by the analysis of the triangle formed by $\mathbf{C}_1, \mathbf{C}_2$ and \mathbf{C} : see Figure 5

By the inductive assumption, ϕ_n verifies

$$\tilde{\mathbf{Z}} \cdot d\tilde{\zeta} + \tilde{\mathbf{G}} \cdot d\tilde{\phi} = \tilde{\Psi} \cdot d\tilde{\psi} + \tilde{\mathbf{G}} \cdot d\tilde{\varphi}$$

and hence ϕ_{n+1}^* verifies

$$\begin{aligned} \mathbf{Z} \cdot d\zeta + \mathbf{G} \cdot d\phi &= \Psi^* \cdot d\psi^* + \mathbf{G}^* \cdot d\varphi^* \\ &= \tilde{\Psi} \cdot d\tilde{\psi} + \tilde{\mathbf{G}} \cdot d\tilde{\varphi} + Z_{n+1}d\zeta_{n+1} + G_{n+1}d\phi_{n+1} \\ &= \left(\sum_{j=1}^{n-2} \tilde{\Psi}_j \cdot d\tilde{\psi}_j + \tilde{\mathbf{G}} \cdot d\tilde{\varphi} \right) + \tilde{\Psi}_{n-1} \cdot d\tilde{\psi}_{n-1} + \tilde{\Psi}_n \cdot d\tilde{\psi}_n + Z_{n+1}d\zeta_{n+1} + G_{n+1}d\phi_{n+1} \end{aligned} \quad (47)$$

having split

$$\tilde{\Psi} = (\tilde{\Psi}_1, \dots, \tilde{\Psi}_n), \quad \tilde{\psi} = (\tilde{\psi}_1, \dots, \tilde{\psi}_n).$$

We moreover define a map $\phi_{*,n+1}$ on $(\Psi^*, \mathbf{G}^*, \psi^*, \varphi^*)$ acting as

$$(\Psi_*, \mathbf{G}_*, \psi_*, \varphi_*) = \phi_2((\tilde{\Psi}_n, Z_{n+1}), (\tilde{\Psi}_{n-1}, G_{n+1}), (\tilde{\psi}_n, \zeta_{n+1}), (\tilde{\psi}_{n-1}, \phi_{n+1}))$$

on the designed variables, and as the identity on the remaining ones. Note that the arguments at left hand side have dimension 2, that $\mathbf{G}_* = (\tilde{\Psi}_{n-1}, G_{n+1})$, and put $\varphi_* = (\varphi_{*,1}, \varphi_{*,2})$. Again by the inductive assumption, we have

$$\tilde{\Psi}_{n-1} \cdot d\tilde{\psi}_{n-1} + \tilde{\Psi}_n \cdot d\tilde{\psi}_n + Z_{n+1}d\zeta_{n+1} + G_{n+1}d\phi_{n+1} = \Psi_* \cdot d\psi_* + \mathbf{G}_* \cdot d\varphi_* \quad (48)$$

Let us now look at the composition

$$\phi_{*,n+1} \circ \phi_{n+1}^* \quad (49)$$

It acts as

$$\begin{aligned} (\mathbf{Z}, \mathbf{G}, \zeta, \phi) &\rightarrow ((\tilde{\Psi}_1, \dots, \tilde{\Psi}_{n-2}, \tilde{\Psi}_{n-1}, \Psi_*), (\tilde{\mathbf{G}}, G_{n+1}), (\tilde{\psi}_1, \dots, \tilde{\psi}_{n-2}, \varphi_{*,1}, \psi_*), (\tilde{\varphi}, \varphi_{*,2})) \\ &=: (\Psi, \mathbf{G}, \psi, \varphi) \end{aligned}$$

and, by (47) and (48), verifies

$$\begin{aligned} \mathbf{Z} \cdot d\zeta + \mathbf{G} \cdot d\phi &= \left(\sum_{j=1}^{n-2} \tilde{\Psi}_j \cdot d\tilde{\psi}_j + \tilde{\mathbf{G}} \cdot d\tilde{\varphi} \right) + \Psi_* \cdot d\psi_* + \mathbf{G}_* \cdot d\varphi_* \\ &= \Psi \cdot d\psi + \mathbf{G} \cdot d\varphi. \end{aligned}$$

It is not difficult to recognize – using (43) – that the map (49) coincides with ϕ_{n+1} . For the details, we refer to [11, 3]. \square

1.5 The map \mathcal{K}

The \mathcal{K} -coordinates have been described in [12] for $n = 2$ and generalized to any $n \in \mathbb{N}$, $n \geq 2$ in [14]. Here, for sake of uniformity with the coordinates \mathcal{D}_{ep} , we change³ notations a little bit compared to [14]. We let

$$\mathcal{K} = (\hat{\Theta}, \hat{\chi}, \mathbf{R}, \hat{\vartheta}, \hat{\kappa}, \mathbf{r})$$

³The main changes regard the coordinates that in [14] are called $\tilde{\Theta}_0, \tilde{\chi}_{n-1}$, which here are called $\hat{\chi}_n, \hat{\Theta}_1$. The other coordinates just underwent a different numbering: $(\tilde{\Theta}_j)_{1 \leq j \leq n-1}, \tilde{\chi}_0, (\tilde{\chi}_j)_{1 \leq j \leq n-2}, \Lambda_j$ here are denoted, respectively, as $(\hat{\Theta}_{n-j+1}), \hat{\chi}_{n-1}, (\hat{\chi}_{n-j+1}), \hat{\Lambda}_{n-j+1}$. An analogue change of notations holds of course for the conjugated coordinates.




Figure 7: The reference frames \hat{F}_j and the \mathcal{K} -coordinates $\hat{\kappa}_{j-1}$, r_j , R_j $j = 2, \dots, n$.




Figure 8: The reference \hat{F}_1 and the \mathcal{K} -coordinates $\hat{\Theta}_1$, $\hat{\vartheta}_1$.




Figure 9: The reference frames \hat{G}_j and the \mathcal{K} -coordinates $\hat{\vartheta}_j$, $\hat{\Theta}_j$, $\hat{\chi}_{j-1}$, $j = 1, \dots, n$. When $j = 2$, take $\hat{\chi}_0 := \hat{\Theta}_1$; when $j = 1$, disregard \mathbf{S}_0 , $\hat{\nu}_0$ and $\hat{\chi}_{-1}$.

where \mathbf{R}, \mathbf{r} are as in (37), while

$$\begin{aligned}\hat{\Theta} &= (\hat{\Theta}_1, \dots, \hat{\Theta}_n), & \hat{\vartheta} &= (\hat{\vartheta}_1, \dots, \hat{\vartheta}_n) \\ \hat{\chi} &= (\hat{\chi}_1, \dots, \hat{\chi}_n), & \hat{\kappa} &= (\hat{\kappa}_1, \dots, \hat{\kappa}_n)\end{aligned}$$

are defined as follows. Let \mathbf{S}_j be as in (33). Define the \mathcal{K} -nodes

$$\hat{\nu}_j := \begin{cases} \mathbf{k} \times \mathbf{C} & j = n \\ \mathbf{x}_{j+1} \times \mathbf{S}_j & j = 1, \dots, n-1 \end{cases} \quad \hat{\mathbf{n}}_j := \mathbf{S}_j \times \mathbf{x}_j \quad j = 1, \dots, n. \quad (50)$$

and then the \mathcal{K} coordinates as follows.

$$\begin{aligned}\hat{\Theta}_j &:= \begin{cases} \mathbf{S}_j \cdot \frac{\mathbf{x}_j}{|\mathbf{x}_j|} \\ |\mathbf{C}_1| \end{cases} & \hat{\vartheta}_j &:= \begin{cases} \alpha_{\mathbf{x}_j}(\hat{\mathbf{n}}_j, \hat{\nu}_{j-1}) & 2 \leq j \leq n \\ \alpha_{\mathbf{C}_1}(\hat{\nu}_1, \hat{\mathbf{n}}_1) & j = 1 \end{cases} \\ \hat{\chi}_j &:= \begin{cases} Z := \mathbf{C} \cdot \mathbf{k} \\ C := |\mathbf{C}| \\ |\mathbf{S}_{j+1}| \end{cases} & \hat{\kappa}_j &:= \begin{cases} \zeta := \alpha_{\mathbf{k}}(\mathbf{i}, \hat{\nu}_n) & j = n \\ \gamma := \alpha_{\mathbf{S}_n}(\hat{\nu}_n, \hat{\mathbf{n}}_n) & j = n-1 \\ \alpha_{\mathbf{S}_{j+1}}(\hat{\nu}_{j+1}, \hat{\mathbf{n}}_{j+1}) & 1 \leq j \leq n-2 \quad (n \geq 3) \end{cases} \end{aligned} \quad (51)$$

Remark 1.2 Note that the node $\hat{\nu}_n$ coincides with $\bar{\nu} = \nu_{n+1}$ in (34); the coordinates Z and ζ are the same as in (37) and, finally, the coordinates χ coincide with the coordinates Ψ in (37). In particular, \mathcal{D}_{ep} and \mathcal{K} share the construction in Figure 4. The geometrical meaning of the other \mathcal{K} -coordinates is pointed out in the next section.

A chain of reference frames We consider the following chain of vectors

$$\begin{array}{ccccccccccc} \mathbf{k} & \rightarrow & \mathbf{S}_n = \mathbf{C} & \rightarrow & \mathbf{x}_n & \rightarrow & \cdots & \rightarrow & \mathbf{S}_j & \rightarrow & \mathbf{x}_j & \rightarrow & \mathbf{S}_{j-1} & \rightarrow & \cdots & \rightarrow & \mathbf{S}_1 = \mathbf{C}_1 \\ & & \Downarrow & & \Downarrow & & \vdots & & \Downarrow & & \Downarrow & & \Downarrow & & \vdots & & \Downarrow \\ & & \hat{\nu}_n & & \hat{\mathbf{n}}_1 & & \vdots & & \hat{\nu}_j & & \hat{\mathbf{n}}_j & & \hat{\nu}_{j-1} & & \vdots & & \hat{\nu}_1 \end{array} \quad (52)$$

where $\hat{\nu}_j, \hat{\mathbf{n}}_j$ are the \mathcal{K} -nodes in (50), given by the skew-product of the two consecutive vectors in the chain.

We associate to this chain of vectors the following chain of frames

$$\hat{\mathbf{G}}_{n+1} \rightarrow \hat{\mathbf{F}}_n \rightarrow \hat{\mathbf{G}}_n \rightarrow \cdots \rightarrow \hat{\mathbf{F}}_j \rightarrow \hat{\mathbf{G}}_j \rightarrow \hat{\mathbf{F}}_{j-1} \rightarrow \cdots \rightarrow \hat{\mathbf{G}}_1 \quad (53)$$

where $\hat{\mathbf{G}}_{n+1} = (\mathbf{i}, \mathbf{j}, \mathbf{k})$ is the initial prefixed frame and the frames, while $\hat{\mathbf{F}}_j, \hat{\mathbf{G}}_j$ are frames defined via

$$\hat{\mathbf{F}}_j = (\hat{\nu}_j, \cdot, \mathbf{S}_j) \quad \hat{\mathbf{G}}_j = (\hat{\mathbf{n}}_j, \cdot, \mathbf{x}_j) \quad j = 1, \dots, n. \quad (54)$$

By construction, each frame in the chain has its first axis coinciding with the intersection of horizontal plane with the horizontal plane of the previous frame (hence, in particular, $\hat{\nu}_j \perp \mathbf{S}_j$ and $\hat{\mathbf{n}}_j \perp \mathbf{x}_j$).

Explicit expression of the \mathcal{K} -map We now derive the explicit formulae of the map which relates the coordinates (51) to the coordinates $(\mathbf{y}_1, \dots, \mathbf{y}_n, \mathbf{x}_1, \dots, \mathbf{x}_n)$. We shall prove that such map has the expression

$$\begin{cases} \mathbf{x}_j = \mathbf{x}_j^n := \mathcal{R}_j^n \tilde{\mathbf{x}}_j \\ \mathbf{y}_j = \mathbf{y}_j^n := \mathcal{R}_j^n \tilde{\mathbf{y}}_j \end{cases} \quad (55)$$

where

$$\begin{cases} \mathcal{R}_j^n := \hat{\mathcal{T}}_n \hat{\mathcal{S}}_n \cdots \hat{\mathcal{T}}_{j+1} \hat{\mathcal{S}}_{j+1} \hat{\mathcal{T}}_j \hat{\mathcal{S}}_j \\ \tilde{\mathbf{x}}_j := r_j \mathbf{k} \\ \tilde{\mathbf{y}}_j := R_j \mathbf{k} + \frac{1}{r_j} \tilde{\mathbf{C}}_j \times \mathbf{k} \\ \tilde{\mathbf{C}}_j := \begin{cases} \hat{\mathcal{S}}_j^{-1} (\hat{\chi}_{j-1} \mathbf{k} - \hat{\chi}_{j-2} \hat{\mathcal{S}}_j \hat{\mathcal{T}}_{j-1} \mathbf{k}) = \tilde{\mathbf{x}}_j \times \tilde{\mathbf{y}}_j & i = 2, \dots, n \\ \hat{\Theta}_1 \hat{\mathcal{S}}_1^{-1} \mathbf{k} & j = 1 \end{cases} \end{cases} \quad (56)$$

where $\hat{\mathcal{T}}_j, \hat{\mathcal{S}}_j$ have the expressions

$$\hat{\mathcal{T}}_j := \begin{cases} \mathcal{R}_3(\zeta) \mathcal{R}_1(\iota_n) & j = n \\ \mathcal{R}_3(\hat{\vartheta}_{j+1}) \mathcal{R}_1(\iota_j) & 1 \leq j \leq n-1 \end{cases} \quad \hat{\mathcal{S}}_j := \begin{cases} \mathcal{R}_3(\hat{\kappa}_{j-1}) \mathcal{R}_1(i_j), & 2 \leq j \leq n \\ \mathcal{R}_3(\hat{\vartheta}_1) \mathcal{R}_1(\frac{\pi}{2}), & j = 1 \end{cases} \quad (57)$$

with

$$\begin{cases} \cos \iota_n = \frac{Z}{\hat{\chi}_{n-1}} \\ \cos \iota_j = \frac{\hat{\Theta}_{j+1}}{\hat{\chi}_{j-1}} & 2 \leq j \leq n-1 \quad (n \geq 3) \\ \cos \iota_1 = \frac{\hat{\Theta}_2}{\hat{\Theta}_1} \\ \cos i_j := \frac{\hat{\Theta}_j}{\hat{\chi}_{j-1}}, & 2 \leq j \leq n \\ i_1 = \frac{\pi}{2} \end{cases} \quad (58)$$

Indeed, $\hat{\mathcal{T}}_j$ is the rotation matrix which describes the change of coordinates from $\hat{\mathcal{G}}_{j+1}$ to $\hat{\mathcal{F}}_j$, while $\hat{\mathcal{S}}_j$ describes the change of coordinates from $\hat{\mathcal{F}}_j$ to $\hat{\mathcal{G}}_j$, as it follows from the definitions of $(\hat{\Theta}, \hat{\chi}, \hat{\vartheta}, \hat{\kappa})$ in (51) (see also Figures 7, 8 and 9). The formulae (55)–(58) are obtained considering the following sequence of transformations

$$\begin{array}{ccccccccccc} \hat{\mathcal{T}}_n & \hat{\mathcal{S}}_n & \cdots & \hat{\mathcal{S}}_j & \hat{\mathcal{T}}_{j-1} & \cdots & \hat{\mathcal{S}}_1 \\ \hat{\mathcal{G}}_{n+1} & \rightarrow \hat{\mathcal{F}}_n & \rightarrow \hat{\mathcal{G}}_n & \rightarrow \cdots & \rightarrow \hat{\mathcal{F}}_j & \rightarrow \hat{\mathcal{G}}_j & \rightarrow \hat{\mathcal{F}}_{j-1} & \rightarrow \cdots & \rightarrow \hat{\mathcal{G}}_1 \end{array}$$

connecting \hat{G}_n to any other frame in the chain. From this, and the definitions of the frames (54), one finds

$$\mathbf{S}_j = \begin{cases} \chi_{j-1} \hat{\mathcal{T}}_n \hat{\mathcal{S}}_n \cdots \hat{\mathcal{T}}_{j+1} \hat{\mathcal{S}}_{j+1} \hat{\mathcal{T}}_j \mathbf{k} & j = 2 \dots n \\ \hat{\Theta}_1 \hat{\mathcal{T}}_n \hat{\mathcal{S}}_n \cdots \hat{\mathcal{T}}_2 \hat{\mathcal{S}}_2 \hat{\mathcal{T}}_1 \mathbf{k} & j = 1 \end{cases} \quad \mathbf{x}_j = r_j \hat{\mathcal{T}}_n \hat{\mathcal{S}}_n \cdots \hat{\mathcal{T}}_{j+1} \hat{\mathcal{S}}_{j+1} \hat{\mathcal{T}}_j \hat{\mathcal{S}}_j \mathbf{k}$$

whence

$$\mathbf{C}_j = \begin{cases} \mathbf{S}_j - \mathbf{S}_{j-1} = \hat{\mathcal{T}}_n \hat{\mathcal{S}}_n \cdots \hat{\mathcal{T}}_{j+1} \hat{\mathcal{S}}_{j+1} \hat{\mathcal{T}}_j (\hat{\chi}_{j-1} \mathbf{k} - \hat{\chi}_{j-2} \hat{\mathcal{S}}_j \hat{\mathcal{T}}_{j-1} \mathbf{k}) & j = 2, \dots, n \\ \mathbf{S}_1 = \hat{\Theta}_1 \hat{\mathcal{T}}_n \hat{\mathcal{S}}_n \cdots \hat{\mathcal{T}}_2 \hat{\mathcal{S}}_2 \hat{\mathcal{T}}_1 \mathbf{k} & j = 1 \end{cases}$$

and finally

$$\mathbf{y}_j = \frac{R_j}{r_j} \mathbf{x}_j + \frac{1}{r_j^2} \mathbf{C}_j \times \mathbf{x}_j$$

Collecting such formulae, one finds (55)–(58).

Canonical character of \mathcal{K}

Lemma 1.3 \mathcal{K} preserves the standard Liouville 1-form:

$$\sum_{j=1}^n \mathbf{y}_j \cdot d\mathbf{x}_j = \hat{\Theta} \cdot d\hat{\vartheta} + \hat{\chi} \cdot d\hat{\kappa} + \mathbf{R} \cdot d\mathbf{r}. \quad (59)$$

The proof of Lemma 1.3 again relies in Lemma 1.1.

Proof We use the expression in (55). We also define

$$\mathbf{C}_j^n := \mathcal{R}_j^n \tilde{\mathbf{C}}_j, \quad \bar{\mathbf{C}}_j^n := \bar{\mathcal{R}}_j^n \tilde{\mathbf{C}}_j, \quad \bar{\mathcal{R}}_j^n := \hat{\mathcal{T}}_n^{-1} \mathcal{R}_j^n$$

Applying Lemma 1.1 twice, we get

$$\mathbf{y}_j^n \cdot d\mathbf{x}_j^n = \mathbf{C}_j^n \cdot \mathbf{k} d\zeta + \bar{\mathbf{C}}_j^n \cdot \mathbf{i} dt_n + \bar{\mathbf{C}}_j^n \cdot \mathbf{k} d\hat{\kappa}_{n-1} + \mathbf{C}_j^{n-1} \cdot \mathbf{i} di_n + \mathbf{y}_j^{n-1} \cdot d\mathbf{x}_j^{n-1}.$$

Continuing in this way, after $n - j + 1$ iterates we arrive at

$$\begin{aligned} \mathbf{y}_j \cdot d\mathbf{x}_j &= \mathbf{C}_j^n \cdot \mathbf{k} d\zeta + \bar{\mathbf{C}}_j^n \cdot \mathbf{i} dt_n + \bar{\mathbf{C}}_j^n \cdot \mathbf{k} d\hat{\kappa}_{n-1} + \mathbf{C}_j^{n-1} \cdot \mathbf{i} di_n \\ &+ \sum_{k=j}^{n-1} \left(\mathbf{C}_j^k \cdot \mathbf{k} d\hat{\vartheta}_{k+1} + \bar{\mathbf{C}}_j^k \cdot \mathbf{i} dt_k + \bar{\mathbf{C}}_j^k \cdot \mathbf{k} d\hat{\kappa}_{k-1} + \mathbf{C}_j^{k-1} \cdot \mathbf{i} di_k \right) \\ &+ \tilde{\mathbf{y}}_j \cdot d\tilde{\mathbf{x}}_j \end{aligned} \quad (60)$$

with

$$i_1 := \frac{\pi}{2}, \quad \kappa_0 := \hat{\vartheta}_1, \quad \mathbf{C}_j^{j-1} := \tilde{\mathbf{C}}_j = \tilde{\mathbf{x}}_j \times \tilde{\mathbf{y}}_j.$$

We take the sum of (60) with $j = 1, \dots, n$. Exchanging the sums

$$\sum_{j=1}^n \sum_{k=j}^{n-1} = \sum_{k=1}^{n-1} \sum_{j=1}^k$$

and recognizing that

$$\left\{ \begin{array}{l} \sum_{j=1}^k \mathbf{C}_j^k = \begin{cases} \hat{\mathcal{S}}_{k+1}^{-1} \hat{\mathcal{T}}_{k+1}^{-1} \cdots \hat{\mathcal{S}}_n^{-1} \hat{\mathcal{T}}_n^{-1} \mathbf{S}_k = \chi_{k-1} \hat{\mathcal{T}}_k \mathbf{k} & 1 \leq k \leq n-1 \\ \mathbf{S}_n = \chi_{n-1} \hat{\mathcal{T}}_n \mathbf{k} & k = n \end{cases} \\ \sum_{j=1}^k \mathbf{C}_j^{k-1} = \begin{cases} \hat{\mathcal{S}}_k^{-1} \hat{\mathcal{T}}_k^{-1} \cdots \hat{\mathcal{S}}_n^{-1} \hat{\mathcal{T}}_n^{-1} \mathbf{S}_k = \chi_{k-1} \hat{\mathcal{S}}_k^{-1} \mathbf{k} & 1 \leq k \leq n-1 \\ \hat{\mathcal{S}}_n^{-1} \hat{\mathcal{T}}_n^{-1} \mathbf{S}_n = \chi_{n-1} \hat{\mathcal{S}}_n^{-1} \mathbf{k} & k = n \end{cases} \\ \sum_{j=1}^k \bar{\mathbf{C}}_j^k = \begin{cases} \hat{\mathcal{T}}_k^{-1} \hat{\mathcal{S}}_{k+1}^{-1} \hat{\mathcal{T}}_{k+1}^{-1} \cdots \hat{\mathcal{S}}_n^{-1} \hat{\mathcal{T}}_n^{-1} \mathbf{S}_k = \chi_{k-1} \mathbf{k} & 1 \leq k \leq n-1 \\ \hat{\mathcal{T}}_n^{-1} \mathbf{S}_n = \chi_{n-1} \mathbf{k} & k = n \end{cases} \end{array} \right.$$

with $\hat{\chi}_0 := \hat{\Theta}_1$ and that, by (56), the last term in (60) is

$$\tilde{\mathbf{y}}_j \cdot d\tilde{\mathbf{x}}_j = R_j dr_j$$

we get

$$\begin{aligned} \sum_{j=1}^n \mathbf{y}_j \cdot d\mathbf{x}_j &= \sum_{j=1}^n \left(\mathbf{C}_j^n \cdot \mathbf{k} d\zeta + \bar{\mathbf{C}}_j^n \cdot \mathbf{i} dl_n + \bar{\mathbf{C}}_j^n \cdot \mathbf{k} d\hat{\kappa}_{n-1} + \mathbf{C}_j^{n-1} \cdot \mathbf{i} di_n \right) \\ &+ \sum_{k=1}^{n-1} \sum_{j=1}^k \left(\mathbf{C}_j^k \cdot \mathbf{k} d\hat{\vartheta}_{k+1} + \bar{\mathbf{C}}_j^k \cdot \mathbf{i} dl_k + \bar{\mathbf{C}}_j^k \cdot \mathbf{k} d\hat{\kappa}_{k-1} + \mathbf{C}_j^{k-1} \cdot \mathbf{i} di_k \right) \\ &+ \sum_{j=1}^n R_j dr_j \\ &= \hat{\chi}_{n-1} \hat{\mathcal{T}}_n \mathbf{k} \cdot \mathbf{k} d\zeta + \hat{\chi}_{n-1} \mathbf{k} \cdot \mathbf{i} dl_n + \hat{\chi}_{n-1} \mathbf{k} \cdot \mathbf{k} d\hat{\kappa}_{n-1} + \hat{\chi}_{n-1} \mathbf{k} \cdot \hat{\mathcal{S}}_n \mathbf{i} di_n \\ &+ \sum_{k=1}^{n-1} \left(\hat{\chi}_{k-1} \hat{\mathcal{T}}_k \mathbf{k} \cdot \mathbf{k} d\hat{\vartheta}_{k+1} + \hat{\chi}_{k-1} \mathbf{k} \cdot \mathbf{i} dl_k + \hat{\chi}_{k-1} \mathbf{k} \cdot \mathbf{k} d\hat{\kappa}_{k-1} + \hat{\chi}_{k-1} \mathbf{k} \cdot \hat{\mathcal{S}}_k \mathbf{i} di_k \right) \\ &+ \sum_{j=1}^n R_j dr_j \\ &= \sum_{k=1}^n \hat{\Theta}_k d\hat{\vartheta}_k + \sum_{k=1}^n \hat{\chi}_k d\hat{\kappa}_k + \sum_{j=1}^n R_j dr_j \end{aligned}$$

having used

$$\hat{\mathcal{T}}_k \mathbf{k} \cdot \mathbf{k} = \cos \iota_k \quad \hat{\mathcal{S}}_k \mathbf{i} \cdot \mathbf{k} = 0, \quad \mathbf{k} \cdot \mathbf{k} = 1, \quad \mathbf{i} \cdot \mathbf{k} = 0$$

and $\cos \iota_k$ as in (58). In the following section, we shall use the following byproduct of Lemma 1.3. Recall the coordinates \mathcal{D}_{el} in (38) and denote

$$\phi_{\mathcal{D}_{el}}^{\mathcal{K}} : \mathcal{D}_{el} = (\mathbf{Z}, \mathbf{G}, \mathbf{R}, \zeta, \phi, \mathbf{r}) \rightarrow \mathcal{K} = (\hat{\Theta}, \hat{\chi}, \mathbf{R}, \hat{\vartheta}, \hat{\kappa}, \mathbf{r})$$

Consider the family of projections

$$\hat{\phi}_{\mathcal{D}_{el}}^{\mathcal{K}} : \mathcal{D}_{el} = (\mathbf{Z}, \mathbf{G}, \zeta, \phi) \rightarrow \mathcal{K} = (\hat{\Theta}, \hat{\chi}, \hat{\vartheta}, \hat{\kappa}) \quad (61)$$

which, as it is immediate to see, is independent of \mathbf{r} and \mathbf{R} .

Lemma 1.4 *The projections (61) verify*

$$\mathbf{Z} \cdot d\zeta + \mathbf{G} \cdot d\phi = \hat{\Theta} \cdot d\hat{\vartheta} + \hat{\chi} \cdot d\hat{\kappa} \quad \forall \mathbf{r}$$

1.6 The reduction of perihelia \mathcal{P}

The \mathcal{P} -coordinates have been described in [14]. Here, as in the case of \mathcal{K} , we change⁴ notations a little bit and denote them as

$$\mathcal{P} = (\Theta, \chi, \Lambda, \vartheta, \kappa, \ell) \in \mathbb{R}^n \times \mathbb{R}_+^n \times \mathbb{R}_+^n \times \mathbb{T}^n \times \mathbb{T}^n \times \mathbb{T}^n \quad (62)$$

where Λ, ℓ are as in (10), while

$$\begin{aligned} \Theta &= (\Theta_1, \dots, \Theta_n), & \vartheta &= (\vartheta_1, \dots, \vartheta_n) \\ \chi &= (\chi_1, \dots, \chi_n), & \kappa &= (\kappa_1, \dots, \kappa_n) \end{aligned}$$

are defined as follows. Consider a phase space where the Kepler Hamiltonians (6) take negative values. Let \mathbf{S}_j be as in (33) and \mathbf{P}_j the perihelia of the instantaneous ellipses generated by (6), assuming they are not circles. The coordinates Λ, ℓ are the same as in Delaunay, while, roughly, $(\Theta, \chi, \vartheta, \kappa)$ in (62) are defined as the $(\hat{\Theta}, \hat{\chi}, \hat{\vartheta}, \hat{\kappa})$ of \mathcal{K} , “replacing \mathbf{x}_j with \mathbf{P}_j ” (see Figures 10, 11, 12). Exact definitions are below.

Define the \mathcal{P} -nodes

$$\tilde{\nu}_j := \begin{cases} \mathbf{k} \times \mathbf{C} & j = n \\ \mathbf{P}_{j+1} \times \mathbf{S}_j & j = 1, \dots, n-1 \end{cases} \quad \tilde{\mathbf{n}}_j := \mathbf{S}_j \times \mathbf{P}_j \quad j = 1, \dots, n. \quad (63)$$

Then the \mathcal{P} -coordinates are

$$\begin{aligned} \Theta_j &:= \begin{cases} \mathbf{S}_j \cdot \mathbf{P}_j \\ |\mathbf{C}_1| \end{cases} & \vartheta_j &:= \begin{cases} \alpha_{\mathbf{P}_j}(\tilde{\mathbf{n}}_j, \tilde{\nu}_{j-1}) & 2 \leq j \leq n \\ \alpha_{\mathbf{C}_1}(\tilde{\nu}_1, \tilde{\mathbf{n}}_1) & j = 1 \end{cases} \\ \chi_j &:= \begin{cases} Z := \mathbf{C} \cdot \mathbf{k} \\ C := |\mathbf{C}| \\ |\mathbf{S}_{j+1}| \end{cases} & \kappa_j &:= \begin{cases} \zeta := \alpha_{\mathbf{k}}(\mathbf{i}, \tilde{\nu}_n) & j = n \\ \gamma := \alpha_{\mathbf{S}_n}(\tilde{\nu}_n, \tilde{\mathbf{n}}_n) & j = n-1 \\ \alpha_{\mathbf{S}_{j+1}}(\tilde{\nu}_{j+1}, \tilde{\mathbf{n}}_{j+1}) & 1 \leq j \leq n-2 \quad (n \geq 3) \end{cases} \end{aligned} \quad (64)$$

To prove that (62) are canonical, we consider the map

$$\phi_{\mathcal{D}_{el,aa}}^{\mathcal{P}} : \mathcal{D}_{el,aa} = (\mathbf{Z}, \mathbf{G}, \Lambda, \zeta, \mathbf{g}, \ell) \rightarrow \mathcal{P} = (\Theta, \chi, \Lambda, \vartheta, \kappa, \ell)$$

relating action-angle Delaunay (9) and \mathcal{P} and its projection

$$\hat{\phi}_{\mathcal{D}_{el,aa}}^{\mathcal{P}} : \mathcal{D}_{el,aa} = (\mathbf{Z}, \mathbf{G}, \zeta, \mathbf{g}) \rightarrow \mathcal{P} = (\Theta, \chi, \vartheta, \kappa)$$

which is independent of Λ, ℓ (even though this will not be used).

Lemma 1.5 $\hat{\phi}_{\mathcal{D}_{el,aa}}^{\mathcal{P}}$ coincides with the map $\hat{\phi}_{\mathcal{D}_{el}}^{\mathcal{K}}$ in (61).

Combining Lemmas 1.4 and 1.5, we have

Lemma 1.6 *The map*

$$\phi_{\mathcal{D}_{el,aa}}^{\mathcal{P}} : \mathcal{D}_{el,aa} = (\mathbf{Z}, \mathbf{G}, \Lambda, \zeta, \mathbf{g}, \ell) \rightarrow \mathcal{P} = (\Theta, \chi, \Lambda, \vartheta, \kappa, \ell)$$

verifies

$$\Theta \cdot d\vartheta + \chi \cdot d\kappa + \Lambda \cdot d\ell = \mathbf{Z} \cdot d\zeta + \mathbf{G} \cdot d\mathbf{g} + \Lambda \cdot d\ell.$$

⁴The coordinates named in [14] $\Theta_0, (\Theta_j)_{1 \leq j \leq n-1}, \chi_0, (\chi_j)_{1 \leq j \leq n-2}, \chi_{n-1}, \Lambda_j$ here are denoted, respectively, as $\chi_n, (\Theta_{n-j+1}), \chi_{n-1}, (\chi_{n-j+1}), \Theta_1, \Lambda_{n-j+1}$. An analogue change of notations holds for the conjugated coordinates.




Figure 10: The references \mathbf{F}_j and the \mathcal{P} -coordinates κ_{j-1} , $j = 2, \dots, n$.




Figure 11: The reference \mathbf{F}_1 and the \mathcal{P} -coordinates Θ_1, ϑ_1 .




Figure 12: The references \mathbf{G}_j and the \mathcal{P} -coordinates $\Theta_j, \vartheta_j, \chi_{j-2}$, $j = 1, \dots, n$. When $j = 2$, take $\chi_0 := \Theta_1$; when $j = 1$, disregard $\mathbf{S}_0, \boldsymbol{\nu}_0$ and χ_{-1} .

Explicit expression of the \mathcal{P} -map We now provide the explicit formulae of the map which relates the coordinates (64) to the coordinates $(\mathbf{y}_1, \dots, \mathbf{y}_n, \mathbf{x}_1, \dots, \mathbf{x}_n)$. We shall prove that such map has the expression

$$\begin{cases} \mathbf{x}_j = \mathbf{x}_j^n := \mathcal{R}_j^n \tilde{\mathbf{x}}_j \\ \mathbf{y}_j = \mathbf{y}_j^n := \mathcal{R}_j^n \tilde{\mathbf{y}}_j \end{cases} \quad (65)$$

where

$$\begin{cases} \mathcal{R}_j^n := \mathcal{T}_n \mathcal{S}_n \cdots \mathcal{T}_{j+1} \mathcal{S}_{j+1} \mathcal{T}_j \mathcal{S}_j \\ \tilde{\mathbf{x}}_j := a_j \left((\cos \xi_j - e_j) \mathbf{k} + \sqrt{1 - e_j^2} \sin \xi_j \tilde{\mathbf{Q}}_j \right) \\ \tilde{\mathbf{y}}_j := \frac{\mu_j n_j a_j}{1 - e_j \sin \xi_j} \left(-\sin \xi_j \mathbf{k} + \sqrt{1 - e_j^2} \cos \xi_j \tilde{\mathbf{Q}}_j \right) \end{cases} \quad (66)$$

where $\mathcal{T}_j, \mathcal{S}_j$ have the expressions

$$\mathcal{T}_j := \begin{cases} \mathcal{R}_3(\zeta) \mathcal{R}_1(\iota_n) & j = n \\ \mathcal{R}_3(\vartheta_{j+1}) \mathcal{R}_1(\iota_j) & 1 \leq j \leq n-1 \end{cases} \quad \mathcal{S}_j := \begin{cases} \mathcal{R}_3(\kappa_{j-1}) \mathcal{R}_1(i_j), & 2 \leq j \leq n \\ \mathcal{R}_3(\vartheta_1) \mathcal{R}_1(\frac{\pi}{2}), & j = 1 \end{cases} \quad (67)$$

with

$$\begin{cases} \cos \iota_n = \frac{Z}{\chi_{n-1}} \\ \cos \iota_j = \frac{\Theta_{j+1}}{\chi_{j-1}} & 2 \leq j \leq n-1 \quad (n \geq 3) \\ \cos \iota_1 = \frac{\Theta_2}{\Theta_1} \end{cases} \quad \begin{cases} \cos i_j := \frac{\Theta_j}{\chi_{j-1}}, & 2 \leq j \leq n \\ i_1 = \frac{\pi}{2} \end{cases} \quad (68)$$

and

$$\tilde{\mathbf{Q}}_j = \frac{\tilde{\mathbf{C}}_j}{C_j} \times \mathbf{k}$$

with

$$C_j = |\mathbf{C}_j| = \begin{cases} \sqrt{\chi_{j-1}^2 + \chi_{j-2}^2 - 2\Theta_j^2 + 2\sqrt{\chi_{j-1}^2 - \Theta_j^2} \sqrt{\chi_{j-2}^2 - \Theta_j^2} \cos \vartheta_j} & j = 2, \dots, n \\ \Theta_1 & j = 1 \end{cases}$$

$$\tilde{\mathbf{C}}_j := \begin{cases} \mathcal{S}_j^{-1} (\chi_{j-1} \mathbf{k} - \chi_{j-2} \mathcal{S}_j \mathcal{T}_{j-1} \mathbf{k}) = \tilde{\mathbf{x}}_j \times \tilde{\mathbf{y}}_j & i = 2, \dots, n \\ \Theta_1 \mathcal{S}_1^{-1} \mathbf{k} & j = 1 \end{cases}$$

$$e_j = \sqrt{1 - \frac{C_j^2}{\Lambda_j^2}}$$

a_j as in (10), $n_j = \sqrt{\frac{M_j}{a_j^3}}$ the mean motion, and ξ_j the eccentric anomaly, solving

$$\xi_j - e_j \sin \xi_j = \ell_j.$$

These formulae are easily obtained using the well-known relations

$$\begin{aligned}\mathbf{x}_j &= a_j \left((\cos \xi_j - e_j) \mathbf{P}_j + \sqrt{1 - e_j^2} \sin \xi_j \mathbf{Q}_j \right) \\ \mathbf{y}_j &:= \frac{\mu_j n_j a_j}{1 - e_j \sin \xi_j} \left(-\sin \xi_j \mathbf{P}_j + \sqrt{1 - e_j^2} \cos \xi_j \mathbf{Q}_j \right)\end{aligned}$$

with \mathbf{P}_j the j^{th} perihelion and $\mathbf{Q}_j = \frac{\mathbf{C}_j}{C_j} \times \mathbf{P}_j$, and the relations which relate \mathbf{C}_j , \mathbf{P}_j , \mathbf{Q}_j to \mathcal{P} , which, similarly to how done for \mathcal{K} , are:

$$\mathbf{C}_j = \mathcal{R}_j^n \tilde{\mathbf{C}}_j, \quad \mathbf{P}_j = \mathcal{R}_j^n \mathbf{k}, \quad \mathbf{Q}_j = \mathcal{R}_j^n \tilde{\mathbf{Q}}_j.$$

1.7 The behavior of \mathcal{K} and \mathcal{P} under reflections

The maps \mathcal{K} and \mathcal{P} have a nice behavior under reflections, which turns to be useful if they are applied to Hamiltonians which are reflection-invariant.

We denote as

$$\mathbf{x}^* = (x_1, -x_2, x_3) \tag{69}$$

the vector obtained from $\mathbf{x} = (x_1, x_2, x_3)$ by reflecting its second coordinate, and as

$$\mathcal{R}_2^- \left((\mathbf{y}_1, \dots, \mathbf{y}_n), (\mathbf{x}_1, \dots, \mathbf{x}_n) \right) := \left((\mathbf{y}_1^*, \dots, \mathbf{y}_n^*), (\mathbf{x}_1^*, \dots, \mathbf{x}_n^*) \right)$$

the simultaneous reflection of the second coordinate of all the \mathbf{y}_j and all the \mathbf{x}_j in the system of Cartesian coordinates $(\mathbf{y}, \mathbf{x}) = \left((\mathbf{y}_1, \dots, \mathbf{y}_n), (\mathbf{x}_1, \dots, \mathbf{x}_n) \right)$. We aim to show that

Lemma 1.7 *Using \mathcal{K} , the reflection \mathcal{R}_2^- is obtained by changing*

$$\left((\hat{\Theta}_2, \dots, \hat{\Theta}_n, Z), (\hat{\vartheta}_2, \dots, \hat{\vartheta}_n, \zeta) \right) \rightarrow \left((-\hat{\Theta}_2, \dots, -\hat{\Theta}_n, -Z), (-\hat{\vartheta}_2, \dots, -\hat{\vartheta}_n, -\zeta) \right)$$

Similarly, using \mathcal{P} , it is obtained by changing

$$\left((\Theta_2, \dots, \Theta_n, Z), (\vartheta_2, \dots, \vartheta_n, \zeta) \right) \rightarrow \left((-\Theta_2, \dots, -\Theta_n, -Z), (-\vartheta_2, \dots, -\vartheta_n, -\zeta) \right)$$

Proof We prove for \mathcal{K} . We write (69) as

$$\mathbf{x}^* = \mathcal{I}_2^- \mathbf{x} \quad \mathcal{I}_2^- = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Now use the formulae in (55)–(58) and that

$$\mathcal{I}_2^- \mathcal{R}_3(\alpha) = \mathcal{R}_3(-\alpha) \mathcal{I}_2^-, \quad \mathcal{I}_2^- \mathcal{R}_1(\beta) = \mathcal{R}_1(\pi - \beta) \mathcal{I}_2^-$$

and finally that the change

$$(\hat{\Theta}_2, \dots, \hat{\Theta}_n, Z) \rightarrow (-\hat{\Theta}_2, \dots, -\hat{\Theta}_n, -Z)$$

acts on the functions in (58) as

$$(\iota_1, \dots, \iota_n, \mathbf{i}_2, \dots, \mathbf{i}_n) \rightarrow (\pi - \iota_1, \dots, \pi - \iota_n, \pi - \mathbf{i}_2, \dots, \pi - \mathbf{i}_n).$$

The proof for \mathcal{P} is similar. \square

Lemma 1.7 reflects on the Hamiltonian (1) as well as in all Hamiltonians which are \mathcal{R}_2^- -invariant as follows.

Lemma 1.8 Let $\mathcal{H}(\mathbf{y}, \mathbf{x})$ be \mathcal{R}_2^- -invariant. Using the coordinates \mathcal{K} , the manifolds

$$\hat{\Theta}_j = 0, \quad \hat{\vartheta}_j \in \{0, \pi\} \quad j = 2, \dots, n \quad Z = 0, \quad \zeta \in \{0, \pi\}$$

are equilibria. Similarly, using the coordinates \mathcal{P} , the manifolds

$$\Theta_j = 0, \quad \vartheta_j \in \{0, \pi\} \quad j = 2, \dots, n \quad Z = 0, \quad \zeta \in \{0, \pi\}$$

are equilibria.

2 Applications

2.1 Arnold's Theorem

Here we retrace the main ideas of the proof of Theorem 1.1 given in [5]. Such proof uses on the coordinates (35). The first step is to switch from the coordinates (35) to a new set of coordinates which are well fitted with the close-to-be-integrable form of the Hamiltonian (4). Then we modify the coordinates (35) to the following form

$$\mathcal{D}_{ep,aa} = (\mathbf{\Lambda}, \mathbf{G}, \mathbf{\Psi}, \boldsymbol{\ell}, \boldsymbol{\gamma}, \boldsymbol{\psi}) \quad (70)$$

which we call *action-angle Deprit coordinates*, where $\mathbf{\Psi} = (\Psi_1, \dots, \Psi_n)$, $\boldsymbol{\psi} = (\psi_1, \dots, \psi_n)$ are left unvaried, while $\mathbf{\Lambda} = (\Lambda_1, \dots, \Lambda_n)$, $\mathbf{G} = (\Gamma_1, \dots, \Gamma_n)$, $\boldsymbol{\ell} = (\ell_1, \dots, \ell_n)$, $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_n)$ are obtained replacing the quadruplets $(R_i, G_i, r_i, \varphi_i)$ with the quadruplets $(\Lambda_i, \Gamma_i, \ell_i, \gamma_i)$ (with $G_i = \Gamma_i$), through the symplectic maps (depending on μ_i, M_i)

$$(R_i, G_i, r_i, \varphi_i) \rightarrow (\Lambda_i, \Gamma_i, \ell_i, \gamma_i)$$

which integrate Kepler Hamiltonian (6). This step is necessary to carry the integrable part in (4) to the form

$$h_K(\mathbf{\Lambda}) = \sum_{1 \leq i \leq n} \left(-\frac{\mu_i^3 M_i^2}{2\Lambda_i^2} \right).$$

Recall that the new angles γ_i provide the direction of the perihelion of the instantaneous ellipse generated by (6), however they have a different meaning compared to the analogous angles g_i appearing in the set of Delaunay coordinates (10), as, by construction, the γ_i 's are measured *relatively to the nodes* $\boldsymbol{\nu}_i$ in (34) (because the φ_i were), while the angles g_i in the Delaunay set are measured relatively to \boldsymbol{n}_i in (8).

The $3n - 2$ degrees of freedom Hamiltonian which is obtained is still singular. Singularities appear when the coordinates are not defined and in correspondence of collisions among the planets. The latter case will be later excluded through a careful choice of the reference frame. The singularities of the coordinates appear when the some of the convex angles (*Deprit inclinations*)

$$i_j^* := (\mathbf{S}_j, \mathbf{S}_{j+1}) \quad j = 1, \dots, n, \quad \mathbf{S}_{n+1} := \mathbf{k} \quad (71)$$

take the values 0 or π , because in such situations the angle ψ_j is not defined (see Figures 4, 5, 6) and when the instantaneous orbits of some of the Kepler Hamiltonians (6) is a circle, because in that case, the corresponding γ_i is not defined. Such singularities are important from the physical point of view, because the eccentricities and the inclinations of the planets of the solar system are very small, hence the system is in a configuration pretty close to the singularity. To deal with this situation, a regularization similar to the Poincaré regularization (11) of Delaunay coordinates has been introduced in [5]. Note that, in principle, there are 2^n singular configurations (corresponding to any choice of $i_j^* \in \{0, \pi\}$, besides $e_j = 0$ for some j). Here we discuss the case $i_j = 0$ for some j . Another regularization will be discussed in Section 2.3.

RPS coordinates and Birkhoff normal form The RPS variables are given by $(\mathbf{\Lambda}, \boldsymbol{\lambda}, \mathbf{z}) := (\mathbf{\Lambda}, \boldsymbol{\lambda}, \boldsymbol{\eta}, \boldsymbol{\xi}, \mathbf{p}, \mathbf{q})$ with (again) the Λ 's as in (10) and

$$\lambda_i = \ell_i + \gamma_i + \psi_{i-1}^n \quad \begin{cases} \eta_i = \sqrt{2(\Lambda_i - \Gamma_i)} \cos(\gamma_i + \psi_{i-1}^n) \\ \xi_i = -\sqrt{2(\Lambda_i - \Gamma_i)} \sin(\gamma_i + \psi_{i-1}^n) \end{cases} \quad (72)$$

$$\begin{cases} p_i = \sqrt{2(\Gamma_{i+1} + \Psi_{i-1} - \Psi_i)} \cos \psi_i^n \\ q_i = -\sqrt{2(\Gamma_{i+1} + \Psi_{i-1} - \Psi_i)} \sin \psi_i^n \end{cases}$$

where

$$\Psi_0 := \Gamma_1, \quad \Gamma_{n+1} := 0, \quad \psi_0 := 0, \quad \psi_i^n := \sum_{i \leq j \leq n} \psi_j. \quad (73)$$

Let ϕ_{RPS} denote the map

$$\phi_{\mathcal{C}}^{\text{RPS}} : (\mathbf{y}, \mathbf{x}) \rightarrow (\mathbf{\Lambda}, \boldsymbol{\lambda}, \mathbf{z}). \quad (74)$$

The main point is that

Lemma 2.1 ([5]) *The map $\phi_{\mathcal{C}}^{\text{RPS}}$ can be extended to a symplectic diffeomorphism on a set $\mathcal{P}_{\text{RPS}}^{6n}$ where the eccentricities e_j and the angles i_j^* in (71) are allowed to be zero. In particular,*

- $e_j = 0$ corresponds to the RPS coordinates $\eta_j = 0 = \xi_j$;
- $i_j^* = 0$ corresponds to the RPS coordinates $p_j = 0 = q_j$.

From the definitions (72)–(73) it follows that the variables

$$\begin{cases} p_n = \sqrt{2(\Psi_{n-1} - \Psi_n)} \cos \psi_n = \sqrt{2(C - Z)} \cos \zeta \\ q_n = -\sqrt{2(\Psi_{n-1} - \Psi_n)} \sin \psi_n = -\sqrt{2(C - Z)} \sin \zeta \end{cases} \quad (75)$$

are integrals (as they are defined only in terms of the integral \mathbf{C}), hence, cyclic for the Hamiltonian (4). Therefore, if \mathcal{H}_{RPS} denotes the planetary Hamiltonian expressed in RPS variables, we have that

$$\mathcal{H}_{\text{RPS}}(\mathbf{\Lambda}, \boldsymbol{\lambda}, \bar{\mathbf{z}}) := \mathcal{H} \circ \phi_{\mathcal{C}}^{\text{RPS}} = h_{\text{K}}(\mathbf{\Lambda}) + \mu f_{\text{RPS}}(\mathbf{\Lambda}, \boldsymbol{\lambda}, \bar{\mathbf{z}}) \quad (76)$$

where \mathcal{H} is as in (4) and ϕ_{RPS} as in (74) has $3n - 1$ degrees of freedom, as it depends on $\mathbf{\Lambda}, \boldsymbol{\lambda}, \bar{\mathbf{z}}$, where

$$\bar{\mathbf{z}} = (\boldsymbol{\eta}, \bar{\mathbf{p}}, \boldsymbol{\xi}, \bar{\mathbf{q}}) \quad \text{with} \quad \bar{\mathbf{p}} = (p_1, \dots, p_{n-1})$$

We denote as $a_i = \frac{1}{M_i} \left(\frac{\Lambda_i}{\mu_i} \right)^2$ the semi-major axis associated to Λ_i . The next result solves the problem of the construction of the Birkhoff normal form for the Hamiltonian (4), mentioned in Section 1.1.

Theorem 2.1 ([5, 4]) *For any $s \in \mathbb{N}$ there exists an open set $\mathcal{A} \subset \{a_1 < \dots < a_n\}$, a set $\mathcal{M}_{\varepsilon}^{6n-2} \subseteq \mathcal{A} \times \mathbb{T}^n \times \mathbb{R}^{4n}$ containing the strip $\mathcal{M}_0^{6n-2} = \mathcal{A} \times \mathbb{T}^n \times \{0\}_{\mathbb{R}^{4n}}$, a positive number ε and a symplectic map (“Birkhoff transformation”)*

$$\Phi_{\text{B}} : (\mathbf{\Lambda}, \mathbf{l}, \bar{\mathbf{w}}) \in \mathcal{M}_{\varepsilon}^{6n-2} \rightarrow (\mathbf{\Lambda}, \boldsymbol{\lambda}, \bar{\mathbf{z}}) \in \Phi_{\text{B}}(\mathcal{M}_{\varepsilon}^{6n-2}) \quad (77)$$

which carries the Hamiltonian (76) into

$$\mathcal{H}_{\text{B}}(\mathbf{\Lambda}, \mathbf{l}, \bar{\mathbf{w}}) := \tilde{\mathcal{H}}_{\text{RPS}} \circ \Phi_{\text{B}} = h_{\text{K}}(\mathbf{\Lambda}) + \mu f_{\text{B}}(\mathbf{\Lambda}, \mathbf{l}, \bar{\mathbf{w}}) \quad (78)$$

where the average $f_{\text{B}}^{\text{av}}(\mathbf{\Lambda}, w) := \int_{\mathbb{T}^n} f_{\text{B}} dl$ is in BNF of order s :

$$f_{\text{B}}^{\text{av}}(\mathbf{\Lambda}, \bar{\mathbf{w}}) = C_0 + \Omega \cdot \mathbf{r} + P_s(\mathbf{r}) + O(|\bar{\mathbf{w}}|^{2s+1}) \quad \bar{\mathbf{w}} := (\mathbf{u}, \mathbf{v}) \quad r_i := \frac{u_i^2 + v_i^2}{2}, \quad (79)$$

P_s being homogeneous polynomial in r of order s , parameterized by Λ . Furthermore, the normal form (78)–(79) is non-degenerate, in the sense that, if $s \geq 4$, the $(2n-1) \times (2n-1)$ matrix $\tau(\Lambda)$ of the coefficients of the monomial

$$\sum_{i,j=1}^{2n-1} \tau(\Lambda)_{ij} r_i r_j \quad (80)$$

with degree 2 in $P_s(\mathbf{r})$ is non singular, for all $\Lambda \in \mathcal{A}$.

Denote by $B_\varepsilon = B_\varepsilon^{2n_2} = \{y \in \mathbb{R}^{2n_2} : |y| < \varepsilon\}$ the $2n_2$ -ball of radius ε and let

$$\mathcal{P}_\varepsilon := V \times \mathbb{T}^{n_1} \times B_\varepsilon. \quad (81)$$

The second ingredient is a KAM theorem for properly-degenerate Hamiltonian systems. This has been stated and proved (with a proof of about 100 pages) by Arnold in [1], who named it the *Fundamental Theorem*. Here we present a refined version appeared in [2].

Theorem 2.2 (Fundamental Theorem, V.I. Arnold, 1963) *Let*

$$H(\mathbf{I}, \boldsymbol{\varphi}, \mathbf{p}, \mathbf{q}) := H_0(\mathbf{I}) + \mu P(\mathbf{I}, \boldsymbol{\varphi}, \mathbf{p}, \mathbf{q}), \quad (82)$$

be real-analytic on \mathcal{P}_ε and assume

(A1) $\mathbf{I} \in V \rightarrow \partial_{\mathbf{I}} H_0$ *is a diffeomorphism;*

(A2) $P_{\text{av}}(\mathbf{p}, \mathbf{q}; \mathbf{I}) = P_0(\mathbf{I}) + \sum_{i=1}^{n_2} \Omega_i(\mathbf{I}) r_i + \frac{1}{2} \sum_{i,j=1}^{n_2} \beta_{ij}(\mathbf{I}) r_i r_j + o_4$ *where $r_i := \frac{p_i^2 + q_i^2}{2}$ and $o_4/|(\mathbf{p}, \mathbf{q})|^4 \rightarrow 0$ as $(\mathbf{p}, \mathbf{q}) \rightarrow 0$;*

(A3) *The matrix $\beta(\mathbf{I}) = (\beta_{ij}(\mathbf{I}))$ is non-singular for all $\mathbf{I} \in V$.*

Then, there exist positive numbers ε_ , μ_* , C_* and b such that, for*

$$0 < \varepsilon < \varepsilon_*, \quad 0 < \mu < \mu_*, \quad \mu < \frac{1}{C_* (\log \varepsilon^{-1})^{2b}}, \quad (83)$$

one can find a set $\mathcal{T} \subset \mathcal{P}$ formed by the union of H -invariant $(n_1 + n_2)$ -dimensional tori, on which the H -motion is analytically conjugated to linear Diophantine quasi-periodic motions. The set \mathcal{T} is of positive Liouville–Lebesgue measure and satisfies

$$\text{meas } \mathcal{P}_\varepsilon > \text{meas } \mathcal{T} > \left(1 - C_* \left(\sqrt{\mu} (\log \varepsilon^{-1})^b + \sqrt{\varepsilon}\right)\right) \text{meas } \mathcal{P}_\varepsilon. \quad (84)$$

An application of Theorem 2.2 with $n_0 = n$, $n_1 = 2n - 1$ to the system in (78) with $s = 4$ now leads to the proof of Theorem 1.1.

2.2 Global Kolmogorov tori

The quasi-periodic motions of Theorem 1.1 provide almost circular and almost planar orbits. This is because the normal form of Theorem 2.1 is constructed around the strip \mathcal{M}_0^{6n-2} , and the origin corresponds to zero eccentricities and zero mutual inclinations. The question whether similar motions may exist outside such regime is therefore natural and important from the physical point of view. To this end, one has to understand that the Birkhoff normal form (assumption (A2) of

Theorem 2.2) is used in the proof only to construct a reasoning integrable approximation for the whole Hamiltonian, in fact given by

$$H_{\text{int}}(\mathbf{I}, \mathbf{r}) = H_0(\mathbf{I}) + \mu \left(P_0(\mathbf{I}) + \sum_{i=1}^{n_2} \Omega_i(\mathbf{I}) r_i + \frac{1}{2} \sum_{i,j=1}^{n_2} \beta_{ij}(\mathbf{I}) r_i r_j \right)$$

Therefore, a possible construction of full dimensional quasi-periodic motions outside the small eccentricities and small inclinations regime should start from a different integrable approximation. In this section we describe an approach in such direction, where we look at the first terms of the series expansion of the ℓ -averaged f with respect to a small parameter. The small parameter will be taken to be the inverse distance between the planets (the idea goes back to S. Harrington [9]). In addition, the use of the coordinates \mathcal{P} will allow to construct $(3n-2)$ -dimensional quasi-periodic motions without singularities when the inclinations become zero. Recall that the tori of Theorem 1.1 may be reduced to $(3n-2)$ frequencies (as shown in [5]), in a almost co-planar, co-centric configuration, but away from it, due to singularities. Here we show discuss the following result.

Theorem 2.3 (Global Kolmogorov tori in the planetary problem, [14]) *Fix numbers $0 < \underline{e}_i < \bar{e}_i < 0.6627\dots$, $i = 1, \dots, n$. There exists a number N depending only on n and a number α_0 depending on $\underline{e}_i, \bar{e}_i$, and n such that, if $\alpha < \alpha_0$, $\mu \leq \alpha^N$, in a domain of planetary motions where the semi-major axes $a_n < a_{n-1} < \dots < a_1$ are spaced as follows*

$$a_i^- \leq a_i \leq a_i^+ \quad \text{with} \quad a_i^\pm := \frac{a_n^\pm}{\alpha^{\frac{1}{3}(2^{n+1}-2^{i+1}+i-n)}} \quad (*)$$

there exists a positive measure set $\mathcal{K}_{\mu,\alpha}$, the density of which in phase space can be bounded below as

$$\text{dens}(\mathcal{K}_{\mu,\alpha}) \geq 1 - (\log \alpha^{-1})^p \sqrt{\alpha},$$

consisting of quasi-periodic motions with $3n-2$ frequencies where the planets' eccentricities e_i verify

$$\underline{e}_i \leq e_i \leq \bar{e}_i.$$

Let us consider a general set of coordinates $\mathcal{C} = (\mathbf{\Lambda}, \boldsymbol{\ell}, \mathbf{u}, \mathbf{v})$ which puts the Kepler Hamiltonians (6) into integrated form and hence carries the Hamiltonian (4) to

$$\mathcal{H}_{\mathcal{C}}(\mathbf{\Lambda}, \boldsymbol{\ell}, \mathbf{u}, \mathbf{v}) := \mathcal{H} \circ \mathcal{C} = - \sum_{j=1}^n \frac{\mu_j^3 M_j^2}{2\Lambda_j^2} + \mu f_{\mathcal{C}}(\mathbf{\Lambda}, \mathbf{u}, \mathbf{v}),$$

where

$$f_{\mathcal{C}}(\mathbf{\Lambda}, \boldsymbol{\ell}, \mathbf{u}, \mathbf{v}) := \sum_{1 \leq i < j \leq n} \left(\frac{\mathbf{y}_i \cdot \mathbf{y}_j}{m_0} - \frac{m_i m_j}{|\mathbf{x}_i - \mathbf{x}_j|} \right) \circ \mathcal{C}.$$

We denote

$$\overline{f_{\mathcal{C}}}(\mathbf{\Lambda}, \mathbf{u}, \mathbf{v}) := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} f_{\mathcal{C}}(\mathbf{\Lambda}, \boldsymbol{\ell}, \mathbf{u}, \mathbf{v}) d\boldsymbol{\ell}, \quad (85)$$

so that

$$f_{\mathcal{C}} = \sum_{1 \leq i < j \leq n} f_{\mathcal{C}}^{ij}, \quad \overline{f_{\mathcal{C}}} = \sum_{1 \leq i < j \leq n} \overline{f_{\mathcal{C}}^{ij}}$$

$$f_{\mathcal{C}}^{ij} := \left(\frac{\mathbf{y}_i \cdot \mathbf{y}_j}{m_0} - \frac{m_i m_j}{|\mathbf{x}_i - \mathbf{x}_j|} \right) \circ \mathcal{C}, \quad \overline{f_{\mathcal{C}}^{ij}} := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} f_{\mathcal{C}}^{ij} d\ell_1 \cdots d\ell_n.$$

For such any \mathcal{C} one always has, as a consequence of the motion equations of (6), the following identities

$$\begin{aligned}\frac{1}{2\pi} \int_{\mathbb{T}} \frac{1}{\mathbf{x}_j} dl_j &= \frac{1}{a_j} \\ \frac{1}{2\pi} \int_{\mathbb{T}} \mathbf{y}_j dl_j &= \frac{\mu_j}{2\pi} \int_{\mathbb{T}} \dot{\mathbf{x}}_j dl_j = 0 \\ \frac{1}{2\pi} \int_{\mathbb{T}} \frac{\mathbf{x}_j}{|\mathbf{x}_j|^3} dl_j &= \frac{1}{2\pi\mu_j M_j} \int_{\mathbb{T}} \dot{\mathbf{y}}_j dl_j = 0\end{aligned}\tag{86}$$

with a_j the semi-major axes. Consider now the average $\overline{f_{\mathcal{C}}}(\boldsymbol{\Lambda}, \mathbf{u}, \mathbf{v})$ in (85) with respect to ℓ . Due to the fact that \mathbf{y}_j has zero-average, one has that only the Newtonian part contributes to $\overline{f_{\mathcal{C}}}(\boldsymbol{\Lambda}, \mathbf{u}, \mathbf{v})$:

$$\overline{f_{\mathcal{C}}} = - \sum_{1 \leq i < j \leq n} \frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{dl_i dl_j}{|\mathbf{x}_i - \mathbf{x}_j|}.$$

We now consider any of the contributions to this sum

$$\overline{f_{\mathcal{C}}^{ij}} = - \frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{dl_i dl_j}{|\mathbf{x}_i - \mathbf{x}_j|} \quad 1 \leq i < j \leq n$$

and expand any such terms

$$\overline{f_{\mathcal{C}}^{ij}} = \overline{f_{\mathcal{C}}^{ij}}^{(0)} + \overline{f_{\mathcal{C}}^{ij}}^{(1)} + \overline{f_{\mathcal{C}}^{ij}}^{(2)} + \dots$$

where

$$\overline{f_{\mathcal{C}}^{ij}}^{(h)} := - \frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{1}{h!} \frac{d^h}{d\varepsilon^h} \frac{1}{|\mathbf{x}_i - \varepsilon \mathbf{x}_j|} \Big|_{\varepsilon=0} dl_i dl_j$$

is proportional to $\frac{1}{a_i} \left(\frac{a_i}{a_j}\right)^h$. Then the formulae in (86) imply that the two first terms of this expansion are given by

$$\overline{f_{\mathcal{C}}^{ij}}^{(0)} = - \frac{m_i m_j}{a_i}, \quad \overline{f_{\mathcal{C}}^{ij}}^{(1)} = 0.$$

Namely, whatever is the map \mathcal{C} that is used, the first non-trivial term is the double average of the second order term, which is given by

$$\overline{f_{\mathcal{C}}^{ij}}^{(2)}(\boldsymbol{\Lambda}, \mathbf{u}, \mathbf{v}) = - \frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{3(\mathbf{x}_i \cdot \mathbf{x}_j)^2 - |\mathbf{x}_i|^2 |\mathbf{x}_j|^2}{|\mathbf{x}_i|^5} dl_i dl_j.$$

Using Jacobi coordinates, S. Harrington noticed that

Lemma 2.2 ([9]) *If $n = 2$, $\overline{f_{\mathcal{J}}^{12}}^{(2)}$ depends on one only angle: the perihelion argument of the inner planet, hence is integrable.*

When $n = 2$, Lemma 2.2 provides an effective good starting point to construct quasi-periodic motions without the constraint of small eccentricities and inclinations, because in that case one can take, as initial approximation,

$$\mathcal{H}_{Harr} = - \sum_{j=1}^2 \frac{\mu_j^3 M_j^2}{2\Lambda_i^2} + \mu \left(- \frac{m_1 m_2}{a_2} + \overline{f_{\mathcal{J}}^{12}}^{(2)}(\Lambda_1, \Lambda_2, \Gamma_1, \Gamma_2, \gamma_1) \right)\tag{87}$$

The motions of \mathcal{H}_{Harr} have indeed widely studied in the literature, after [9]. When $n > 2$, the argument does not seem to have an immediate extension using Deprit coordinates (which, as said, are the natural extension of Jacobi reduction). The generalization of (87) for such a case is

$$\mathcal{H}_{\mathcal{D}_{ep,aa}} = - \sum_{j=1}^n \frac{\mu_j^3 M_j^2}{2\Lambda_i^2} + \mu \sum_{1 \leq i < j \leq n} \left(- \frac{m_i m_j}{a_j} + \overline{f_{\mathcal{J}}^{ij}}^{(2)} \right)$$

It turns out that, even looking at the nearest neighbors interactions

$$\mathcal{H}_{nn} = - \sum_{j=1}^n \frac{\mu_j^3 M_j^2}{2\Lambda_i^2} + \mu \sum_{i=1}^{n-1} \left(-\frac{m_i m_{i+1}}{a_i} + \overline{f_{\mathcal{D}_{ep}}^{i,i+1}}^{(2)} \right) \quad (88)$$

the terms $\overline{f_{\mathcal{D}_{ep}}^{i,i+1}}^{(2)}$ with $1 \leq i \leq n-2$ depend on two angles: γ_i and ψ_{i-1} , so the effective study of the unperturbed motions of (88) is involved. Using the \mathcal{P} -coordinates

$$\mathcal{H}_{nn} = - \sum_{j=1}^n \frac{\mu_j^3 M_j^2}{2\Lambda_i^2} + \mu \sum_{i=1}^{n-1} \left(-\frac{m_i m_{i+1}}{a_i} + \overline{f_{\mathcal{P}}^{i,i+1}}^{(2)} \right) \quad (89)$$

one has that the terms $\overline{f_{\mathcal{P}}^{i,i+1}}^{(2)}$ with $1 \leq i \leq n-2$ depend on 3 angles: κ_{i-1} , ϑ_i and ϑ_{i+1} , but the dependence upon κ_{i-1} and ϑ_i is at a higher order term. This is shown by the following formula, discussed in [14]:

$$\begin{aligned} \overline{f_{\mathcal{P}}^{i,i+1}}^{(2)} &= m_i m_{i+1} \frac{a_{i+1}^2}{4a_i^3} \frac{\Lambda_i^3}{\chi_{i-1}^2 (\chi_{i-1} - \chi_{i-2})^3} \left[\frac{5}{2} (3\Theta_{i+1}^2 - \chi_{i-1}^2) \right. \\ &- \frac{3}{2} \frac{4\Theta_{i+1}^2 - \chi_{i-1}^2}{\Lambda_{i+1}^2} \left(\chi_i^2 + \chi_{i-1}^2 - 2\Theta_{i+1}^2 + 2\sqrt{(\chi_i^2 - \Theta_{i+1}^2)(\chi_{i-1}^2 - \Theta_{i+1}^2)} \cos \vartheta_{i+1} \right) \\ &+ \frac{3}{2} \frac{(\chi_{i-1}^2 - \Theta_{i+1}^2)(\chi_i^2 - \Theta_{i+1}^2)}{\Lambda_{i+1}^2} \sin^2 \vartheta_{i+1} \\ &\left. + O(\Theta_i^2 + (\vartheta_i - \vartheta_i^0)^2) \right] \quad i = 1, \dots, n-1 \end{aligned} \quad (90)$$

where $\chi_0 := \Theta_1$, $\chi_{-1} := 0$, $\vartheta_i^0 \in \{0, \pi\}$ and the $O(\Theta_i^2 + (\vartheta_i - \vartheta_i^0)^2)$ term vanishes identically when $i = 1$.

The proof of Theorem 2.3 is based on two steps: in the first step the Hamiltonian is transformed to a similar one, but with a much smaller remainder. In the second step, a well fitted KAM theory is applied. Note that, as the terms of the unperturbed part are smaller and smaller as and when the distance from the sun increases, such KAM theory will be required to take such different scales into account.

Step 1: Normal Form Theory

Definition 2.1 Given $m, \nu_1, \dots, \nu_m \in \mathbb{N}$, $\nu := \nu_1 + \dots + \nu_m$; $\gamma_1, \dots, \gamma_m, \tau \in \mathbb{R}_+$. We call *m-scale Diophantine set*, and denote it as $\mathcal{D}_{\gamma_1, \dots, \gamma_m, \tau}$, the set of $\omega = (\omega_1, \dots, \omega_m)$, with $\omega_j \in \mathbb{R}^{\nu_j}$ such that, for any $k = (k_1, \dots, k_m) \in \mathbb{Z}^\nu \setminus \{0\}$, with $k_j \in \mathbb{Z}^{\nu_j}$, the following inequalities hold:

$$|\omega \cdot k| = \left| \sum_{j=1}^m \omega_j \cdot k_j \right| \geq \begin{cases} \frac{\gamma_1}{|k|^\tau} & \text{if } k_1 \neq 0; \\ \frac{\gamma_2}{|k|^\tau} & \text{if } k_1 = 0, \quad k_2 \neq 0; \\ \vdots & \\ \frac{\gamma_m}{|k_m|^\tau} & \text{if } k_1 = \dots = k_{m-1} = 0, \quad \dots, \quad k_m \neq 0. \end{cases} \quad (91)$$

The set $\mathcal{D}_{\gamma_1, \dots, \gamma_m, \tau}$ reduces to the usual diophantine set taking $\gamma_j = \gamma \forall j$. The first multi-scale Diophantine set was proposed by Arnold in [1] with $m = 2$.

Proposition 2.1 Let μ_j, M_j be as in (2) and $m_j := \sum_{i=1}^{j-1} m_i$, with $j = 2, \dots, n$. There exists a number c , depending only on $n, m_0, \dots, m_n, a_1^\pm, \varepsilon_j, \bar{\varepsilon}_j$, and a number $0 < \bar{c} < 1$, depending only on n such that, for any fixed positive numbers $\bar{\gamma} < 1 < \bar{K}, \alpha > 0$ verifying

$$\bar{K} \leq \frac{c}{\alpha^{3/2}} \quad (92)$$

and

$$\frac{1}{c} \max \left\{ \mu \left(\frac{a_n^+}{a_1^-} \right)^5 \frac{\bar{K}^{2\bar{\gamma}+2}}{\bar{\gamma}^2}, \frac{\bar{K}^{2(\bar{\gamma}+1)} \alpha}{\bar{\gamma}^2} \right\} < 1 \quad (93)$$

there exist natural numbers ν_1, \dots, ν_{2n-1} , with $\sum_j \nu_j = 3n - 2$, open sets $B_j^* \subset B_{\varepsilon_j}^2, \mathcal{X}^* \subset \mathcal{X}$, positive real numbers $\gamma_1 > \dots > \gamma_{2n-1} \varepsilon_1, \dots, \varepsilon_{n-1}, \bar{r}_1, \dots, \bar{r}_{n-1}, \tilde{r}_1, \dots, \tilde{r}_n$, a domain

$$D_n := B_{\sqrt{2\bar{\gamma}}} \times \mathcal{X}_{\bar{\gamma}} \times \mathcal{A}_{\bar{\gamma}} \times \mathbb{T}_{\bar{c}s}^n \times \mathbb{T}_{\bar{c}s}^n$$

a sub-domain of the form

$$D_n^* := B_{\sqrt{2\bar{\gamma}}}^* \times \mathcal{X}_{\bar{\gamma}}^* \times \mathcal{A}_{\bar{\gamma}} \times \mathbb{T}_{\bar{c}s}^n \times \mathbb{T}_{\bar{c}s}^n$$

verifying

$$\text{meas } D_n^* \geq \left(1 - \frac{\bar{\gamma}}{c}\right) \text{meas } D_n \quad (94)$$

a real-analytic transformation

$$\phi_n : (p, q, \chi, \Lambda, \kappa, \ell) \in D_n^* \rightarrow D_{\mathcal{P}}$$

which conjugates $\mathcal{H}_{\mathcal{P}}$ to

$$\mathcal{H}_n(p, q, \chi, \Lambda, \kappa, \ell) := \mathcal{H}_{\mathcal{P}} \circ \phi_n = h_{fast,sec}(p, q, \chi, \Lambda) + \mu f_{exp}(p, q, \chi, \Lambda, \kappa, \ell)$$

where $f_{exp}(p, q, \chi, \Lambda, \kappa, \ell)$ is independent of κ_{n-1} , and the following holds.

1. The function $h_{fast,sec}(p, q, \chi, \Lambda)$ is a sum

$$h_{fast,sec}(p, q, \chi, \Lambda) = h_{fast}(\Lambda) + \mu h_{sec}(p, q, \chi, \Lambda)$$

where, if

$$\hat{y}_i := \left(\frac{p_2^2 + q_2^2}{2}, \dots, \frac{p_{i+1}^2 + q_{i+1}^2}{2}, \chi_0, \dots, \chi_i, \Lambda_1, \dots, \Lambda_{i+1} \right) \quad i = 1, \dots, n-1$$

then h_{fast} and h_{sec} are given by

$$h_{fast}(\Lambda) = - \sum_{j=1}^n \frac{m_j^3 M_j^2}{2\Lambda_j^2} - \mu \sum_{j=1}^{n-1} \frac{M_j m_j^2 m_j m_j}{\Lambda_j^2}, \quad h_{sec}(p, q, \chi, \Lambda) = \sum_{i=1}^{n-1} h_{sec}^i(\hat{y}_i)$$

where the functions h_{sec}^i have an analytic extension on D_n and verify

$$c \frac{(a_{j+1}^+)^2}{(a_j^-)^3} \leq |h_{sec}^j(\hat{y}_j)| \leq \frac{1}{c} \frac{(a_{j+1}^+)^2}{(a_j^-)^3}.$$

2. The function f_{exp} satisfies

$$|f_{exp}| \leq \frac{1}{c} \frac{e^{-c\bar{K}}}{a_n^-}.$$

3. If ζ is \hat{y}_1 deprived of χ_0 , the frequency-map

$$\zeta \rightarrow \omega_{fast,sec}(\zeta) := \partial_{\zeta} h_{fast,sec}(\zeta)$$

is a diffeomorphism of $\Pi_{\zeta}(B_{\sqrt{2\bar{\tau}}}^* \times \mathcal{X}_{\bar{\tau}}^* \times \mathcal{A}_{\bar{\tau}}^*)$ and, moreover, it satisfies (91), with $m = 2n - 1$, $\tau = \bar{\tau} > 2$, and

$$\nu_j := \begin{cases} 1 & j = 1, \dots, n \\ 2 & j = 3, n = 2 \\ 3 & j = n + 1, n \geq 3 \\ 2 & n + 2 \leq j \leq 2n - 2, n \geq 4 \\ 1 & j = 2n - 1, n \geq 3 \end{cases}$$

$$\omega_j := \begin{cases} \partial_{\Lambda_j} h_{fast,sec} & j = 1, \dots, n \\ \partial_{(\frac{p_2^2+q_2^2}{2}, \chi_0)} h_{fast,sec} & j = 3, n = 2 \\ \partial_{(\frac{p_2^2+q_2^2}{2}, \chi_1, \chi_0)} h_{fast,sec} & j = n + 1, n \geq 3 \\ \partial_{(\frac{p_{j-n+1}^2+q_{j-n+1}^2}{2}, \chi_{j-n})} h_{fast,sec} & n + 2 \leq j \leq 2n - 2, n \geq 4 \\ \partial_{\frac{p_n^2+q_n^2}{2}} h_{fast,sec} & j = 2n - 1, n \geq 3 \end{cases}$$

$$\gamma_j := \begin{cases} \frac{1}{a_j^-} \frac{\bar{\gamma}}{\theta_j} & 1 \leq j \leq n \\ \frac{\mu(a_{2n-j+1}^+)^2}{(a_{2n-j}^-)^3} \frac{\bar{\gamma}}{\theta_{j-n}} & n + 1 \leq j \leq 2n - 1 \end{cases} \quad (95)$$

4. The mentioned constants are

$$\varepsilon_j := c \sqrt{\theta_j}, \quad \bar{r}_j := \frac{\theta_j \bar{\gamma}}{K^{\bar{\tau}+1}}, \quad \tilde{r}_i := c \theta_j$$

with $\bar{\tau} > 2$.

Step 2: KAM theory

Theorem 2.4 (Multi-scale KAM Theorem, [14]) *Let $m, \ell, \nu_1, \dots, \nu_m \in \mathbb{N}$, $\nu := \nu_1 + \dots + \nu_m \geq \ell$, $\tau_* > \nu$, $\gamma_1 \geq \dots \geq \gamma_m > 0$, $0 < 4s \leq \bar{s} < 1$, $\rho_1, \dots, \rho_\ell, r_1, \dots, r_{\nu-\ell}, \varepsilon_1, \dots, \varepsilon_\ell > 0$, $B_1, \dots, B_\ell \subset \mathbb{R}^2$, $D_j := \{\frac{x^2+y^2}{2} \in \mathbb{R} : (x, y) \in B_j\} \subset \mathbb{R}$, $B := B_1 \times \dots \times B_\ell \subset \mathbb{R}^{2\ell}$, $D := D_1 \times \dots \times D_\ell \subset \mathbb{R}^\ell$, $C \subset \mathbb{R}^{\nu-\ell}$, $A := D_\rho \times C_r$. Let*

$$H(\mathbf{p}, \mathbf{q}, \mathbf{I}, \psi) = h(\mathbf{p}, \mathbf{q}, \mathbf{I}) + f(\mathbf{p}, \mathbf{q}, \mathbf{I}, \psi)$$

be real-analytic on $B_{\sqrt{2\rho}} \times C_r \times \mathbb{T}_{\bar{s}+s}^{\nu-\ell}$, where $h(\mathbf{p}, \mathbf{q}, \mathbf{I})$ depends on (\mathbf{p}, \mathbf{q}) only via

$$J(\mathbf{p}, \mathbf{q}) := \left(\frac{p_1^2 + q_1^2}{2}, \dots, \frac{p_\ell^2 + q_\ell^2}{2} \right).$$

Assume that $\omega_0 := \partial_{J(\mathbf{p}, \mathbf{q}, \mathbf{I})} \mathbf{h}$ is a diffeomorphism of A with non singular Hessian matrix $U := \partial_{J(\mathbf{p}, \mathbf{q}, \mathbf{I})}^2 \mathbf{h}$ and let U_k denote the $(\nu_k + \dots + \nu_m) \times \nu$ submatrix of U , i.e., the matrix with entries $(U_k)_{ij} = U_{ij}$, for $\nu_1 + \dots + \nu_{k-1} + 1 \leq i \leq \nu$, $1 \leq j \leq \nu$, where $2 \leq k \leq m$. Let

$$\begin{aligned} M &\geq \sup_A |U|, \quad M_k \geq \sup_A |U_k|, \quad \bar{M} \geq \sup_A |U^{-1}|, \quad E \geq |f|_{\rho, \bar{s}+s} \\ \bar{M}_k &\geq \sup_A |T_k| \quad \text{if} \quad U^{-1} = \begin{pmatrix} T_1 \\ \vdots \\ T_m \end{pmatrix} \quad 1 \leq k \leq m. \end{aligned}$$

Define

$$\begin{aligned} K &:= \frac{6}{s} \log_+ \left(\frac{EM_1^2 L}{\gamma_1^2} \right)^{-1} \quad \text{where} \quad \log_+ a := \max\{1, \log a\} \\ \hat{\rho}_k &:= \frac{\gamma_k}{3M_k K^{\tau_*+1}}, \quad \hat{\rho} := \min\{\hat{\rho}_1, \dots, \hat{\rho}_m, \rho_1, \dots, \rho_\ell, r_1, \dots, r_{\nu-\ell}\} \\ L &:= \max\{\bar{M}, M_1^{-1}, \dots, M_m^{-1}\} \\ \hat{E} &:= \frac{EL}{\hat{\rho}^2}. \end{aligned}$$

Then one can find two numbers $\hat{c}_\nu > c_\nu$ depending only on ν such that, if the perturbation f is so small that the following ‘‘KAM condition’’ holds

$$\hat{c}_\nu \hat{E} < 1,$$

for any $\omega \in \Omega_* := \omega_0(D) \cap \mathcal{D}_{\gamma_1, \dots, \gamma_m, \tau_*}$, one can find a unique real-analytic embedding

$$\begin{aligned} \phi_\omega : \quad \vartheta = (\hat{\vartheta}, \bar{\vartheta}) \in \mathbb{T}^\nu &\rightarrow (\hat{v}(\vartheta; \omega), \hat{\vartheta} + \hat{u}(\vartheta; \omega), \mathcal{R}_{\bar{\vartheta} + \bar{u}(\vartheta; \omega)} w_1, \dots, \mathcal{R}_{\bar{\vartheta} + \bar{u}(\vartheta; \omega)} w_\ell) \\ &\in \text{Re } C_r \times \mathbb{T}^{\nu-\ell} \times \text{Re } B_{\sqrt{2r}}^{2\ell} \end{aligned}$$

where $r := c_\nu \hat{E} \hat{\rho}$ such that $\mathbb{T}_\omega := \phi_\omega(\mathbb{T}^\nu)$ is a real-analytic ν -dimensional \mathbf{H} -invariant torus, on which the \mathbf{H} -flow is analytically conjugated to $\vartheta \rightarrow \vartheta + \omega t$. Furthermore, the map $(\vartheta; \omega) \rightarrow \phi_\omega(\vartheta)$ is Lipschitz and one-to-one and the invariant set $\mathbf{K} := \bigcup_{\omega \in \Omega_*} \mathbb{T}_\omega$ satisfies the following measure estimate

$$\text{meas} \left(\text{Re} (D_r) \times \mathbb{T}^n \setminus \mathbf{K} \right) \leq c_\nu \left(\text{meas} (D \setminus D_{\gamma_1, \dots, \gamma_m, \tau_*}) \times \mathbb{T}^n + \text{meas} (\text{Re} (D_r) \setminus D) \times \mathbb{T}^n \right),$$

where $D_{\gamma_1, \dots, \gamma_m, \tau_*}$ denotes the ω_0 -pre-image of $\mathcal{D}_{\gamma_1, \dots, \gamma_m, \tau_*}$ in D . Finally, on $\mathbb{T}^\nu \times \Omega_*$, the following uniform estimates hold

$$\begin{aligned} |v_k(\cdot; \omega) - I_k^0(\omega)| &\leq c_\nu \left(\frac{\bar{M}_k}{M} + \frac{M_k}{M_1} \right) \hat{E} \hat{\rho} \\ |u(\cdot; \omega)| &\leq c_\nu \hat{E} s \end{aligned}$$

where v_k denotes the projection of $v = (\hat{v}, \bar{v}) \in \mathbb{R}^{\nu_1} \times \dots \times \mathbb{R}^{\nu_m}$ over \mathbb{R}^{ν_k} , $\bar{v}_k := \frac{|w_k|^2}{2}$ and $I^0(\omega) = (I_1^0(\omega), \dots, I_\nu^0(\omega)) \in D$ is the ω_0 -pre-image of $\omega \in \Omega_*$.

Theorem 2.4 generalizes Theorem 3 in [2] and hence the Fundamental Theorem of [1], to which Theorem 3 in [2] is inspired.

Proof of Theorem 2.3 Let

$$\bar{\gamma} := \bar{c}\sqrt{\alpha}(\log \alpha^{-1})^{\bar{\tau}+1}, \quad \bar{K} = \frac{1}{\bar{c}} \log \frac{1}{\alpha}$$

where \bar{c} is as in (94) and \bar{c} will be fixed later. We aim to apply Theorem 2.4 to the Hamiltonian \mathcal{H}_n of Proposition 2.1, with these choices of $\bar{\gamma}$ and \bar{K} . To this end, we take

$$\begin{aligned} M_j &= \begin{cases} \frac{1}{c_1 a_j^- \theta_j^2} & 1 \leq j \leq n \\ \frac{\mu(a_{2n-j+1}^+)^2}{c_1 (a_{2n-j}^-)^3 \theta_j^2} & n+1 \leq j \leq 2n-1 \end{cases} & L = \bar{M} = \frac{1}{c_2} \frac{\theta_1^2 (a_n^+)^3}{\mu (a_{n-1}^-)^2} \\ E &= \frac{1}{c_3} \frac{\mu}{a_n^-} e^{-c\bar{K}} & K = \frac{1}{c_4} \log_+ \left(\frac{1}{\bar{\gamma}^2} \frac{(a_n)^3}{(a_{n-1}^-)^3} e^{-c\bar{K}} \right)^{-1} \\ \hat{\rho}_j &= \begin{cases} c_5 \frac{\bar{\gamma} \theta_j}{K^{\tau_*+1}} & 1 \leq j \leq n \\ c_5 \frac{\bar{\gamma} \theta_{j-n}}{K^{\tau_*+1}} & n+1 \leq j \leq 2n-1 \end{cases} & \hat{\rho} := \frac{\theta_1 \bar{\gamma}}{\bar{K}^{\tau_*+1}} \quad \tau_* > 3n-2 \\ \hat{E} &= \frac{1}{c_6} \frac{1}{\bar{\gamma}^2} \frac{(a_n)^3}{(a_{n-1}^-)^3} e^{-c\bar{K}} \hat{K}^{2(\tau_*+1)} \end{aligned}$$

where $\hat{K} := \max\{K, \bar{K}\}$. The number $\frac{1}{\bar{\gamma}^2} \frac{(a_n)^3}{(a_{n-1}^-)^3}$ can be bounded by $\frac{1}{\alpha^N}$ for a sufficiently large N depending only on n . Hence, if $\bar{c} < \frac{c}{N}$ and $\alpha < c_6$, we have $\hat{E} < 1$ and the theorem is proved. \square

2.3 On the co-existence of stable and whiskered tori

In this section we discuss how the use two different sets of coordinates may lead to prove the co-existence of stable and unstable motions. Specifically, we deal with the following situation, which we shall refer to as *outer, retrograde configuration* (ORC):

Two planets describe almost co-planar orbits, revolving around their common sun, in opposite sense. The outer planet has a lower angular momentum and retrograde motion, as seen from the total angular momentum of the system.

We aim to discuss the following

Theorem 2.5

1. *There exists a region \mathcal{D}_s in the phase space almost completely filled with a positive measure set of five-dimensional KAM tori, in ORC configuration;*
2. *There exists a region \mathcal{D}_u in the phase space including an invariant region \mathcal{D}_u^0 consisting of co-planar, retrograde motions for the outer planet as a hyperbolic equilibrium manifold for the three-body system;*
3. *\mathcal{D}_s and \mathcal{D}_u^0 have a non-empty intersection.*

Theorem 2.5 leads to the following conjecture, which is likely to be proved somewhere.

Conjecture 2.1 *Full dimensional quasi-periodic motions and hyperbolic 3-dimensional tori co-exist in \mathcal{D}_s .*

The proof of statements 1. and 2. in Theorem 2.5 relies on the use of two different sets of coordinates for the Hamiltonian (4) with $n = 2$:

$$\mathcal{H}_{3BP} = \frac{|\mathbf{y}_1|^2}{2\mu_1} - \frac{\mu_1 M_1}{|\mathbf{x}_1|} + \frac{|\mathbf{y}_2|^2}{2\mu_2} - \frac{\mu_2 M_2}{|\mathbf{x}_2|} + \mu \left(\frac{\mathbf{y}_1 \cdot \mathbf{y}_2}{m_0} - \frac{m_1 m_2}{|\mathbf{x}_i - \mathbf{x}_j|} \right) \quad (96)$$

Proof of 1. We consider the coordinates (70) with $n = 2$. It will turn to be useful to work with regularizing complex coordinates, which we denote as

$$\text{RPS}_\pi^{\mathbb{C}} := (\mathbf{\Lambda}, \mathbf{\lambda}, \mathbf{t}, \mathbf{t}^*, T, T^*) = (\Lambda_1, \Lambda_2, \lambda_1, \lambda_2, t_1, t_2, t_3, t_1^*, t_2^*, t_3^*, T, T^*) \quad (97)$$

and define via the formulae

$$\begin{cases} \Lambda_1 = \Lambda_1 \\ \Lambda_2 = \Lambda_2 \\ t_1 = -i\sqrt{\Lambda_1 - \Gamma_1} e^{i(-\gamma_1 + \gamma + \zeta)} \\ t_2 = \sqrt{\Lambda_2 - \Gamma_2} e^{i(\gamma_2 + \gamma + \zeta)} \\ t_3 = -i\sqrt{C - \Gamma_2 + \Gamma_1} e^{i(\gamma + \zeta)} \\ T = \sqrt{C - Z} e^{i\zeta} \end{cases} \quad \begin{cases} \lambda_2 = \ell_2 + \gamma_2 + \gamma + \zeta \\ \lambda_1 = \ell_1 + \gamma_1 - \gamma - \zeta \\ t_1^* = -\sqrt{\Lambda_1 - \Gamma_1} e^{-i(-\gamma_1 + \gamma + \zeta)} \\ t_2^* = -i\sqrt{\Lambda_2 - \Gamma_2} e^{-i(\gamma_2 + \gamma + \zeta)} \\ t_3^* = -\sqrt{C - \Gamma_2 + \Gamma_1} e^{-i(\gamma + \zeta)} \\ T^* = -i\sqrt{C - Z} e^{-i\zeta} \end{cases} \quad (98)$$

We also define, for later need, $\eta_1, \eta_2, p, \xi_1, \xi_2, q$ via

$$\begin{aligned} t_2 &:= \frac{\eta_2 - i\xi_2}{\sqrt{2}} & t_1 &:= \frac{i\eta_1 - \xi_1}{\sqrt{2}} & t_3 &:= \frac{ip - q}{\sqrt{2}} & T &:= \frac{P - iQ}{\sqrt{2}} \\ t_2^* &:= \frac{\eta_2 + i\xi_2}{\sqrt{2i}} & t_1^* &:= \frac{i\eta_1 + \xi_1}{\sqrt{2i}} & t_3^* &:= \frac{ip + q}{\sqrt{2i}} & T^* &:= \frac{P + iQ}{\sqrt{2i}}. \end{aligned} \quad (99)$$

Observe that

$$\mathcal{M}_\pi := \{(\mathbf{\Lambda}, \mathbf{\lambda}, \mathbf{t}, \mathbf{t}^*) : (\mathbf{t}, \mathbf{t}^*) = (0, 0)\} \quad (100)$$

corresponds to co-circular, co-planar orbits for the two planets, with the outer planet in retrograde motion.

We denote as

$$\mathcal{H}_{\text{RPS}_\pi^{\mathbb{C}}} = -\frac{\mu_1^3 M_1^2}{2\Lambda_1^2} - \frac{\mu_2^3 M_2^2}{2\Lambda_2^2} + \mu f_{\text{RPS}_\pi^{\mathbb{C}}}(\mathbf{\Lambda}, \mathbf{\lambda}, \mathbf{t}, \mathbf{t}^*) \quad (101)$$

the expression of the Hamiltonian (96) using the coordinates $\text{RPS}_\pi^{\mathbb{C}}$ in (97), which, similarly to the prograde case, $\mathcal{H}_{\text{RPS}_\pi^{\mathbb{C}}}$ is independent of (T, T^*) . Abusively, we shall continue calling $\text{RPS}_\pi^{\mathbb{C}}$ the coordinates (97) deprived of (T, T^*) .

We now define a domain where letting the $\text{RPS}_\pi^{\mathbb{C}}$ coordinates vary. First of all, we observe that ORC configuration can be realized only if the planetary masses are tuned with the semi-major axes. More precisely, that, if we denote as “2” and “1” the inner⁵, outer planet; as a_2, a_1 , the semi-major axes of their respective instantaneous orbits around the sun; α_-, α_+ , with $0 < \alpha_- < \alpha_+ < 1$, two numbers such that the semi-axes ratio $\alpha := \frac{a_2}{a_1}$ verifies

$$\alpha_- < \alpha < \alpha_+, \quad (102)$$

then the following inequality needs to be satisfied

$$\frac{m_2}{m_1} \sqrt{\alpha_-} > 1. \quad (103)$$

Indeed, since the motions are almost-circular, the lengths of the angular momenta of the planets, C_1, C_2 are arbitrarily close to the action coordinates Λ_1, Λ_2 related to their semi-major axes, which in turn are related to the semi-axes and the mass ratio via

$$1 < \frac{C_2}{C_1} \sim \frac{\Lambda_2}{\Lambda_1} = \frac{\mu_2}{\mu_1} \sqrt{\frac{M_2}{M_1}} \sqrt{\alpha}$$

⁵Compared to [13], here “2” and “1” are exchanged, in order to keep uniform notations along the paper.

where μ_i, M_i are as in (5). This inequality does not make conflict with (102) if one assumes that

$$k_{\pm} := \frac{\mu_2}{\mu_1} \sqrt{\frac{M_2}{M_1}} \alpha_{\pm} > 1. \quad (104)$$

whence the necessity of (103).

We then fix the domain as follows. The coordinates Λ_1, Λ_2 will be taken to vary in the set

$$\mathcal{L} := \left\{ \Lambda = (\Lambda_1, \Lambda_2) : \Lambda_- \leq \Lambda_1 \leq \Lambda_+, k_- \Lambda_1 \leq \Lambda_2 \leq k_+ \Lambda_1 \right\} \quad (105)$$

with k_{\pm} as in (104), and $0 < \Lambda_- < \Lambda_+$ to be chosen later.

The coordinates $\lambda = (\lambda_1, \lambda_2)$ will be taken to run in the torus \mathbb{T}^2 .

As for the coordinates $(\mathbf{t}, \mathbf{t}^*)$, we take a domain of the form

$$\mathcal{U}_s := \left\{ (\mathbf{t}, \mathbf{t}^*) \in \mathbb{C}^6 : |(\mathbf{t}, \mathbf{t}^*)| \leq \varepsilon \right\}$$

The domain for $\text{RPS}_{\pi}^{\mathbb{C}}$ will then be

$$\mathcal{D}_s = \mathcal{L} \times \mathbb{T}^2 \times \mathcal{U}_s. \quad (106)$$

The following statement is a more precise version of statement 1. in Theorem 2.5.

Theorem 2.6 ([13]) *There exist two numbers $0 < \varepsilon_+ < \varepsilon_0, 0 < \alpha_+ < 1$, such that, for any $0 < \varepsilon < \varepsilon_+, 0 < \alpha_- < \alpha_+, 0 < \Lambda_- < \Lambda_+$, one can find $\mu_+(\varepsilon) > 0$ such that, for any $0 < \mu < \mu_+(\varepsilon)$, in the domain \mathcal{D}_s there exists an invariant set $\mathcal{F}_{\varepsilon, \mu} \subset \mathcal{D}_s$ with density going to 1 as $\varepsilon \rightarrow 0$ which is foliated as*

$$\mathcal{F}_{\varepsilon, \mu} = \bigcup_{\omega} \mathcal{T}_{\omega, \varepsilon, \mu} \quad (107)$$

where $\mathcal{T}_{\omega, \varepsilon, \mu}$ is diffeomorphic to \mathbb{T}^5 , where $\mathbb{T} := \mathbb{R}/(2\pi\mathbb{Z})$ is the standard, “flat” torus. Moreover, on $\mathcal{T}_{\omega, \varepsilon, \mu}$ the motions are quasi-periodic, in ORC configuration, with suitable (“diophantine”) irrational frequencies.

Theorem 2.6 extends Theorem 1.1 to ORC motions. As we briefly discuss below, even though the setting is similar, the extension is not completely trivial. Here we provide a sketch of the proof.

In [13] it is shown that $\mathcal{H}_{\text{RPS}_{\pi}^{\mathbb{C}}}$ is related to the Hamiltonian \mathcal{H}_{RPS} in (76) with $n = 2$ by a simple relation. If, in order to avoid confusions, we equip with “tildas” the coordinates (72) with $n = 2$ and denote as

$$\text{RPS}^{\mathbb{C}} := (\mathbf{\Lambda}, \tilde{\boldsymbol{\lambda}}, \tilde{\mathbf{t}}, \tilde{\mathbf{t}}^*, \tilde{T}, \tilde{T}^*) = (\Lambda_1, \Lambda_2, \tilde{\lambda}_1, \tilde{\lambda}_2, \tilde{t}_1, \tilde{t}_2, \tilde{t}_3, \tilde{t}_1^*, \tilde{t}_2^*, \tilde{t}_3^*, \tilde{T}, \tilde{T}^*)$$

their complex version, defined via

$$\begin{aligned} \tilde{t}_1 &:= \frac{\tilde{\eta}_1 - i\tilde{\xi}_1}{\sqrt{2}} & \tilde{t}_2 &:= \frac{\tilde{\eta}_2 - i\tilde{\xi}_2}{\sqrt{2}} & \tilde{t}_3 &:= \frac{\tilde{p} - i\tilde{q}}{\sqrt{2}} & \tilde{T} &:= \frac{\tilde{P} - i\tilde{Q}}{\sqrt{2}} \\ \tilde{t}_1^* &:= \frac{\tilde{\eta}_1 + i\tilde{\xi}_1}{\sqrt{2i}} & \tilde{t}_2^* &:= \frac{\tilde{\eta}_2 + i\tilde{\xi}_2}{\sqrt{2i}} & \tilde{t}_3^* &:= \frac{\tilde{p} + i\tilde{q}}{\sqrt{2i}} & \tilde{T}^* &:= \frac{\tilde{P} + i\tilde{Q}}{\sqrt{2i}} \end{aligned} \quad (108)$$

and, finally, introduce the involution

$$\phi_1^- (\Lambda_1, \Lambda_2, \lambda_1, \lambda_2, t, t^*, T, T^*) := (-\Lambda_1, \Lambda_2, -\lambda_1, \lambda_2, t, t^*, T, T^*). \quad (109)$$

Then we have

Proposition 2.2 ([13]) $\mathcal{H}_{\text{rps}\mathbb{C}} = \mathcal{H}_{\text{rps}\mathbb{C}} \circ \phi_1^-$.

In particular, the coefficients of the expansion

$$f_{\text{RPS}\mathbb{C}}^{\text{av}} = C_0(\mathbf{\Lambda}) + i\mathbf{t}_h \cdot \sigma(\mathbf{\Lambda})\mathbf{t}^* + i\varsigma(\mathbf{\Lambda})t_3t_3^* + O_4(\mathbf{t}, \mathbf{t}^*; \mathbf{\Lambda}) \quad (110)$$

of $f_{\text{RPS}\mathbb{C}}^{\text{av}}$ are obtained from the corresponding coefficients $\tilde{\sigma}(\mathbf{\Lambda})$, $\tilde{\varsigma}(\mathbf{\Lambda})$ computed in [5] by applying the projection on $(\mathbf{\Lambda}, \lambda)$ of the transformation in (109). This immediately provides

$$\begin{cases} \sigma(\Lambda_1, \Lambda_2) = \tilde{\sigma}(-\Lambda_1, \Lambda_2) = \begin{pmatrix} -\frac{s}{\Lambda_1} & -i\frac{\tilde{s}}{\sqrt{\Lambda_1\Lambda_2}} \\ -i\frac{\tilde{s}}{\sqrt{\Lambda_1\Lambda_2}} & \frac{s}{\Lambda_2} \end{pmatrix} \\ \varsigma(\Lambda) = \tilde{\varsigma}(-\Lambda_1, \Lambda_2) = -\left(\frac{1}{\Lambda_2} - \frac{1}{\Lambda_1}\right)s \end{cases} \quad (111)$$

with

$$s := -m_1m_2\frac{\alpha}{2a_1}b_{3/2}^{(1)}(\alpha) \quad \tilde{s} := m_1m_2\frac{\alpha}{2a_1}b_{3/2}^{(2)}(\alpha) \quad \alpha = \frac{a_2}{a_1} \quad (112)$$

where $b_s^{(j)}(\alpha)$'s being the Laplace coefficients⁶. It is to be remarked, from the formulae in (111)–(112) that the matrix σ is symmetric but *not* real. This is a remarkable difference with the prograde case studied in [8, 5], which, in particular, does not ensure “a priori” the reality of its eigenvalues. However, the following turns true:

Lemma 2.3 *The eigenvalues of the (2×2) matrix $\sigma(\mathbf{\Lambda})$ in (110) are real. Hence, $(\mathbf{t}, \mathbf{t}^*) = (\mathbf{0}, \mathbf{0}) \in \mathbb{R}^3 \times \mathbb{R}^3$ is an elliptic equilibrium point for $f_{\text{RPS}\mathbb{C}}^{\text{av}}$.*

Proof The eigenvalues of σ can be explicitly computed:

$$\sigma_1, \sigma_2 = \frac{\text{tr } \sigma}{2} \pm \frac{1}{2}\sqrt{(\text{tr } \sigma)^2 - 4 \det \sigma}. \quad (113)$$

Since $\text{tr } \sigma = \left(\frac{1}{\Lambda_2} - \frac{1}{\Lambda_1}\right)s$ is real, we have to check that the discriminant

$$\Delta := (\text{tr } \sigma)^2 - 4 \det \sigma = \left(\frac{1}{\Lambda_2} - \frac{1}{\Lambda_1}\right)^2 s^2 + \frac{4}{\Lambda_1\Lambda_2}(s^2 - \tilde{s}^2)$$

is positive. Recalling that the Laplace coefficients verify

$$b_s^{(j)}(\beta) > b_s^{(j+1)}(\beta) \quad \text{for all } s > 0, \quad j \in \mathbb{Z}, \quad 0 < |\beta| < 1,$$

(see Ref.[8] for a proof), one has

$$s^2 - \tilde{s}^2 = (m_1m_2\frac{\alpha}{a_1})^2((b_{3/2}^{(1)}(\alpha))^2 - (b_{3/2}^{(2)}(\alpha))^2) > 0. \quad (114)$$

and we have the assertion. \square

The formulae in (111)–(112) show that, as in the prograde case, the eigenvalues of $\sigma(\mathbf{\Lambda})$ and the number $\varsigma(\mathbf{\Lambda})$ verify, identically

$$\sigma_1 + \sigma_2 + \varsigma \equiv 0 \quad (115)$$

⁶The Laplace coefficients defined via the Fourier expansion

$$\frac{1}{(1 - 2\alpha \cos \theta + \alpha^2)^s} = \sum_{k \in \mathbb{Z}} b_s^{(k)}(\alpha) e^{ik\theta} \quad i := \sqrt{-1}.$$

By analogy with the latter identity in (24), we shall refer to (115) as *Herman resonance*. The asymptotic values of the eigenvalues σ_1 , σ_2 and ς in the well-spaced regime (105) can be computed directly from (113)–(114), or from the corresponding ones in [8, 5] applying the transformation (109). In any case, the result is

$$\begin{cases} \sigma_1 = +\frac{3}{4\Lambda_1} \frac{a_2^2}{a_1^3} + O\left(\frac{a_2^3}{a_1^4\Lambda_1}\right) \\ \sigma_2 = -\frac{3}{4\Lambda_2} \frac{a_2^2}{a_1^3} + O\left(\frac{a_2^3}{a_1^4\Lambda_2}\right) \\ \varsigma = \frac{3}{4} \frac{a_2^2}{a_1^2} \left(\frac{1}{\Lambda_2} - \frac{1}{\Lambda_1} \right) + O\left(\frac{a_2^3}{a_1^4\Lambda_2}\right) \end{cases}$$

It shows that there is no other resonance besides Herman resonance in (115), provided the semi-axes are well spaced. Recall the definition of \mathcal{L} in (105).

Lemma 2.4 *For any $K > 0$, there exist Λ_{\pm} , α_{\pm} such that the triple $\Omega^{\mathbb{C}}(\mathbf{\Lambda}) := (\sigma_1(\mathbf{\Lambda}), \sigma_2(\mathbf{\Lambda}), \varsigma(\mathbf{\Lambda}))$ verifies*

$$\Omega^{\mathbb{C}}(\mathbf{\Lambda}) \cdot k \neq 0 \quad \forall k \in \mathbb{Z}^3, \quad 0 < |k| \leq K, \quad k \neq N(1, 1, 1) \quad \forall \mathbf{\Lambda} \in \mathcal{L} \quad (116)$$

with some $N \in \mathbb{Z}$.

At first sight, Lemma 2.4 might seem an obstruction towards the construction of the Birkhoff normal form for the Hamiltonian (101). However, as in the prograde case, the conservation of the angular momentum length

$$C = \Lambda_2 - \Lambda_1 - \mathbf{it} \cdot \mathbf{t}^* \quad (117)$$

is of great help. Indeed, by the commutation of $f_{\text{RPS}\frac{\mathbb{C}}{\pi}}$ and C , it turns out that, in the Taylor expansion (110), only monomials with literal part $\mathbf{t}^{\mathbf{a}}\mathbf{t}^{*\mathbf{a}^*}$ verifying

$$\sum_i a_i = \sum_i a_i^* \quad (118)$$

appear. In [4] it is shown that, because of (118), then (116) is sufficient for constructing a Birkhoff normal form (i.e., Theorem 2.1 with $n = 2$) for the Hamiltonian (101). Moreover, the torsion matrix (i.e., the matrix $\tau(\mathbf{\Lambda})$ defined via (80)) for this case can be computed from the analogue one from the prograde case again applying (109) to the torsion of the prograde problem. The computation is omitted (see [13] for the details), apart for stating that it is non-singular. An application of Theorem 2.2 then leads to the proof of Theorem 2.6.

Proof of 2. As a second set of coordinates, we use the \mathcal{P} -coordinates defined in Section 1.6. In the case $n = 2$, they reduce to

$$\mathcal{P} = (Z, C, \mathbf{\Theta}, \mathbf{\Lambda}, \zeta, \kappa_2 \mathbf{\vartheta}, \ell)$$

with

$$\mathbf{\Lambda} = (\Lambda_1, \Lambda_2), \quad \mathbf{\Theta} = (\Theta_1, \Theta_2), \quad \ell = (\ell_1, \ell_2), \quad \mathbf{\vartheta} = (\vartheta_1, \vartheta_2)$$

We denote as

$$\mathcal{H}_{\mathcal{P}} = -\sum_{j=1}^2 \frac{\mu_j^3 M_j^2}{2\Lambda_j^2} + \mu f_{\mathcal{P}}(\mathbf{\Lambda}, \mathbf{\Theta}, \ell, \mathbf{\vartheta}; C)$$

the four–degrees–of–freedom Hamiltonian (96) written using \mathcal{P} –coordinates, which is independent of Z , ζ and κ_2 .

The manifolds

$$\mathcal{D}_u^0 := \left\{ (\mathbf{\Lambda}, \Theta, \ell, \vartheta; C) : (\Theta_2, \vartheta_2) = (0, 0) \right\} \quad (119)$$

corresponds to retrograde motions. It is invariant as $f_{\mathcal{P}}$ has an equilibrium on it and includes, in particular, the manifold \mathcal{M}_π in (100).

We establish a suitable domain (including \mathcal{D}_u^0) for the coordinates \mathcal{P} where $\mathcal{H}_{\mathcal{P}}$ is regular. We check below that the following domain is suited to the scope:

$$\mathcal{D}_{\mathcal{P}}(C) := \left\{ (\mathbf{\Lambda}, \Theta_1) \in \mathcal{A}(C) \right\} \times \left\{ (\ell, \vartheta_1) \in \mathbb{T}^3 \right\} \times \left\{ (\Theta_2, \vartheta_2) \in \mathcal{B}(\Theta_1, C) \right\} \quad (120)$$

where

$$\begin{aligned} \mathcal{A}(C) &:= \left\{ (\Lambda_1, \Lambda_2, \Theta_1) : (\Lambda_1, \Lambda_2) \in \mathcal{L}(C), \Theta_1 \in \mathcal{G}(\Lambda_1, \Lambda_2, C) \right\} \\ \mathcal{B}(\Theta_1, C) &:= \left\{ (\Theta_2, \vartheta_2) : |\Theta_2| < \frac{1}{2} \min\{C, \Theta_1\}, |\vartheta_2| < \frac{\pi}{2} \right\} \\ \mathcal{L}(C) &:= \left\{ \mathbf{\Lambda} : \mathbf{\Lambda} \in \mathcal{L}, \quad \Lambda_2 > C + \frac{2}{c} \sqrt{\alpha_+} \Lambda_1 \right\} \\ \mathcal{G}(\Lambda_1, \Lambda_2, C) &:= (C_-, C_+), \quad C_- := \frac{2}{c} \sqrt{\alpha_+} \Lambda_1, \quad C_+ := \min\{\Lambda_2 - C, \Lambda_1\}. \end{aligned} \quad (121)$$

with \mathcal{L} is as in (105), while c is an arbitrarily fixed number in $(0, 1)$. We need to establish two kinds of conditions.

a) existence of the perihelia We need that the planets' eccentricities e_1, e_2 stay strictly confined in $(0, 1)$. Namely, that the following inequalities are satisfied:

$$0 < \Theta_1 < \Lambda_1, \quad 0 < C_2 < \Lambda_2 \quad (122)$$

with $C_2 := |\mathbf{C}_2|$, \mathbf{C}_2 as in (25). The expression of C_2 using \mathcal{P} is

$$C_2 = \sqrt{C^2 + \Theta_1^2 - 2\Theta_2^2 + 2\sqrt{(C^2 - \Theta_2^2)(\Theta_1^2 - \Theta_2^2)} \cos \vartheta_2}$$

We observe that C_2 may vanish only for $(\Theta_2, \vartheta_2) = (0, \pi)$. Since we deal with the equilibrium (119), the occurrence of this equality is automatically excluded, limiting the values of the coordinates (Θ_2, ϑ_2) in the set \mathcal{B} in (121) since in this case

$$C_2^2 \geq \frac{3}{4} C^2. \quad (123)$$

Moreover, the two right inequalities in (122) are satisfied taking

$$\Theta_1 < \min\{\Lambda_2 - C, \Lambda_1\} = C_+$$

where we have used the triangular inequality $C_2 = |\mathbf{C} - \mathbf{C}_1| \leq |\mathbf{C}| + |\mathbf{C}_1| = C + \Theta_1$.

b) non–collision conditions We have to exclude possible encounters of the planets with the sun and each other. Collisions of the inner planet with the sun are excluded by (121). Indeed, using (123),

$$1 - e_2^2 = \frac{C_2^2}{\Lambda_2^2} \geq \frac{3}{4} \frac{C^2}{\Lambda_2^2}$$

whence the minimum distance of the inner planet with the sun $a_2(1 - e_2)$ is positive. In order to avoid planetary collisions, it is typical to ensure the following inequality:

$$a_2(1 + e_2) < c^2 a_1(1 - e_1)$$

with $0 < c < 1$. A sufficient condition for it is

$$\Theta_1 \geq \frac{2}{c} \sqrt{\alpha_+} \Lambda_1 = C_-.$$

Indeed, if this inequality is satisfied, one has

$$a_2(1 + e_2) < 2a_2 < \frac{a_1}{2} \frac{\Theta_1^2 c^2}{\Lambda_1^2} = \frac{a_1}{2} (1 - e_1^2) c^2 < a_1(1 - e_1) c^2.$$

The hyperbolic equilibrium [13] By the formulae (89)–(90) with $n = 2$, the ℓ of $\mathcal{H}_{\mathcal{P}}$ is given by

$$\bar{\mathcal{H}}_{\mathcal{P}} = - \sum_{j=1}^2 \frac{\mu_j^3 M_j^2}{2\Lambda_j^2} + \mu \left(-\frac{m_1 m_2}{a_1} + \overline{f_{\mathcal{P}}^{12}}^{(2)} \right) + \frac{\mu}{a_1} \mathcal{O} \left(\frac{a_2^2}{a_1^2} \right)$$

with

$$\begin{aligned} \overline{f_{\mathcal{P}}^{12}}^{(2)} &= m_1 m_2 \frac{a_2^2}{4a_1^3} \frac{\Lambda_1^3}{\Theta_1^5} \left[\frac{5}{2} (3\Theta_2^2 - \Theta_1^2) \right. \\ &\quad - \frac{3}{2} \frac{4\Theta_2^2 - \Theta_1^2}{\Lambda_2^2} \left(C^2 + \Theta_1^2 - 2\Theta_2^2 + 2\sqrt{(C^2 - \Theta_2^2)(\Theta_1^2 - \Theta_2^2)} \cos \vartheta_2 \right) \\ &\quad \left. + \frac{3}{2} \frac{(\Theta_1^2 - \Theta_2^2)(C^2 - \Theta_2^2)}{\Lambda_2^2} \sin^2 \vartheta_2 \right]. \end{aligned}$$

We shall now prove that, restricting the domain (120) a little bit, so that the manifolds (119) are hyperbolic for $\overline{f_{\mathcal{P}}^{12}}^{(2)}$. We fix the following domain

$$\mathcal{D}_u := \mathcal{A}_u \times \mathcal{B}_u \times \mathbb{T}^3 \quad (124)$$

with

$$\begin{aligned} \mathcal{A}_u(C) &:= \left\{ (\Lambda_1, \Lambda_2) \in \mathcal{L}_u(C), \quad \Theta_1 \in \mathcal{G}_u(\Lambda_1, \Lambda_2, C) \right\} \\ \mathcal{B}_u(C) &:= \left\{ (\Theta_2, \vartheta_2) : |\Theta_2| < \frac{C}{2}, |\vartheta_2| < \frac{\pi}{2} \right\} \end{aligned} \quad (125)$$

where

$$\begin{aligned} \mathcal{L}_u(C) &:= \left\{ \Lambda = (\Lambda_1, \Lambda_2) \in \mathcal{L} : 5\Lambda_2^2 C - (C + \frac{2}{c} \sqrt{\alpha_+} \Lambda_2)^2 (4C + \frac{2}{c} \sqrt{\alpha_+} \Lambda_2) > 0, \right. \\ &\quad \left. \Lambda_1 > C, \Lambda_2 > \max \left\{ C + \frac{2}{c} \sqrt{\alpha_+} \Lambda_1, 2C \right\} \right\} \\ \mathcal{G}_u(\Lambda_1, \Lambda_2, C) &:= (\bar{C}_-, \bar{C}_+) \end{aligned} \quad (126)$$

where \mathcal{L} is as in (105) and, if $C^*(\Lambda_2, C)$ is the unique positive root of the cubic polynomial $C_2 \rightarrow 5\Lambda_2^2 C - (C + C_2)^2 (4C + C_2)$, then

$$\bar{C}_- := \max \left\{ \frac{2}{c} \sqrt{\alpha_+} \Lambda_1, C \right\} \quad \bar{C}_+ := \min \{ \Lambda_1, C^* \}. \quad (127)$$

Implicitly, we shall prove that

$$\bar{C}_- < \bar{C}_+ . \quad (128)$$

We check that the coefficients in front of $\Theta_2^2, \vartheta_2^2$ in the Taylor expansion about $(\Theta_2, \vartheta_2) = (0, 0)$ have opposite sign in the domain (124), so that the equilibrium manifold (119) is hyperbolic. Indeed, the part of degree 2 in such expansion is

$$m_1 m_2 \frac{a_2^2}{a_1^3} \frac{1}{8} \frac{\Lambda_1^3}{\Lambda_2^2 \Theta_1^5} \times \left[\frac{3A}{C} \Theta_2^2 + 3C \Theta_1^2 B \vartheta_2^2 + O(\Theta_2^4 + \vartheta_2^4) \right]$$

where

$$A := 5\Lambda_2^2 C - (C + \Theta_1)^2 (4C + \Theta_1) \quad \text{and} \quad B := C - \Theta_1 . \quad (129)$$

Both $\Theta_1 \rightarrow a(\Lambda_1, \Theta_1; C)$ and $\Theta_1 \rightarrow b(\Theta_1; C)$, as functions of Θ_1 decrease monotonically from a positive value (respectively, $C(5\Lambda_2^2 - 4C^2)$ and C) to $-\infty$ as Θ_1 increases from $\Theta_1 = 0$ to $\Theta_1 = +\infty$. The function $a(\Lambda_1, \Theta_1; C)$ changes its sign for Θ_1 equal to a suitable unique positive value $C^*(\Lambda_2, C)$, while $b(\Theta_1; C)$ does it for $\Theta_1 = C$. We note that (i) inequality $C < \min\{C_+, C^*\}$ follows immediately from the assumptions (126) (in particular, the two last ones) and (ii), more generally, that $C^* \leq C$ is equivalent to $\Lambda_2 \leq 2C$. Since, for our purposes, we have to exclude $C^* = C$ (otherwise, $a(\Lambda_1, \Theta_1; C)$ and $b(\Theta_1; C)$ would be simultaneously positive and simultaneously negative, and no hyperbolicity would be possible), we distinguish two cases.

- (a) $C > \frac{2}{c} \sqrt{\alpha_+} \Lambda_1$ and $C + \frac{2}{c} \sqrt{\alpha_+} \Lambda_1 < \Lambda_2 < 2C$. In this case $C^* < C$. We show that no such \mathcal{G}_u can exist in this case. In fact, since $C^* < C$, in order that the interval (C^*, C) and the set \mathcal{G} have a non-empty intersection, one should have, necessarily, $C_+ = \sup \mathcal{G} > C^*$, hence, in particular, $\Lambda_2 - C > C^*$. Using the definition of C^* , this would imply $\Lambda_2 > 2C$, which is a contradiction.
- (b) $\Lambda_2 > \max\{2C, C + \frac{2}{c} \sqrt{\alpha_+} \Lambda_1\}$. In this case $C < C^* < \Lambda_2 - C$. In order that the interval (C, C^*) and the set \mathcal{G} have a non-empty intersection, we need

$$C_- < C^* \quad \text{and} \quad C_+ > C \quad (130)$$

and such intersection will be given by the interval \mathcal{G}_u as in (126). Note that the definition of \bar{C}_+ does not include $\Lambda_2 - C$ in the brackets because, as noted, $C^* < \Lambda_2 - C$. But (130) are equivalent to (126).

Proof of 3. Here we prove that

Theorem 2.7 *Let $\alpha_+ < \frac{1}{16}$. There exist universal numbers $1 < \underline{k} < \bar{k}$ such that, if*

$$\alpha_- < \frac{\underline{k}^2}{\bar{k}^2} \alpha_+ , \quad \frac{\bar{k}}{\sqrt{\alpha_+}} < \frac{\mu_2}{\mu_1} \sqrt{\frac{M_2}{M_1}} < \frac{\underline{k}}{\sqrt{\alpha_-}}$$

then $\mathcal{D}_s \cap \mathcal{D}_u^0$ is non-empty. The following values work:

$$\underline{k} = \frac{1}{4} \sqrt{\frac{3}{10} (69 + 11\sqrt{33})} \sim 1.57 , \quad \bar{k} = 2 . \quad (131)$$

Proof The sets \mathcal{D}_s in (106) and \mathcal{D}_u^0 in (119) are expressed with different sets of coordinates. To prove that \mathcal{D}_s and \mathcal{D}_u^0 have a non-empty intersection, we need to use the same set for both. We choose to use the coordinates \mathcal{P} , so we rewrite \mathcal{D}_s in terms of \mathcal{P} .

Using \mathcal{P} , the set \mathcal{D}_s becomes (at the expenses of diminishing ε , if necessary)

$$\mathcal{D}_s = \mathcal{A}_s \times \mathcal{B}_s \times \mathbb{T}^3$$




Figure 13: The blue curve is \mathcal{C} ; the orange line has slope \underline{k} , the green one has slope \bar{k} (MATHEMATICA).




Figure 14: The blue strip corresponds to the set \mathcal{L}_1 , the green one to \mathcal{L}_2 (MATHEMATICA).




Figure 15: \mathcal{L}_1 : the blue region; \mathcal{L}_2 : the green region; \mathcal{L}_3 : the violet region (MATHEMATICA).

where, if

$$\mathcal{L}_s(C) := \left\{ \Lambda = (\Lambda_1, \Lambda_2) \in \mathcal{L}_0 : |\Lambda_2 - \Lambda_1 - C| < \varepsilon \right\}, \quad \mathcal{G}_s(\Lambda_1) := \left\{ \Theta_1 : 0 < \Lambda_1 - \Theta_1 < \varepsilon \right\}, \quad (132)$$

then

$$\mathcal{A}_s := \left\{ (\Lambda_1, \Lambda_2, \Theta_1) : (\Lambda_1, \Lambda_2) \in \mathcal{L}_s, \Theta_1 \in \mathcal{G}_s(\Lambda_1) \right\}, \quad \mathcal{B}_s := \left\{ (\Theta_2, \vartheta_2) : |(\Theta_2, \vartheta_2)| < \varepsilon \right\}. \quad (133)$$

All we have to do is to check that the intersection $\mathcal{A}_s \cap \mathcal{A}_u$ is non-empty.

Recalling the definition of \mathcal{A}_u in (125)–(126) and the definition of \mathcal{A}_s in (132)–(133), asserting that $\mathcal{A}_s \cap \mathcal{A}_u \neq \emptyset$ is equivalent to asserting that

$$\mathcal{L}_s(C) \cap \mathcal{L}_u(C) \neq \emptyset$$

and

$$\mathcal{G}_s(\Lambda_1) \cap \mathcal{G}_u(\Lambda_1, \Lambda_2, C) \neq \emptyset \quad \forall (\Lambda_1, \Lambda_2) \in \mathcal{L}_s(C) \cap \mathcal{L}_u(C).$$

It will be enough to check that

$$\mathcal{L}_s(C) \cap \mathcal{L}_u(C) \cap \mathcal{L}_{su}(C) \neq \emptyset \quad (134)$$

and

$$\mathcal{G}_s(\Lambda_1) \cap \mathcal{G}_u(\Lambda_1, \Lambda_2, C) \neq \emptyset \quad \forall (\Lambda_1, \Lambda_2) \in \mathcal{L}_s(C) \cap \mathcal{L}_u(C) \cap \mathcal{L}_{su}(C), \quad (135)$$

where, if \overline{C}_\pm are as in (127), \mathcal{L}_{su} is defined as

$$\mathcal{L}_{su} := \left\{ (\Lambda_1, \Lambda_2) : \overline{C}_+ = \Lambda_1 \right\}. \quad (136)$$

Note that (135) is certainly satisfied provided (134) is, since in fact, for $(\Lambda_1, \Lambda_2) \in \mathcal{L}_s(C) \cap \mathcal{L}_u(C) \cap \mathcal{L}_{su}(C)$,

$$\mathcal{G}_s(\Lambda_1) \cap \mathcal{G}_u(\Lambda_1, \Lambda_2, C) = \left\{ \Theta_1 : \max\{\overline{C}_-, \Lambda_1 - \varepsilon\} < \Theta_1 < \Lambda_1 \right\}$$

which is well-defined by (127)–(128).

On the other hand, in view of the definition of \overline{C}_+ in (127), and of C^* a few lines above, \mathcal{L}_{su} in (136) is equivalently defined as

$$\mathcal{L}_{su} = \left\{ (\Lambda_1, \Lambda_2) : 5\Lambda_2^2 C - (C + \Lambda_1)^2(4C + \Lambda_1) > 0 \right\}. \quad (137)$$

Therefore, in view of this definition and the definitions of \mathcal{L}_s , \mathcal{L}_u in (126) and (132), one sees that the set on the left hand side in (134) is determined by inequalities

$$\begin{aligned} \Lambda_- &< \Lambda_1 < \Lambda_+ \\ k_- \Lambda_1 &\leq \Lambda_2 \leq k_+ \Lambda_1 \\ 5\Lambda_2^2 C - (C + 2\sqrt{\alpha_+} \Lambda_2)^2(4C + 2\sqrt{\alpha_+} \Lambda_2) &> 0 \\ \Lambda_1 &> C \\ \Lambda_2 &> \max\{C + 2\sqrt{\alpha_+} \Lambda_1, 2C\} \\ |\Lambda_2 - \Lambda_1 - C| &< \varepsilon \\ 5\Lambda_2^2 C - (C + \Lambda_1)^2(4C + \Lambda_1) &> 0 \end{aligned} \quad (138)$$

We observe that no phase point⁷ (Λ_1, Λ_2) with $\Lambda_2 - \Lambda_1 - C < 0$ will ever satisfy (138), and that inequality $\Lambda_2 > 2C$ is implied by $\Lambda_1 > C$ and (137). Then, we divide such inequalities in three groups, so as to rewrite the set (134) as the intersection of the sets

⁷Inequalities $\Lambda_1 < C_*$ (which is equivalent to (137)) and $C_* < \Lambda_2 - C$ (which is equivalent to $\Theta_1 > C$, in turn implied by the definition of \mathcal{G}_{su} above) imply $\Lambda_2 - \Lambda_1 - C > 0$.

$$\begin{aligned}
\widehat{\mathcal{L}}_1 &:= \left\{ (\Lambda_1, \Lambda_2) : \Lambda_- < \Lambda_1 < \Lambda_+, \Lambda_1 > C, \Lambda_2 > 2C, \right. \\
&\quad \left. \max\{k_- \Lambda_1, (C + \Lambda_1) \sqrt{\frac{4C + \Lambda_1}{5C}}\} < \Lambda_2 \leq k_+ \Lambda_1 \right\} \\
\widehat{\mathcal{L}}_2 &:= \left\{ (\Lambda_1, \Lambda_2) : 0 < \Lambda_2 - \Lambda_1 - C < \varepsilon, \Lambda_2 > C + 2\sqrt{\alpha_+} \Lambda_1, \Lambda_1 > C \right\} \\
\widehat{\mathcal{L}}_3 &:= \left\{ (\Lambda_1, \Lambda_2) : 5\Lambda_2^2 C - (C + 2\sqrt{\alpha_+} \Lambda_2)^2 (4C + 2\sqrt{\alpha_+} \Lambda_2) > 0, \Lambda_2 > 2C \right\}
\end{aligned}$$

We now aim to choose the parameters Λ_{\pm} , k_{\pm} and α_+ so as to find a non-empty intersection of the sets above.

Let us denote as \mathcal{C} the curve, in the (Λ_2, Λ_1) -plane, having equation

$$\mathcal{C} : \quad \Lambda_2 = (C + \Lambda_1) \sqrt{\frac{4C + \Lambda_1}{5C}} \quad (139)$$

Let

$$\Lambda_1 = k \Lambda_2$$

be any straight line through the origin. The straight line intersecting \mathcal{C} into the point $(\underline{\Lambda}_1, \underline{\Lambda}_2) = (C, 2C)$ has $\bar{k} = 2$, and intersects this curve, also in the higher point

$$(\overline{\Lambda}_1, \overline{\Lambda}_2) = \left(\frac{1}{2} (13 + \sqrt{185}), (13 + \sqrt{185}) \right) C .$$

Any other line with $k > \bar{k}$ has a lower intersection $(\underline{\Lambda}'_1, \underline{\Lambda}'_2)$, with $\underline{\Lambda}'_1 < C$ and $\underline{\Lambda}'_2 < 2C$ and a higher intersection $(\overline{\Lambda}'_1, \overline{\Lambda}'_2)$ with $\overline{\Lambda}'_1 > \overline{\Lambda}_1$ and $\overline{\Lambda}'_2 > \overline{\Lambda}_2$.

The last straight line, in the plane (Λ_1, Λ_2) , through the origin intersecting \mathcal{C} is the tangent line, and it is easy to compute (see below) that such a tangent line has slope \bar{k} as in (131) (Figure 13). We then conclude that, as soon as we choose $k_- < \bar{k}$, $k_+ > \bar{k}$, $\Lambda_- < \underline{\Lambda}_1$, $\Lambda_+ > \overline{\Lambda}_1$, we have the inclusion

$$\widehat{\mathcal{L}}_1 \supset \mathcal{L}_1 := \left\{ (\Lambda_1, \Lambda_2) : (C + \Lambda_1) \sqrt{\frac{4C + \Lambda_1}{5C}} < \Lambda_2 \leq 2\Lambda_1 \right\} .$$

Let us now turn to $\widehat{\mathcal{L}}_2$. Since we are assuming $\alpha_+ < \frac{1}{16}$, we conclude that the strip

$$\mathcal{L}_2 := \left\{ (\Lambda_1, \Lambda_2) : 0 < \Lambda_2 - \Lambda_1 - C < \varepsilon, \Lambda_1 > C \right\}$$

is all included in the region

$$\widetilde{\mathcal{L}}_2 = \left\{ (\Lambda_1, \Lambda_2) : \Lambda_2 > C + 2\sqrt{\alpha_+} \Lambda_1, \Lambda_1 > C \right\}$$

and this allows to conclude

$$\widehat{\mathcal{L}}_2 = \mathcal{L}_2 \cap \widetilde{\mathcal{L}}_2 = \mathcal{L}_2 .$$

Since the sets \mathcal{L}_1 and \mathcal{L}_2 have a non-empty intersection, independently of α_+ (see Figure 14), a fortiori, $\widehat{\mathcal{L}}_1$ and $\widehat{\mathcal{L}}_2$ have one:

$$\widehat{\mathcal{L}}_1 \cap \widehat{\mathcal{L}}_2 \supset \mathcal{L}_1 \cap \mathcal{L}_2 \neq \emptyset .$$

Observe, in particular, that $\mathcal{L}_1 \cap \mathcal{L}_2$ (hence, $\widehat{\mathcal{L}}_1 \cap \widehat{\mathcal{L}}_2$) has non-empty intersection with any strip $\mathbb{R} \times [2C, y]$, with $y > 2C$ (see Figure 15).

On the other hand, it is immediate to check that $\widehat{\mathcal{L}}_3$ includes the horizontal strip

$$\mathcal{L}_3 := \left\{ (\Lambda_1, \Lambda_2) : 2C < \Lambda_2 < \frac{C}{2\sqrt{\alpha_+}}, \Lambda_1 \in \mathbb{R} \right\} \quad 0 < \alpha_+ < \frac{1}{16}$$

and so we conclude

$$\mathcal{L}_s(C) \cap \mathcal{L}_u(C) \cap \mathcal{L}_{su}(C) = \widehat{\mathcal{L}}_1 \cap \widehat{\mathcal{L}}_2 \cap \widehat{\mathcal{L}}_3 \supset \mathcal{L}_1 \cap \mathcal{L}_2 \cap \mathcal{L}_3 \neq \emptyset$$

In order to complete the proof, it remains to prove that the tangent straight line to \mathcal{C} through the origin has slope \underline{k} as in (131).

We switch to the homogenized variables

$$x := \frac{\Lambda_1}{C} \quad y = \frac{\Lambda_2}{C}$$

so that the curve \mathcal{C} in (139) becomes

$$\widehat{\mathcal{C}}: \quad y = (1+x)\sqrt{\frac{4+x}{5}}.$$

We look for a straight line through the origin $y = \underline{k}x$ with $\underline{k} > 0$ which is tangent to $\widehat{\mathcal{C}}$ at some point (a, b) , with $a > 0$.

The intersections between $\widehat{\mathcal{C}}$ and any straight line through the origin $y = kx$ are ruled by a complete cubic equation, given by

$$x^3 + (6 - 5k^2)x^2 + 9x + 4 = 0. \quad (140)$$

In order that such an equation has a double solution $x = a$ for $k = \underline{k}$, one needs that, when $k = \underline{k}$, it can factorized as

$$(x - a)^2(x - c) = 0 \quad (141)$$

Therefore, equating the respective coefficients of (140) and (141) one finds the equations

$$\begin{cases} -(c + 2a) = 6 - 5\underline{k}^2 \\ 2ac + a^2 = 9 \\ -a^2c = 4 \end{cases}$$

Two last equations, allow to eliminate b so as to obtain the equation for a

$$a^3 - 9a - 8 = 0$$

which has the following three roots:

$$a_0 = -1, \quad a_{\pm} = \frac{1 \pm \sqrt{33}}{2}.$$

The only admissible (positive) value is then

$$a = a_+ = \frac{1 + \sqrt{33}}{2}$$

and it provides the values

$$c = \frac{-17 + \sqrt{33}}{32}, \quad \underline{k} = \frac{1}{4}\sqrt{\frac{3}{10}(69 + 11\sqrt{33})}. \quad \square$$

A The m_0 -centric reduction

The Hamiltonian of $1 + n$ masses m_0, \dots, m_n interacting through gravity is

$$\mathcal{H} = \sum_{i=0}^n \frac{|\mathbf{u}_i|^2}{2m_i} - \sum_{0 \leq i < j \leq n} \frac{m_i m_j}{|\mathbf{v}_i - \mathbf{v}_j|}. \quad (142)$$

We switch from the position coordinates \mathbf{v}_i to new ones, denoted \mathbf{x}_i , where \mathbf{x}_0 is the coordinate of m_0 , while \mathbf{x}_i is the coordinate of m_i relatively to m_0 . The change is

$$\mathbf{v}_i = \begin{cases} \mathbf{x}_0 & i = 0 \\ \mathbf{x}_i + \mathbf{x}_0 & i = 1, \dots, n. \end{cases} \quad (143)$$

As the change does not involve the \mathbf{u}_i 's, the coordinates \mathbf{y}_i conjugated to \mathbf{x}_i may be computed imposing the conservation of the standard 1-form

$$\Lambda = \sum_{i=0}^n \mathbf{y}_i \cdot d\mathbf{x}_i = \sum_{i=0}^n \mathbf{u}_i \cdot d\mathbf{v}_i.$$

We find

$$\begin{aligned} \sum_{i=0}^n \mathbf{u}_i \cdot \mathbf{v}_i &= \mathbf{u}_0 \cdot \mathbf{x}_0 + \sum_{i=1}^n \mathbf{u}_i \cdot (\mathbf{x}_i + \mathbf{x}_0) \\ &= \left(\sum_{i=0}^n \mathbf{u}_i \right) \cdot \mathbf{x}_0 + \sum_{i=1}^n \mathbf{u}_i \cdot \mathbf{x}_i. \end{aligned}$$

So we identify

$$\mathbf{y}_i = \begin{cases} \sum_{i=0}^n \mathbf{u}_i & i = 0 \\ \mathbf{u}_i & i = 1, \dots, n. \end{cases}$$

We recognize that \mathbf{y}_0 is the total linear momentum, which keeps constant along the motions of \mathcal{H} , as \mathcal{H} is translation-invariant. Fixing a reference frame moving with the the centre of mass of m_0, \dots, m_n , we have $\mathbf{y}_0 = 0$ and hence

$$\mathbf{u}_i = \begin{cases} -\sum_{i=1}^n \mathbf{y}_i & i = 0 \\ \mathbf{y}_i & i = 1, \dots, n. \end{cases} \quad (144)$$

Replacing (144) and (143) into (142) we arrive at (1), with μ_i, M_i as in (2).

References

- [1] V.I. Arnold. Small denominators and problems of stability of motion in classical and celestial mechanics. *Russian Math. Surveys*, 18(6):85–191, 1963.
- [2] L. Chierchia and G. Pinzari. Properly–degenerate KAM theory (following V.I. Arnold). *Discrete Contin. Dyn. Syst. Ser. S*, 3(4):545–578, 2010.
- [3] L. Chierchia and G. Pinzari. Deprit’s reduction of the nodes revised. *Celestial Mech.*, 109(3):285–301, 2011.
- [4] L. Chierchia and G. Pinzari. Planetary Birkhoff normal forms. *J. Mod. Dyn.*, 5(4):623–664, 2011.
- [5] L. Chierchia and G. Pinzari. The planetary N -body problem: symplectic foliation, reductions and invariant tori. *Invent. Math.*, 186(1):1–77, 2011.
- [6] L. Chierchia and G. Pinzari. Metric stability of the planetary n -body problem. *Proceedings of the International Congress of Mathematicians*, 2014.
- [7] A. Deprit. Elimination of the nodes in problems of n bodies. *Celestial Mech.*, 30(2):181–195, 1983.
- [8] J. Féjoz. Démonstration du ‘théorème d’Arnold’ sur la stabilité du système planétaire (d’après Herman). *Ergodic Theory Dynam. Systems*, 24(5):1521–1582, 2004.
- [9] R. S. Harrington. The stellar three-body problem. *Celestial Mech. and Dyn. Astronom.*, 1(2):200–209, 1969.
- [10] H. Hofer, E. Zehnder. *Symplectic Invariants and Hamiltonian Dynamics*. Birkhäuser Verlag, Basel, 1994.
- [11] G. Pinzari. *On the Kolmogorov set for many–body problems*. PhD thesis, Università Roma Tre, April 2009.
- [12] G. Pinzari. Aspects of the planetary Birkhoff normal form. *Regul. Chaotic Dyn.*, 18(6):860–906, 2013.
- [13] G. Pinzari. On the co-existence of maximal and whiskered tori in the planetary three-body problem. *J. Math. Phys.*, 59(5):052701, 37, 2018.
- [14] G. Pinzari. Perihelia reduction and global Kolmogorov tori in the planetary problem. *Mem. Amer. Math. Soc.*, 255(1218), 2018.
- [15] R. Radau. Sur une transformation des équations différentielles de la dynamique. *Ann. Sci. Ec. Norm. Sup.*, 5:311–375, 1868.